arXiv:1410.0080v1 [hep-th] 1 Oct 2014

Higher Spin Currents in Orthogonal Wolf Space

Changhyun Ahn and **Jinsub Paeng**

Department of Physics, Kyungpook National University, Taegu 702-701, Korea

ahn@knu.ac.kr, jdp2r@knu.ac.kr

Abstract

For the $\mathcal{N} = 4$ superconformal coset theory by $\frac{SO(N+4)}{SO(N)\times SU(2)} \times U(1)$ (that contains an orthogonal Wolf space) with $N = 4$, the $\mathcal{N} = 2$ WZW affine current algebra is obtained. The 16 generators (or 11 generators) of the large $\mathcal{N} = 4$ linear (or nonlinear) superconformal algebra are described by these WZW affine currents explicitly. Along the line of large $\mathcal{N} = 4$ holography, the extra 16 currents with spins $(2, \frac{5}{2}, \frac{5}{2}, 3)$, $(\frac{5}{2}, 3, 3, \frac{7}{2})$, $(\frac{5}{2}, 3, 3, \frac{7}{2})$, and $(3, \frac{7}{2}, \frac{7}{2}, 4)$ are obtained in terms of the WZW affine currents. The lowest spin of this $\mathcal{N} = 4$ multiplet is two rather than one which is for an unitary Wolf space. The operator product expansions (OPEs) between the above 11 currents and these extra 16 higher spin currents are found explicitly.

1 Introduction

It is known in [1] that the large $\mathcal{N} = 4$ higher spin theory on AdS_3 based on the higher spin algebra is dual to the 't Hooft limit of the two dimensional large $\mathcal{N} = 4$ superconformal coset theory. This is a supersymmetric version of the original nonsupersymmetric duality described in [2, 3]. The $\mathcal{N} = 4$ higher spin algebra contains eight fields of spin $s = \frac{3}{2}, 2, \frac{5}{2}, \cdots$ and seven fields of spin $s = 1$. The lowest spin for the fields is given by one. There is no field of spin $\frac{1}{2}$. The exceptional superalgebra $D(2, 1|\frac{\mu}{1-\mu})$ is the largest finite dimensional subalgebra of this $\mathcal{N} = 4$ higher spin algebra. On the other hand, the $\mathcal{N} = 4$ coset theory [4] is described by the Wolf [5, 6, 7] space 'bosonic' coset $\frac{G}{H} = \frac{SU(N+M)_k \times SO(2NM)_1}{SU(N)_{k+M} \times U(1)_{NM(N+M)(N+M+k)}}$ with $M = 2$ [1]. The first level k is the one for $SU(2)_+$ affine algebra while the second level for other $SU(2)_-$ affine algebra is given by N of the large $\mathcal{N} = 4$ nonlinear superconformal algebra. The 't Hooft coupling constant $\lambda = \frac{N+1}{N+k+2}$ is identified with the free parameter of the large $\mathcal{N} = 4$ nonlinear superconformal algebra. In other words, the numerator of λ is given by the second level N plus one and the denominator of λ is given by the sum of first and second levels $(k + N)$ plus two. Furthermore, this 't Hooft parameter λ in two dimensional CFT is equal to the above μ parameter, in the subalgebra of the $\mathcal{N} = 4$ higher spin algebra, which appears in the mass of scalar field in the AdS_3 bulk theory.

For the $N = 3$ (and $M = 2$), the $\mathcal{N} = 2$ WZW affine current algebra [17, 18, 19] was obtained in [20]. The 16 generators (or 11 generators) of the large $\mathcal{N} = 4$ linear [21, 22, 23, 24, 25, 26, 27] (nonlinear [28, 29, 30, 31]) superconformal algebra were obtained. Based on the large $\mathcal{N} = 4$ holography [1], the extra 16 higher spin currents, with spin contents $(1, \frac{3}{2}, \frac{3}{2}, 2)$, $(\frac{3}{2}, 2, 2, \frac{5}{2})$, $(\frac{3}{2}, 2, 2, \frac{5}{2})$, $(2, \frac{5}{2}, \frac{5}{2}, 3)$ described in terms of four $\mathcal{N} = 2$ multiplets, were found and realized by the above $\mathcal{N} = 2$ WZW affine currents. Then the operator product expansions (OPEs) between the above 11 currents and these extra 16 higher spin currents were described explicitly. By calculating the OPEs between the above 16 higher spin currents, the next 16 higher spin currents of spins $(2, \frac{5}{2}, \frac{5}{2}, 3)$, $(\frac{5}{2}, 3, 3, \frac{7}{2})$, $(\frac{5}{2}, 3, 3, \frac{7}{2})$ and $(3, \frac{7}{2}, \frac{7}{2}, 4)$ were determined from the right hand sides of these OPEs [32]. Moreover, the composite fields consisting of both the 11 currents in the large $\mathcal{N} = 4$ nonlinear superconformal algebra and the above 16 lowest higher spin currents also occurred in the right hand sides of these OPEs. The latter appeared quadratically (and linearly) in the fusion rules together with large $\mathcal{N} = 4$ nonlinear superconformal family of the identity operator.

[1] The 'supersymmetric' version of this coset can be written as $\frac{SU(N+M)_{k+N+M}}{SU(N)_{k+N+M} \times SU(M)_{k+N+M} \times U(1)}$ where the central charge is given by $c = \frac{3kMN}{(k+M+N)}$. For $M = 1$, the minimal model holography was described in [8]. Moreover, the duality property has been studied in [9, 10, 11, 12, 13, 14] further. For general M, further analysis has been done in [15, 16].

The asymptotic symmetry of the AdS_3 higher spin theory [33] based on super higher spin algebra $shs_{M=2}[\lambda]$ has been studied further and matches with those of the two-dimensional CFT Wolf space coset in the 't Hooft limit [34]. The extension of large $\mathcal{N}=4$ superconformal algebra which contains one $\mathcal{N}=4$ multiplet for each integer superspin $s=1,2,\cdots$ as well as the currents of large $\mathcal{N}=4$ superconformal algebra has been studied in [35]. For the particular level at the Kazama-Suzuki model [36, 37], the $\mathcal{N}=3$ (enhanced) supersymmetry is observed in [38]. This behavior also appeared in many previous examples of [39, 40, 41]. Furthermore, the full spectrum of the tensionless string theory in $AdS_3\times\mathbf{S}^3\times\mathbf{T}^4$ can be reorganized in terms of representations of the $\mathcal{N}=4$ super $\mathcal{W}_\infty$ algebra where the large level limit is taken [42].

In [43], the possible 'bosonic' coset theory with $O(M)$ Chan-Paton factor is described by

$$\frac{SO(N+M)_k\times SO(NM)_1}{SO(N)_{k+M}\times SO(M)_{k+N}}. \tag{1.1}$$

The central charge, by inserting the dimension of group, the level and the dual Coxeter number of the group into the central charge formula of each factor group, is given by [2]

$$c=\frac{3kMN}{2(M+N+k-2)}. \tag{1.2}$$

In the 'stringy' limit where M,N and k go to infinity simultaneously, this central charge is proportional to N^2 as appropriate for a stringy model [44]. This central charge looks similar to the one in the unitary Wolf space coset in the footnote 1. The 'supersymmetric' coset theory in the supersymmetric version of (1.1) is

$$\frac{SO(N+M)_{k+N+M-2}}{SO(N)_{k+N+M-2}\times SO(M)_{k+N+M-2}}. \tag{1.3}$$

Let us describe each case for this supersymmetric coset model.

For $M=1$, the above coset (1.1) or (1.3), by multiplying the appropriate $SO(1)_{k+N-1}$ in the numerator of the coset (1.1), reproduces the one studied in [45, 46] where the supersymmetry is equal to $\mathcal{N}=1$.

[2] One has $\frac{1}{2}(N+M)(N+M-1)\left[\frac{k}{k+(N+M-2)}\right]+\frac{1}{2}NM(NM-1)\left[\frac{1}{1+(NM-2)}\right]-\frac{1}{2}N(N-1)\left[\frac{(k+M)}{(k+M)+(N-2)}\right]-\frac{1}{2}M(M-1)\left[\frac{(k+N)}{(k+N)+(M-2)}\right]=\frac{3kMN}{2(M+N+k-2)}$. Or splitting the k independent part of this expression as $\frac{1}{4}\left[(N+M)(N+M-1)-N(N-1)-M(M-1)\right]$ and distributing it into other k dependent parts, one can also reexpress the above central charge as $\frac{3}{2}\frac{1}{2}(N+M)(N+M-1)\left[1-\frac{2(N+M-2)}{3(k+M+N-2)}\right]-\frac{3}{2}\frac{1}{2}N(N-1)\left[1-\frac{2(N-2)}{3(k+N+M-2)}\right]-\frac{3}{2}\frac{1}{2}M(M-1)\left[1-\frac{2(M-2)}{3(k+N+M-2)}\right]$ which is the central charge for the 'supersymmetric' coset theory in (1.3).

For $M = 2$, the above coset (1.1) or (1.3) has been described in $\mathcal{N} = 2$ Kazama-Suzuki model [36, 37] [3].

For $M = 4$, the above coset (1.1) or (1.3) reproduces the orthogonal Wolf space [4] by realizing that the $SO(4)$ group in the denominator can be interpreted as a product of two $SU(2)$ groups. In this case, the Wolf space central charge, by substituting $M = 4$ into (1.2), is given by

$$c_W = \frac{6kN}{(N+k+2)}. \tag{1.4}$$

We will see that this central charge appears in the highest singular term in the OPE between the two different fermionic spin-$\frac{3}{2}$ currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra [5].

In this paper, we would like to construct the 16 generators of the large $\mathcal{N} = 4$ linear superconformal algebra in the coset Wolf $\times SU(2) \times U(1) = \frac{SO(N+4)}{SO(N)\times SU(2)} \times U(1)$ (that is the fourth entry of the table 1 in [4]) theory with $N = 4$. See also the papers [49, 50, 51] on the generalization of GKO coset construction. By factoring out the spin-1 current and four spin-$\frac{1}{2}$ currents [28], the 11 generators of the large $\mathcal{N} = 4$ nonlinear algebra in the Wolf space coset are obtained [6]. What are the spin contents for the lowest higher spin currents? From the lessons in [52, 53, 54, 55, 56], one should have the spin-4 current. After putting this spin-4 current into the last place of the $\mathcal{N} = 4$ multiplet and then the other spin contents can be filled automatically. Let us denote them by spin contents as follows:

$$\begin{aligned}
&\left(2, \frac{5}{2}, \frac{5}{2}, 3\right) : (T^{(2)}, T_+^{(\frac{5}{2})}, T_-^{(\frac{5}{2})}, T^{(3)}), \\
&\left(\frac{5}{2}, 3, 3, \frac{7}{2}\right) : (U^{(\frac{5}{2})}, U_+^{(3)}, U_-^{(3)}, U^{(\frac{7}{2})}), \\
&\left(\frac{5}{2}, 3, 3, \frac{7}{2}\right) : (V^{(\frac{5}{2})}, V_+^{(3)}, V_-^{(3)}, V^{(\frac{7}{2})}), \\
&\left(3, \frac{7}{2}, \frac{7}{2}, 4\right) : (W^{(3)}, W_+^{(\frac{7}{2})}, W_-^{(\frac{7}{2})}, W^{(4)}).
\end{aligned} \tag{1.5}$$

[3] It would be interesting to obtain the higher spin currents as done in [47, 11, 13].

[4] This orthogonal Wolf space appeared in [48] previously.

[5] It is natural to ask what happens for the $M = 3$ case. It would be interesting to find the corresponding CFT explicitly. We thank S.-J. Rey for raising this issue.

[6] As in the unitary Wolf space, one can combine the spin-1 current and one spin-$\frac{1}{2}$ current as a single $\mathcal{N} = 1$ super current. The superpartners of the remaining three spin-$\frac{1}{2}$ currents (i.e. three spin-1 currents) do not play an important role in the denominator subgroup. When one divides $SU(2) \times U(1)$ factor, then one obtains the 'supersymmetric' Wolf space coset Wolf $= \frac{SO(N+4)}{SO(N)\times SU(2)\times SU(2)}$ where the second $SU(2)$ factor in the denominator refers only to the above three spin-$\frac{1}{2}$ currents. Their superpartners (three spin-1 currents) appear in the group $SO(N+4)$ and contribute to the $SU(2)_k$ affine algebra. The first $SU(2)$ contains the above $\mathcal{N} = 1$ super current.

We would like to construct these 16 currents (1.5) in terms of $\mathcal{N}=2$ affine Kac-Moody currents in the above orthogonal Wolf space coset theory (1.3) with $N=4$ and $M=4$ explicitly by following the previous works in [20, 32]. Furthermore, we calculate the various OPEs between the 11 generators of the large $\mathcal{N}=4$ nonlinear superconformal algebra and the 16 higher spin currents [7].

In section 2, the 16 currents of $\mathcal{N}=4$ linear superconformal algebra are obtained in the 'supersymmetric' coset Wolf $\times SU(2) \times U(1) = \frac{SO(N+4)}{SO(N)\times SU(2)} \times U(1)$ with $N=4$. After that, the 11 currents of $\mathcal{N}=4$ nonlinear superconformal algebra are determined in the orthogonal Wolf space coset (1.1) or (1.3) with $N=M=4$.

In section 3, the extra 16 currents in (1.5) are obtained in the orthogonal Wolf space coset (1.1) or (1.3). The OPEs between the 11 currents in section 2 and 16 currents in section 3 are obtained.

In section 4, we summarize the main results of this paper together with future directions.

In Appendices *A*-*D*, some details which are necessary in sections 2-3 are presented.

The mathematica package by Thielemans [58] is used.

2 The large $\mathcal{N}=4$ linear and nonlinear superconformal algebras in the coset minimal model

In this section, we would like to construct the large $\mathcal{N}=4$ nonlinear superconformal algebra realization. Instead of making it directly from the $\mathcal{N}=2$ WZW affine currents, we construct the large $\mathcal{N}=4$ linear superconformal algebra realization and then its nonlinear version can be obtained by factoring out the four fermion spin-$\frac{1}{2}$ currents and one spin-1 current.

2.1 $\mathcal{N}=2$ WZW affine current algebra

Let us describe the particular $\mathcal{N}=4$ superconformal coset theory described in [4] which can be written as

$$\text{Wolf} \times SU(2) \times U(1) = \frac{SO(N+4)}{SO(N) \times SU(2)} \times U(1), \ \text{ where } \ \text{Wolf} = \frac{SO(N+4)}{SO(N) \times [SU(2)]^2}. \quad (2.1)$$

More explicitly, one has (1.3) with $M=4$ and the levels $(k+N+M-2)$ at each factor group. The $SO(4)$ group is written in terms of two $SU(2)$ groups in the denominator of the orthogonal Wolf space coset.

[7] The higher spin theory with extended supersymmetries where the two dimensional coset theory contains Wolf space was studied also in [57].

The central charge of this coset model [36] is given by

$$c_{SO(N+4)} - c_{SO(N)} - c_{SU(2)} + c_{U(1)} = \frac{6(k+1)(N+1)}{(k+N+2)}, \tag{2.2}$$

where each contribution can be obtained from the following expressions

$$\begin{aligned} c_{SO(N+4)} &= \frac{3}{2}\left[\frac{1}{2}(N+4)(N+3)\right]\left[1-\frac{2(N+2)}{3(k+N+2)}\right], \\ c_{SO(N)} &= \frac{3}{2}\left[\frac{1}{2}N(N-1)\right]\left[1-\frac{2(N-2)}{3(k+N+2)}\right], \\ c_{SU(2)} &= \frac{3}{2}\left[2^2-1\right]\left[1-\frac{2\cdot 2}{3(k+N+2)}\right], \quad c_{U(1)} = \frac{3}{2}. \end{aligned} \tag{2.3}$$

This central charge (2.2) is the same as the unitary case in [4, 1, 20, 32] and will appear in the OPE between the spin-2 stress energy tensor of large $\mathcal{N}=4$ linear superconformal algebra as usual. The two levels of two $SU(2)$ affine algebras of large $\mathcal{N}=4$ linear superconformal algebra are given by the two positive integers (k, N) of the coset theory as follows $k^+ = k+1$ and $k^- = N+1 = 5$. One can also express the central charge in terms of (k^+, k^-) as $c = \frac{6k^+k^-}{(k^++k^-)}$. In the present case, because one chooses N as 4, the common factor $(k+N+2) = (k+6)$ will be present in all the important expressions later. For the previous case in [20, 32], one could see the $(k+N+2) = (k+5)$ dependence because we took N as 3.

Furthermore, the central charge (1.4) of the orthogonal Wolf space (2.1) can be obtained as follows:

$$c_{\rm Wolf} = c_{SO(N+4)} - c_{SO(N)} - 2\,c_{SU(2)} = \frac{6kN}{(2+k+N)}, \tag{2.4}$$

where the previous relations (2.3) are used. This can be written in terms of $c_{\rm Wolf} = \frac{6(k^+-1)(k^--1)}{(k^++k^-)}$. We will see this central charge in the OPE between the spin-$\frac{3}{2}$ currents of large $\mathcal{N}=4$ nonlinear superconformal algebra. Note that the central charge, in the usual OPE between the stress energy tensor in the large $\mathcal{N}=4$ nonlinear superconformal algebra, will take a different form.

The 28 basic $\mathcal{N}=2$ WZW affine currents of spin $\frac{1}{2}$ living in the orthogonal $SO(N+4=8)$ group are classified as follows:

$$\begin{aligned} &\text{subgroup } SO(4)\times SU(2)\times SU(2) \text{ currents}: \quad K^m = (K^1,\, K^2,\, K^3;\, K^4,\, K^5,\, K^6), \\ &\text{subgroup } SO(4)\times SU(2)\times SU(2) \text{ currents}: \quad K^{\bar m} = (K^{\bar 1},\, K^{\bar 2},\, K^{\bar 3};\, K^{\bar 4},\, K^{\bar 5},\, K^{\bar 6}), \\ &\text{Wolf coset } \frac{SO(8)}{SO(4)\times SU(2)\times SU(2)} \text{ currents}: \quad J^a = (J^7,\, J^8,\, \cdots,\, J^{14}), \\ &\text{Wolf coset } \frac{SO(8)}{SO(4)\times SU(2)\times SU(2)} \text{ currents}: \quad J^{\bar a} = (J^{\bar 7},\, J^{\bar 8},\, \cdots,\, J^{\bar{14}}). \end{aligned} \tag{2.5}$$

For $N = 4$, the 28 adjoint indices of $SO(8)$ can be divided in terms of 14 unbarred indices and 14 barred indices in the complex basis [8]. The nonvanishing structure constants and the generators are described in Appendix A. The $K^m(z)$ and $K^{\bar{m}}(z)$ of (2.5) live in the subgroup $SO(4) \times SU(2) \times SU(2)$ of $SO(8)$. More precisely, the 6 adjoint indices of the subgroup $SO(4)$ are given by $1, 2, 3$ (and their barred indices, $\bar{1}, \bar{2}, \bar{3}$). The indices of the first $SU(2)$ are given by 4, and $\bar{4}$, and some combination between 5 and $\bar{5}$ while those of the second $SU(2)$ are given by 6, and $\bar{6}$, and other combination between 5 and $\bar{5}$. On the other hand, the currents $J^a(z)$ and $J^{\bar{a}}(z)$ live in the Wolf coset $\frac{SO(8)}{SO(4)\times SU(2)\times SU(2)}$ for which the remaining 8 unbarred indices and 8 barred indices are present. One has similar spin-1 currents, denoted by $DK^{\bar{m}}|_{\theta=\bar{\theta}=0}(z)$, $\overline{D}K^m|_{\theta=\bar{\theta}=0}(z)$, $DJ^{\bar{a}}|_{\theta=\bar{\theta}=0}(z)$ and $\overline{D}J^a|_{\theta=\bar{\theta}=0}(z)$, corresponding to the superpartners of the above 28 spin-$\frac{1}{2}$ currents [9].

The complete OPEs for the $\mathcal{N} = 2$ WZW affine currents are summarized in Appendix B.

2.2 Large $\mathcal{N} = 4$ linear superconformal algebra in terms of $\mathcal{N} = 2$ WZW affine currents

We will construct the large $\mathcal{N} = 4$ linear superconformal algebra realization for the coset theory (2.1) using the description of previous presentation.

[8] The adjoint representation **28** of $SO(8)$ breaks into $\mathbf{28} \to (\mathbf{1},\mathbf{3},\mathbf{1},\mathbf{1}) \oplus (\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1}) \oplus (\mathbf{1},\mathbf{1},\mathbf{3},\mathbf{1}) \oplus (\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{3}) \oplus (\mathbf{2},\mathbf{2},\mathbf{2},\mathbf{2})$ under the $SU(2) \times SU(2) \times SU(2) \times SU(2)$ [59] where the $SO(4)$ is replaced with the first two $SU(2)$ factors. The first two representations are given by the currents having the indices $1, 2, 3$ (and their complex conjugated ones). The next two representations are given by the currents having the indices $4, 5, 6$ (and their complex conjugated ones). The remaining representations are given by the currents having the indices $7, 8, \cdots, 14$ (and their complex conjugated ones).

[9] It is also useful to write down the spin-1 affine current in terms of purely bosonic spin-1 current and spin-$\frac{1}{2}$ current. One has the following expressions [17]

$$\begin{aligned}
\overline{D}K^m|_{\theta=\bar{\theta}=0}(z) &= V^m(z) - \frac{1}{(k+N+2)} \left(f_{p\bar{q}}{}^m K^p K^{\bar{q}} + f_{a\bar{b}}{}^m J^a J^{\bar{b}} \right)(z), \\
DK^{\bar{m}}|_{\theta=\bar{\theta}=0}(z) &= V^{\bar{m}}(z) - \frac{1}{(k+N+2)} \left(f_{\bar{p}q}{}^{\bar{m}} K^{\bar{p}} K^q + f_{m\bar{a}}{}^{\bar{b}} J^{\bar{a}} J^b \right)(z), \\
\overline{D}J^a|_{\theta=\bar{\theta}=0}(z) &= V^a(z) - \frac{1}{(k+N+2)} \left(f_{b\bar{m}}{}^a J^b K^{\bar{m}} + f_{\bar{a}m}{}^b K^m J^{\bar{b}} \right)(z), \\
DJ^{\bar{a}}|_{\theta=\bar{\theta}=0}(z) &= V^{\bar{a}}(z) - \frac{1}{(k+N+2)} \left(f_{\bar{b}m}{}^{\bar{a}} J^{\bar{b}} K^m + f_{a\bar{m}}{}^{\bar{b}} K^{\bar{m}} J^b \right)(z).
\end{aligned} \tag{2.6}$$

The OPEs between the purely bosonic spin-1 currents, which commute with the above spin-$\frac{1}{2}$ currents, are given in [32]: $V^A(z)\, V^B(w) = -\frac{1}{(z-w)}\, f_{\bar{A}\bar{B}}{}^{\bar{C}} V^C(w) + \cdots$, $V^{\bar{A}}(z)\, V^{\bar{B}}(w) = -\frac{1}{(z-w)}\, f_{AB}{}^C V^{\bar{C}}(w) + \cdots$ and $V^A(z)\, V^{\bar{B}}(w) = \frac{1}{(z-w)^2}\, k\, \delta^{A\bar{B}} - \frac{1}{(z-w)} \left(f_{\bar{A}B}{}^{\bar{C}} V^C + f_{\bar{A}B}{}^C V^{\bar{C}} \right)(w) + \cdots$, where the indices $A, B, \cdots$ and $\bar{A}, \bar{B}, \cdots$ stand for the indices of the group $SO(N+4=8)$ in the complex basis. Again the two extra terms of last two equations in (2.6) appear due to the nonvanishing structure constants $f_{ab}{}^m$ and $f_{\bar{a}\bar{b}}{}^{\bar{m}}$ which vanish in the unitary case [20].

2.2.1 Construction of two spin-$\frac{3}{2}$ currents

Let us consider the two spin-$\frac{3}{2}$ currents of large $\mathcal{N}=4$ linear superconformal algebra, which appear in the last components of $\mathcal{N}=2$ multiplets $(\frac{1}{2},1,1,\frac{3}{2})$ and $(\frac{1}{2},1,1,\frac{3}{2})$ [20]. Let us denote the first one as $G_{11}(z)$ and the second one as $G_{22}(z)$. As in the unitary case [20], one should consider the spin-$\frac{1}{2}$ current with the Wolf space subgroup index. Among the two $SU(2)$ subgroups, the second $SU(2)$ group corresponds to the above spin-$\frac{3}{2}$ currents. It turns out that the spin-$\frac{1}{2}$ current with the index 6 provides the correct spin-$\frac{3}{2}$ current

$$G_{11}(z) = \sqrt{2}i\left(-\frac{1}{2}\left[D,\overline{D}\right]K^{6} - \frac{(k-4)}{2(6+k)}\partial K^{6}\right)(z). \tag{2.7}$$

Note that $K^{6}(w)$ and $K^{\bar{6}}(w)$ are primary under the $T_{SO(8)}(z)$ and $(T_{SO(8)}-T_{SO(4)})(z)$. The coefficient $\frac{(k-4)}{(6+k)}$ can be written as $\frac{(k^{+}-k^{-})}{(k^{+}+k^{-})}$ with

$$k^{+}=k+1, \qquad \text{and} \qquad k^{-}=N+1=5.$$

The second term in (2.7) is necessary for the primary field condition on $G_{11}(w)$ under the $T_{SO(8)}(z)$.

The second spin-$\frac{3}{2}$ current can be obtained as follows:

$$G_{22}(z) = \sqrt{2}i\left(-\frac{1}{2}\left[D,\overline{D}\right]K^{\bar{6}} + \frac{(k-4)}{2(6+k)}\partial K^{\bar{6}}\right)(z). \tag{2.8}$$

One can find some informations on other currents (or Wolf space central charge c_{Wolf}) by calculating the OPE between (2.7) and (2.8) and reading off the right hand side.

2.2.2 Construction of three spin-1 currents

Now one continues to obtain the other components in the previous two $\mathcal{N}=2$ multiplets of large $\mathcal{N}=4$ linear superconformal algebra. For example, one can construct the following spin-1 currents associated with the indices 6 and $\bar{6}$ as follows:

$$\begin{aligned} A_1(z) &= \frac{1}{2}\left(-\overline{D}K^{6} + DK^{\bar{6}}\right)(z), \\ A_2(z) &= \frac{i}{2}\left(\overline{D}K^{6} + DK^{\bar{6}}\right)(z). \end{aligned} \tag{2.9}$$

Note that there are no singular terms in the OPEs of $\overline{D}K^{6}(z)\,\overline{D}K^{6}(w)$ and $DK^{\bar{6}}(z)\,DK^{\bar{6}}(w)$. It is easy to determine the overall constants in (2.9) because they are the affine currents of $SU(2)_{k^{+}(=k+1)}$ algebra. Then one can fix the remaining current by calculating the OPE

between $A_1(z)$ and $A_2(w)$ and reading off the first-order pole

$$\begin{aligned} A_3(w) &= \frac{1-i}{2\sqrt{2}}(\overline{D}K^5 - i\,DK^{\bar{5}})(w) \\ &- \frac{i}{2(6+k)}\left(\sum_{(m,\bar{m})=(5,\bar{5})}^{(6,\bar{6})} K^m K^{\bar{m}} + \sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})} J^a J^{\bar{a}}\right)(w). \end{aligned} \tag{2.10}$$

Then it turns out that the OPEs between these spin-1 currents are the same as the ones in (2.17) of [20].

2.2.3 Construction of four spin-$\frac{1}{2}$ currents and other two spin-$\frac{3}{2}$ currents

Let us determine the four spin-$\frac{1}{2}$ currents of large $\mathcal{N}=4$ linear superconformal algebra. Let us focus on the OPE between $A_1(z)$ and $G_{11}(w)$ from (2.9) and (2.7) [10]. This OPE is nontrivial and contains the second-order pole and first-order pole. The deformation parameter of large $\mathcal{N}=4$ linear superconformal algebra is defined as

$$\gamma = \frac{k^-}{(k^+ + k^-)} = \frac{5}{(6+k)}, \tag{2.11}$$

and the second-order pole provides the following spin-$\frac{1}{2}$ current as follows:

$$F_{21}(w) = \frac{(1+i)}{2}\,K^5(w). \tag{2.12}$$

Moreover, the first-order pole determines the following spin-$\frac{3}{2}$ current

$$G_{21}(w) = \frac{\sqrt{2}}{(6+k)}\left(\sum_{(m,\bar{m})=(5,\bar{5})}^{(6,\bar{6})} K^m DK^{\bar{m}} + \sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})} J^a DJ^{\bar{a}}\right)(w) + \text{cubic terms}. \tag{2.13}$$

Let us describe the OPE between $A_1(z)$ and $G_{22}(w)$ from (2.9) and (2.8) [11]. In this case, this nontrivial OPE contains the second-order pole and first-order pole. The former determines the spin-$\frac{1}{2}$ current

$$F_{12}(w) = \frac{(1-i)}{2}\,K^{\bar{5}}(w), \tag{2.14}$$

and the latter fixes the spin-$\frac{3}{2}$ current

$$\begin{aligned} G_{12}(w) &= \frac{5(1-i)}{(6+k)}\partial K^{\bar{5}}(w) - \frac{\sqrt{2}}{6+k}\left(\sum_{(m,\bar{m})=(5,\bar{5})}^{(6,\bar{6})} \overline{D}K^m K^{\bar{m}} + \sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})} \overline{D}J^a J^{\bar{a}}\right)(w) \\ &+ \text{cubic terms}. \end{aligned} \tag{2.15}$$

[10] That is, $A_1(z)\,G_{11}(w) = \frac{1}{(z-w)^2}\,\frac{i}{2}\,2(1-\gamma)\,F_{21}(w) - \frac{1}{(z-w)}\,\frac{i}{2}\,G_{21}(w) + \cdots$.

[11] In other words, $A_1(z)\,G_{22}(w) = -\frac{1}{(z-w)^2}\,\frac{i}{2}\,2(1-\gamma)\,F_{12}(w) + \frac{1}{(z-w)}\,\frac{i}{2}\,G_{12}(w) + \cdots$, with (2.11).

Let us determine the remaining spin-$\frac{1}{2}$ currents. From the second order pole of the OPE between $A_3(z)$ and $G_{11}(w)$ [12], one finds the following spin-$\frac{1}{2}$ current

$$F_{11}(w) \;=\; \frac{i}{\sqrt{2}}\,K^{6}(w). \tag{2.16}$$

Furthermore, the OPE between $A_3(z)$ and $G_{22}(w)$ provides the last spin-$\frac{1}{2}$ current as follows [13]:

$$F_{22}(w) \;=\; -\frac{i}{\sqrt{2}}\,K^{\bar{6}}(w). \tag{2.17}$$

Then the Wolf space coset subgroup $SU(2)$ containing the indices 6 and $\bar{6}$ has the three spin-$\frac{1}{2}$ currents, $F_{11}(z)$, $F_{22}(z)$ and $(F_{12}+F_{21})(z)$. The nontrivial OPEs between these currents are given in (2.29) of [20]. The above $\mathcal{N}=2$ multiplets, $(\frac{1}{2},1,1,\frac{3}{2})$ and $(\frac{1}{2},1,1,\frac{3}{2})$, are determined (except the second component of first multiplet and third component of second multiplet) completely (see also (2.44) in [20]). Furthermore, the second and third components of the other $\mathcal{N}=2$ multiplets $(1,\frac{3}{2},\frac{3}{2},2)$ and $(0,\frac{1}{2},\frac{1}{2},1)$ are determined. Because the four spin-$\frac{3}{2}$ currents are known, one calculates their OPEs between themselves and obtains other informations on the other undetermined currents.

2.2.4 Construction of other three spin-1 currents

Let us determine the other spin-1 currents. Let us focus on the OPE between $F_{11}(z)$ and $G_{21}(w)$ (and similarly the OPE between $F_{12}(z)$ and $G_{22}(w)$) from (2.16), (2.13), (2.14) and (2.8). By combining these [14], then the two spin-1 currents are fixed as follows:

$$\begin{aligned}
B_1(w) &= -\frac{1}{2(6+k)}(B^{'}+B^{''})(w)-\frac{(1+i)}{2\sqrt{2}(6+k)}(K^5K^6-iK^{\bar{5}}K^{\bar{6}})(w),\\
B_2(w) &= -\frac{i}{2(6+k)}(B^{'}-B^{''})(w)+\frac{(1-i)}{2\sqrt{2}(6+k)}(K^5K^6+iK^{\bar{5}}K^{\bar{6}})(w),\\
B^{'}(w) &\equiv J^7J^{14}(w)+J^8J^{13}(w)-J^9J^{12}(w)-J^{10}J^{11}(w),\\
B^{''}(w) &\equiv J^{\bar{7}}J^{\bar{14}}(w)+J^{\bar{8}}J^{\bar{13}}(w)-J^{\bar{9}}J^{\bar{12}}(w)-J^{\bar{10}}J^{\bar{11}}(w).
\end{aligned} \tag{2.18}$$

After that, the remaining spin-1 current can be read off from the OPE between $B_1(z)$ and $B_2(w)$ as follows:

$$B_3(w) \;=\; -\frac{i}{2(6+k)}\left(\sum_{(m,\bar{m})=(5,\bar{5})}^{(6,\bar{6})}K^mK^{\bar{m}}+\sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})}J^aJ^{\bar{a}}\right)(w). \tag{2.19}$$

[12] One has $A_3(z)\,G_{11}(w)=\frac{1}{(z-w)^2}\,\frac{i}{2}\,2(1-\gamma)\,F_{11}(w)-\frac{1}{(z-w)}\,\frac{i}{2}\,G_{11}(w)+\cdots$.

[13] We have $A_3(z)\,G_{22}(w)=-\frac{1}{(z-w)^2}\,\frac{i}{2}\,2(1-\gamma)\,F_{22}(w)+\frac{1}{(z-w)}\,\frac{i}{2}\,G_{22}(w)+\cdots$.

[14] Explicitly one has $F_{11}(z)\,G_{21}(w) = \frac{1}{(z-w)}\,(-iB_1-B_2)\,(w)+\cdots$ and $F_{12}(z)\,G_{22}(w) = \frac{1}{(z-w)}\,(-iB_1+B_2)\,(w)+\cdots$.

The three spin-1 currents consist of the affine $SU(2)_{k^-=5}$ algebra. Note that these currents are made of quadratic fermions and contain the Wolf space coset indices as well as the Wolf space subgroup indices. In the next subsection, we will see that the latter will disappear in the new basis.

2.2.5 Construction of other spin-1 current

For the spin-1 current, one can use the OPE between $F_{11}(z)$ and $G_{22}(w)$ [15] and the first-order pole determines the following spin-1 current

$$U(z) \;=\; \frac{(1+i)}{2\sqrt{2}}(\overline{D}K^5 + i\,DK^{\bar{5}})(z). \tag{2.20}$$

This satisfies the correct OPE between $U(z)$ and $U(w)$ and one can also rewrite this spin-1 current from (2.12) and (2.14) as follows:

$$U(z) = \frac{1}{\sqrt{2}}\left(\overline{D}F_{21} - DF_{12}\right)(z). \tag{2.21}$$

One can construct $\mathcal{N}=1$ superfield as the sum of $(F_{21}-F_{12})(z)$ and $U(z)$ and they correspond to the first $SU(2)$ subgroup of the Wolf space coset.

2.2.6 Construction of spin-2 current

So far, we have found the 15 currents of large $\mathcal{N}=4$ linear superconformal algebra and should determine the last unknown spin-2 stress energy tensor. For example, from the OPE between $G_{11}(z)$ and $G_{22}(w)$, one can read off the stress energy tensor as follows [16]:

$$T(z) = T_{SO(8)}(z) - T_{SO(4)} - T^{'}(z), \;\; \text{where} \;\; T^{'}(z) = T_{SU(2)} - T_{U(1)}. \tag{2.24}$$

[15] One has $F_{11}(z)\,G_{22}(w) = \frac{1}{(z-w)}\,(-iA_3 - iB_3 + U)\,(w) + \cdots$.

[16] One can obtain the following OPEs

$$\begin{aligned}
T_{SO(8)}(z)\,T_{SO(8)}(w) &= \frac{1}{(z-w)^4}\,\frac{c_{SO(8)}}{2} + \frac{1}{(z-w)^2}\,2T_{SO(8)}(w) + \frac{1}{(z-w)}\,\partial T_{SO(8)}(w) + \cdots,\\
T_{SO(8)}(z)\,T_{SO(4)}(w) &= \frac{1}{(z-w)^4}\,\frac{c_{SO(4)}}{2} + \frac{1}{(z-w)^2}\,2T_{SO(4)}(w) + \frac{1}{(z-w)}\,\partial T_{SO(4)}(w) + \cdots,\\
T_{SO(8)}(z)\,T^{'}(w) &= \frac{1}{(z-w)^4}\,\frac{c^{'}}{2} + \frac{1}{(z-w)^2}\,2T^{'}(w) + \frac{1}{(z-w)}\,\partial T^{'}(w) + \cdots,\\
T_{SO(4)}(w)\,T_{SO(4)}(w) &= \frac{1}{(z-w)^4}\,\frac{c_{SO(4)}}{2} + \frac{1}{(z-w)^2}\,2T_{SO(4)}(w) + \frac{1}{(z-w)}\,\partial T_{SO(4)}(w) + \cdots,\\
T^{'}(z)\,T^{'}(w) &= \frac{1}{(z-w)^4}\,\frac{c^{'}}{2} + \frac{1}{(z-w)^2}\,2T^{'}(w) + \frac{1}{(z-w)}\,\partial T^{'} + \cdots,\\
T_{SO(4)}(z)\,T^{'}(w) &= +\cdots,
\end{aligned} \tag{2.22}$$

Then the standard OPE between the stress tensor $T(z)$, which can be obtained from (2.22), has the following form

$$T(z)\,T(w) \;=\; \frac{1}{(z-w)^4}\,\frac{c}{2} + \frac{1}{(z-w)^2}\,2T(w) + \frac{1}{(z-w)}\,\partial T(w) + \cdots, \qquad (2.25)$$

where the central charge is given in (2.23). Furthermore, one has

$$(T_{SO(4)} + T')(z)\,\Phi(w) \;=\; +\cdots, \qquad (2.26)$$

where $\Phi(w)$ in (2.26) is the 16 currents of large $\mathcal{N}=4$ linear superconformal algebra. Also the OPEs between $K^m(z)$ with $m=1,2,3$ (or $K^{\bar{m}}(z)$ with $\bar{m}=\bar{1},\bar{2},\bar{3}$) and these 16 currents do not have singular terms.

Summarizing this subsection, the large $\mathcal{N}=4$ linear superconformal algebra has four spin-$\frac{1}{2}$ currents with (2.16), (2.14), (2.12) and (2.17), seven spin-1 currents with (2.9), (2.10), (2.18), (2.19) and (2.20), four spin-$\frac{3}{2}$ currents with (2.7), (2.15), (2.13) and (2.8), and spin-2 current with (2.24). Among spin-1 currents, the three currents corresponding to $SU(2)_{k^+}$ affine algebra and other three currents corresponding to $SU(2)_{k^-}$ affine algebra contain both Wolf space subgroup index and the Wolf space coset index. In particular, the spin-$\frac{1}{2}$ current $F_{11}(z)$, the spin-1 current $A_+(z)(\equiv (A_1+iA_2)(z))$ and the spin-$\frac{3}{2}$ current $G_{11}(z)$ are located at the same $\mathcal{N}=2$ multiplet. Similarly, the spin-$\frac{1}{2}$ current $F_{22}(z)$, the spin-1 current $A_-(z)(\equiv (A_1-iA_2)(z))$ and the spin-$\frac{3}{2}$ current $G_{22}(z)$ are located at the same other $\mathcal{N}=2$ multiplet. The relevant affine spin-1 currents for these live in the second $SU(2)$ subgroup of the Wolf space coset.

2.3 Large $\mathcal{N}=4$ nonlinear superconformal algebra realization

As described before, one should construct the large $\mathcal{N}=4$ nonlinear superconformal algebra from its linear version by decoupling four spin-$\frac{1}{2}$ currents, $F_{11}(z)$, $F_{12}(z)$, $F_{21}(z)$ and $F_{22}(z)$ and one spin-1 current $U(z)$.

2.3.1 Construction of spin-2 stress tensor

Let us consider the stress energy tensor $\hat{T}(z)$ of large $\mathcal{N}=4$ nonlinear superconformal algebra [28, 29, 30]. By definition, this should satisfy the following regular conditions with (2.20) (or

where the central charges are given by

$$\begin{aligned} c_{SO(8)} &= \frac{42(2+k)}{(6+k)}, \qquad c_{SO(4)} = \frac{3(14+3k)}{(6+k)}, \qquad c^{'} = c_{SU(2)} - c_{U(1)} = \frac{3(14+3k)}{2(6+k)} - \frac{3}{2} = \frac{3(4+k)}{(6+k)}, \\ c &= c_{SO(8)} - c_{SO(4)} - c^{'} = \frac{30(1+k)}{(6+k)}. \end{aligned} \qquad (2.23)$$

The above OPEs can be used to calculate the OPE in (2.25).

(2.21)), (2.16), (2.14), (2.12) and (2.17),

$$U(z)\,\hat{T}(w) \;=\; +\cdots, \qquad F_a(z)\,\hat{T}(w) = +\cdots, \qquad a = 11, 12, 21, 22. \tag{2.27}$$

For $N = 4$, the stress energy tensor of the large $\mathcal{N} = 4$ nonlinear algebra can be written as [28, 29, 30]

$$\begin{aligned}
\hat{T}(z) &= T(z) + \frac{1}{(6+k)} UU(z) + \frac{1}{(6+k)} \partial F^a F_a(z) \\
&= T_{SO(8)}(z) - T_{SO(4)}(z) - T_{SU(2)}(z) + T_{U(1)}(z) \\
&+ \frac{1}{(6+k)} \left(UU + \partial F^a F_a\right)(z).
\end{aligned} \tag{2.28}$$

The relative coefficients in the extra terms are fixed from the conditions (2.27). The stress energy tensor for general N case can be obtained by putting $(6+k) \to (k+N+2)$ and $T \to T_{SO(N+4)} - T_{SO(N)} - T_{SU(2)} + T_{U(1)}$ in (2.28) similarly. One can read off the corresponding central charge appearing in the Virasoro algebra by calculating the OPE $\hat{T}(z)\,\hat{T}(w)$. Therefore, the total central charge is given by

$$\hat{c} = \frac{6(k+1)(N+1)}{(k+N+2)} - 3 = \frac{3(k+N+2kN)}{(k+N+2)} \to \frac{3(4+9k)}{(6+k)}, \tag{2.29}$$

where the $N = 4$ is substituted in the last stage. The remaining 10 currents will be primary fields under this stress energy tensor.

2.3.2 Construction of six spin-1 currents

Let us determine the other currents. For the spin-1 current, one can add 6 additional terms coming from four fermions, $F_{11}(z)$, $F_{12}(z)$, $F_{21}(z)$ and $F_{22}(z)$. By substituting (2.9), (2.10), (2.16), (2.14), (2.12) and (2.17) into the equation (3.6) (where $(5+k)$ is replaced with $(6+k)$) of [20], one obtains

$$\begin{aligned}
\hat{A}_1(z) &= -\frac{(1+i)\sqrt{2}}{4(6+k)} (K^5 K^{\bar{6}} + iK^6 K^{\bar{5}})(z) - \frac{1}{2} (\overline{D} K^6 - D K^{\bar{6}})(z), \\
\hat{A}_2(z) &= -\frac{(-1+i)\sqrt{2}}{4(6+k)} (K^5 K^{\bar{6}} - iK^6 K^{\bar{5}})(z) + \frac{i}{2} (\overline{D} K^6 + D K^{\bar{6}})(z), \\
\hat{A}_3(z) &= -\frac{i}{2(6+k)} \left(\sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})} J^a J^{\bar{a}} + 2K^6 K^{\bar{6}} \right)(z) - \frac{(1+i)\sqrt{2}}{4} \left(i\overline{D} K^5 + D K^{\bar{5}} \right)(z).
\end{aligned} \tag{2.30}$$

One can express these in terms of purely bosonic spin-1 currents appeared in (2.6). That is,

$$\hat{A}_+(z) \;=\; -V^6(z), \qquad \hat{A}_-(z) = V^{\bar{6}}(z), \qquad \hat{A}_3(z) = \frac{(1-i)\sqrt{2}}{4} \left(V^5 - iV^{\bar{5}} \right)(z), \tag{2.31}$$

where we introduce $\hat{A}_{\pm}(z) \equiv (\hat{A}_1 \pm i\hat{A}_2)(z)$. Compared to the previous expression (2.30), all the quadratic terms are disappeared. These three currents consist of $SU(2)_{\hat{k}^+}$ affine algebra with $\hat{k}^+ = k^+ - 1 = k$. Let us look at the second subgroup $SU(2)$ in the Wolf space coset. The indices of spin-$\frac{1}{2}$ currents, $F_{11}(z)$, $F_{22}(z)$ and $(F_{12}+F_{21})(z)$ stand for the adjoint representation of this second subgroup $SU(2)$. All the higher spin currents including the 11 currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra should commute with these three fermionic currents. The superpartners of these fermionic currents correspond to the above three spin-1 currents (2.31) plus other terms [17].

Let us move on the other type of spin-1 currents. As done before, the spin-1 currents, $B_1(z)$, $B_2(z)$ and $B_3(z)$ can be modified under the factoring out the fermions and spin-1 current. By substituting (2.18), (2.19), (2.16), (2.14), (2.12) and (2.17) into the equation (3.10) with a replacement of $(6 + k)$ in [20], one has

$$\begin{aligned}
\hat{B}_1(z) &= -\frac{1}{2(6+k)}\left(B^{'} + B^{''}\right)(z), \\
\hat{B}_2(z) &= -\frac{i}{2(6+k)}\left(B^{'} - B^{''}\right)(z), \\
\hat{B}_3(z) &= -\frac{i}{2(6+k)} \sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})} J^a J^{\bar{a}}(z),
\end{aligned} \tag{2.32}$$

where $B^{'}(z)$ and $B^{''}(z)$ are given in (2.18). All the $K^m(z)$ and $K^{\bar{m}}(z)$ dependences are disappeared because the additional quadratic terms in the fermions of $\hat{B}_i(z)$ cancel those terms. Note that these three currents have only the Wolf space coset indices and satisfy the $SU(2)_{\hat{k}^-}$ affine algebra with $\hat{k}^-(= k^- - 1 = N = 4)$. Because the purely bosonic currents $V^m(z)$ or $V^{\bar{m}}(z)$ in (2.31) commute with fermions $J^a(z)$ or $J^{\bar{a}}(z)$ in (2.32), one sees that there are no singular terms in the OPEs between $\hat{A}_i(z)$ and $\hat{B}_j(w)$. In this respect, the expression in (2.31) is more useful than the one in (2.30) [18].

2.3.3 Construction of four spin-$\frac{3}{2}$ currents

Let us construct four spin-$\frac{3}{2}$ currents $\hat{G}_{11}(z), \hat{G}_{22}(z), \hat{G}_{12}(z)$, and $\hat{G}_{21}(z)$ [28, 29, 30]. By substituting the expressions (2.16), (2.14), (2.12), (2.17), (2.20), (2.7), (2.15), (2.13), (2.8),

[17] For general N, it is nontrivial to express the above three spin-1 currents. One expects that they can be written in terms of sum of some tensors with indices a multiplied by $V^a(z)$ and other tensors with indices $\bar{a}$ multiplied by $V^{\bar{a}}(z)$.

[18] One expects that for general N, the three spin-1 currents, $\hat{B}_+(z)$, $\hat{B}_-(z)$ and $\hat{B}_3(z)$ can be written as some tensors (whose values are 1 or -1) with indices a and b multiplied by $J^a J^b(z)$, some tensors (whose values are 1 or -1) with indices $\bar{a}$ and $\bar{b}$ multiplied by $J^{\bar{a}} J^{\bar{b}}(z)$ and other tensors (whose values are 1) with indices a and $\bar{b}$ multiplied by $J^a J^{\bar{b}}(z)$ respectively.

(2.30) and (2.32) into the equation (3.13) of [20], the following results can be obtained

$$\begin{aligned}
\hat{G}_{11}(z) &= \frac{i\sqrt{2}}{(6+k)}\Big(J^{7}\overline{D}J^{14}+J^{8}\overline{D}J^{13}-J^{9}\overline{D}J^{12}-J^{10}\overline{D}J^{11}+J^{11}\overline{D}J^{10} \\
&+ J^{12}\overline{D}J^{9}-J^{13}\overline{D}J^{8}-J^{14}\overline{D}J^{7}\Big)(z)+\text{cubic terms}, \qquad (2.33)\\
\hat{G}_{22}(z) &= \frac{i\sqrt{2}}{(6+k)}\Big(J^{\bar{7}}DJ^{\bar{14}}+J^{\bar{8}}DJ^{\bar{13}}-J^{\bar{9}}DJ^{\bar{12}}-J^{\bar{10}}DJ^{\bar{11}}+J^{\bar{11}}DJ^{\bar{10}} \\
&+ J^{\bar{12}}DJ^{\bar{9}}-J^{\bar{13}}DJ^{\bar{8}}-J^{\bar{14}}DJ^{\bar{7}}\Big)(z)+\text{cubic terms}, \qquad (2.34)\\
\hat{G}_{12}(z) &= \frac{4(1-i)}{(6+k)}\partial K^{\bar{5}}(z)-\frac{\sqrt{2}}{(6+k)}\sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})}\overline{D}J^{a}J^{\bar{a}}(z)+\text{cubic terms}, \qquad (2.35)\\
\hat{G}_{21}(z) &= \frac{\sqrt{2}}{(6+k)}\sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})}J^{a}DJ^{\bar{a}}(z)+\text{other cubic terms}. \qquad (2.36)
\end{aligned}$$

In this case, one also reexpresses them using the purely bosonic spin-1 currents [19]. All the cubic terms appearing in (2.33), (2.35), (2.36) or (2.34) are disappeared in (2.37). All the indices appearing in these four spin-$\frac{3}{2}$ currents are coming from the Wolf space coset indices. The various OPEs between these spin-$\frac{3}{2}$ currents are given in the equation $(A.3)$ of [20] where one should replace $(5+k)$ with $(6+k)$ and the Wolf space central charge c_{Wolf} is given in (2.4) (or the third equation in footnote 20).

2.3.4 $U(1)$ charges of the 11 currents of large $\mathcal{N}=4$ nonlinear superconformal algebra

As in the $U(1)$ current of (3.19) in [20], one can describe the $U(1)$ current here. By realizing that the $U(1)$ current appears in the second-order pole of the OPE $\hat{G}_{21}(z)\,\hat{G}_{12}(w)$, the $U(1)$

[19] One has alternative form as follows:

$$\begin{aligned}
\hat{G}_{11}(z) &= -\frac{i\sqrt{2}}{(6+k)}\left(J^{7}V^{14}+J^{8}V^{13}-J^{9}V^{12}-J^{10}V^{11}+J^{11}V^{10}+J^{12}V^{9}-J^{13}V^{8}-J^{14}V^{7}\right)(z),\\
\hat{G}_{22}(z) &= -\frac{i\sqrt{2}}{(6+k)}\Big(J^{\bar{7}}V^{\bar{14}}+J^{\bar{8}}V^{\bar{13}}-J^{\bar{9}}V^{\bar{12}}-J^{\bar{10}}V^{\bar{11}}+J^{\bar{11}}V^{\bar{10}}+J^{\bar{12}}V^{\bar{9}}-J^{\bar{13}}V^{\bar{8}}-J^{\bar{14}}V^{\bar{7}}\Big)(z),\\
\hat{G}_{12}(z) &= -\frac{\sqrt{2}}{(6+k)}\sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})}J^{\bar{a}}V^{a}(z), \qquad \hat{G}_{21}(z)=\frac{\sqrt{2}}{(6+k)}\sum_{(a,\bar{a})=(7,\bar{7})}^{(14,\bar{14})}J^{a}V^{\bar{a}}(z). \qquad (2.37)
\end{aligned}$$

It is obvious that the last two fermions in (2.37) living in the $\mathcal{N}=2$ stress energy tensor can be written as some tensors (whose values are 1) with indices $\bar{a}$ and b multiplied by $J^{\bar{a}}V^{b}(z)$ and other tensors (whose values are 1) with indices a and $\bar{b}$ multiplied by $J^{a}V^{\bar{b}}(z)$ respectively. For the first two in (2.37), one expects that they can be realized by some tensors (whose values are either 1 or -1) with indices a and b multiplied by $J^{a}V^{b}(z)$ and other tensors (whose values are either 1 or -1) with indices $\bar{a}$ and $\bar{b}$ multiplied by $J^{\bar{a}}V^{\bar{b}}(z)$ respectively.

current is identified with $\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)$ from the OPE (3.18) of [20]. Here

$$\gamma_A = \frac{\hat{k}^-}{(\hat{k}^+ + \hat{k}^- + 2)} = \frac{4}{(6+k)}, \qquad \gamma_B = \frac{\hat{k}^+}{(\hat{k}^+ + \hat{k}^- + 2)} = \frac{k}{(6+k)}. \tag{2.38}$$

Then one obtains the following first-order poles for the 11 currents as follows:

$$\begin{aligned}
\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)\,\hat{A}_\pm(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\mp\frac{8}{(6+k)}\right]\hat{A}_\pm(w), \\
\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)\,\hat{B}_\pm(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\mp\frac{2k}{(6+k)}\right]\hat{B}_\pm(w), \\
\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)\,\hat{A}_3(w)|_{\frac{1}{(z-w)}} &= 0, \\
\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)\,\hat{B}_3(w)|_{\frac{1}{(z-w)}} &= 0, \\
\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)\begin{pmatrix}\hat{G}_{11}\\ \hat{G}_{22}\end{pmatrix}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\pm\frac{(-4+k)}{(6+k)}\right]\begin{pmatrix}\hat{G}_{11}\\ \hat{G}_{22}\end{pmatrix}(w), \\
\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)\begin{pmatrix}\hat{G}_{12}\\ \hat{G}_{21}\end{pmatrix}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\mp\frac{(4+k)}{(6+k)}\right]\begin{pmatrix}\hat{G}_{12}\\ \hat{G}_{21}\end{pmatrix}(w), \\
\left(-2i\gamma_A\hat{A}_3 - 2i\gamma_B\hat{B}_3\right)(z)\,\hat{T}(w)|_{\frac{1}{(z-w)}} &= 0.
\end{aligned} \tag{2.39}$$

We present the $U(1)$ charges for the 11 currents in Table 1 (the relevant Tables 2-10 will appear later).

$U(1)$ charge	11 currents of large $\mathcal{N}=4$ nonlinear superconformal algebra
$\frac{2k}{(6+k)}$	$\hat{B}_-(\equiv \hat{B}_1 - i\hat{B}_2)$
$\frac{(4+k)}{(6+k)}$	$\hat{G}_{21}$
$\frac{(-4+k)}{(6+k)}$	$\hat{G}_{11}$
$\frac{8}{(6+k)}$	$\hat{A}_-(\equiv \hat{A}_1 - i\hat{A}_2)$
0	$\hat{A}_3, \quad \hat{B}_3, \quad \hat{T}$
$-\frac{8}{(6+k)}$	$\hat{A}_+(\equiv \hat{A}_1 + i\hat{A}_2)$
$-\frac{(-4+k)}{(6+k)}$	$\hat{G}_{22}$
$-\frac{(4+k)}{(6+k)}$	$\hat{G}_{12}$
$-\frac{2k}{(6+k)}$	$\hat{B}_+(\equiv \hat{B}_1 + i\hat{B}_2)$

Table 1: The $U(1)$ charges for the 11 currents from the equation (2.39). The first four currents with positive ($k > 4$) $U(1)$ charges have their conjugated currents with each opposite (negative) $U(1)$ charge.

For the affine currents, $K^m(z)$ where $m = 1, 2, \cdots, 6$ has vanishing $U(1)$ charge (and their complex conjugated ones also). The $J^a(z)$ where $a = 7, 8, \cdots, 14$ has the $U(1)$ charge $\frac{k}{(6+k)}$

while the complex conjugated ones has the $U(1)$ charge $-\frac{k}{(6+k)}$. For the spin-1 currents, $\overline{D}K^m(z)$with $m = 1, 2, \cdots, 5$ has vanishing $U(1)$ charge and $DK^{\bar{m}}(z)$with $\bar{m} = \bar{1}, \bar{2}, \cdots, \bar{5}$ has also vanishing $U(1)$ charge.

Summarizing this subsection, the large $\mathcal{N} = 4$ nonlinear superconformal algebra is constructed. There are six spin-1 currents with (2.30) and (2.32), four spin-$\frac{3}{2}$ currents with (2.37) and spin-2 stress energy tensor (2.28). The three spin-1 currents corresponding to $SU(2)_{\hat{k}^+}$ affine algebra live in the second subgroup $SU(2)$ of Wolf space coset and the other three spin-1 currents corresponding to $SU(2)_{\hat{k}^-}$ affine algebra live in the first subgroup $SU(2)$ of Wolf space coset. Compared to the linear version of previous subsection, the quadratic terms in $A_i(z)$ are disappeared and also the cubic terms in $G_a(z)$ are disappeared. Furthermore, some of the quadratic terms in $B_i(z)$ are disappeared [20].

3 The 16 higher spin currents in the orthogonal Wolf space coset $\frac{SO(8)}{SO(4)\times SU(2)\times SU(2)}$

This section will consider the particular supersymmetric Wolf space coset minimal model (2.1) and the following higher spin currents introduced in section 1 will be determined:

$$\left(2, \frac{5}{2}, \frac{5}{2}, 3\right) \quad : \quad (T^{(2)}, T_+^{(\frac{5}{2})}, T_-^{(\frac{5}{2})}, T^{(3)}),$$

[20] The large $\mathcal{N} = 4$ nonlinear superconformal algebra [28, 29, 30, 31] is

$$\begin{aligned}
\hat{T}(z)\,\hat{T}(w) &= \frac{1}{(z-w)^4}\,\frac{\hat{c}}{2} + \frac{1}{(z-w)^2}\,2\hat{T}(w) + \frac{1}{(z-w)}\,\partial\hat{T}(w) + \cdots, \\
\hat{T}(z)\,\Phi(w) &= \frac{1}{(z-w)^2}\,h\,\Phi(w) + \frac{1}{(z-w)}\,\partial\Phi(w) + \cdots, \\
\hat{G}_a(z)\,\hat{G}_b(w) &= \frac{1}{(z-w)^3}\,\frac{2}{3}c_{\rm Wolf}\,\hat{\delta}_{ab} - \frac{1}{(z-w)^2}\,\frac{16}{(6+k)}\left[\hat{k}^-\,\alpha_{ab}^{+i}\,\hat{A}_i + \hat{k}^+\,\alpha_{ab}^{-i}\,\hat{B}_i\right](w) \\
&+ \frac{1}{(z-w)}\left[2\hat{T}\,\hat{\delta}_{ab} - \frac{1}{2}\,\frac{16}{(6+k)}(\hat{k}^-\,\alpha_{ab}^{+i}\,\partial\hat{A}_i + \hat{k}^+\,\alpha_{ab}^{-i}\,\partial\hat{B}_i) + \right. \\
&- \left.\frac{16}{(6+k)}(\alpha^{+i}\,\hat{A}_i - \alpha^{-i}\,\hat{B}_i)_{c(a}(\alpha^{+j}\,\hat{A}_j - \alpha^{-j}\,\hat{B}_j)_{b)}^{c}\right](w) + \cdots, \\
\hat{A}_i(z)\,\hat{G}_a(w) &= \frac{1}{(z-w)}\,\alpha_{ab}^{+i}\,\hat{G}^b(w) + \cdots, \qquad \hat{B}_i(z)\,\hat{G}_a(w) = \frac{1}{(z-w)}\,\alpha_{ab}^{-i}\,\hat{G}^b(w) + \cdots, \\
\hat{A}_i(z)\,\hat{A}_j(w) &= -\frac{1}{(z-w)^2}\,\frac{1}{2}\hat{k}^+\,\hat{\delta}_{ij} + \frac{1}{(z-w)}\,\epsilon_{ijk}\,\hat{A}_k(w) + \cdots, \\
\hat{B}_i(z)\,\hat{B}_j(w) &= -\frac{1}{(z-w)^2}\,\frac{1}{2}\,\hat{k}^-\,\hat{\delta}_{ij} + \frac{1}{(z-w)}\,\epsilon_{ijk}\,\hat{B}_k(w) + \cdots,
\end{aligned}$$

where $\hat{c}$ is given by (2.29), $c_{\rm Wolf}$ is given by (2.4), $h = \frac{3}{2}$ for $\Phi = \hat{G}_a$, $h = 1$ for $\Phi = \hat{A}_i, \hat{B}_i$ and $\hat{\delta}_{ab}$ is normalized to be 1.

$$
\begin{aligned}
\left(\frac{5}{2},3,3,\frac{7}{2}\right) &: (U^{(\frac{5}{2})},U^{(3)}_{+},U^{(3)}_{-},U^{(\frac{7}{2})}),\\
\left(\frac{5}{2},3,3,\frac{7}{2}\right) &: (V^{(\frac{5}{2})},V^{(3)}_{+},V^{(3)}_{-},V^{(\frac{7}{2})}),\\
\left(3,\frac{7}{2},\frac{7}{2},4\right) &: (W^{(3)},W^{(\frac{7}{2})}_{+},W^{(\frac{7}{2})}_{-},W^{(4)}).
\end{aligned}
\tag{3.1}
$$

The highest spin-4 current is located at the last component of the last $\mathcal{N}=2$ multiplet. One can interpret the other 15 currents as the $\mathcal{N}=4$ superpartner of this spin-4 current. As in the unitary case [20], it is important to obtain the lowest higher spin-2 current in the beginning. Then all the other component fields can be determined using the currents of large $\mathcal{N}=4$ nonlinear superconformal algebra. Note that the field contents of (3.1) appear in the unitary case [1, 32].

3.1 Construction of higher spin currents of spins $\left(2,\frac{5}{2},\frac{5}{2},3\right)$

The 28 $\mathcal{N}=2$ WZW affine currents of spin-$\frac{1}{2}$ are given by $K^m(z)$, $K^{\bar{m}}(z)$, $J^a(z)$ and $J^{\bar{a}}(z)$. Let us denote them as $Q^A(z)$ and $Q^{\bar{A}}(z)$ where $A=(m,a)$ and $\bar{A}=(\bar{m},\bar{a})$. Now let us determine the spin-2 current in (3.1). We write down the most general spin-2 currents (with 14596 coefficient functions), which depend on the level k, as

$$
\begin{aligned}
T^{(2)}(z) &= \sum_{A>B=1}^{14}\sum_{\bar{C}>\bar{D}=\bar{1}}^{\bar{14}} c_{A,B,\bar{C},\bar{D}}Q^AQ^BQ^{\bar{C}}Q^{\bar{D}}(z)\\
&+ \sum_{A,C=1}^{14}\sum_{\bar{B}=\bar{1}}^{\bar{14}} c_{A,\bar{B},C}Q^AQ^{\bar{B}}\overline{D}Q^C(z)+\sum_{A=1}^{14}\sum_{\bar{B},\bar{C}=\bar{1}}^{\bar{14}} c_{A,\bar{B},\bar{C}}Q^AQ^{\bar{B}}DQ^{\bar{C}}(z)\\
&+ \sum_{A=1}^{14}\sum_{\bar{B}=\bar{1}}^{\bar{14}} c_{A,\bar{B}}\partial Q^AQ^{\bar{B}}(z)+\sum_{A=1}^{14}\sum_{\bar{B}=\bar{1}}^{\bar{14}} c'_{A,\bar{B}}Q^A\partial Q^{\bar{B}}(z)\\
&+ \sum_{A,B=1}^{14} c_{A,B}\overline{D}Q^A\overline{D}Q^B(z)+\sum_{A=1}^{14}\sum_{\bar{B}=\bar{1}}^{\bar{14}} c''_{A,\bar{B}}\overline{D}Q^ADQ^{\bar{B}}(z)\\
&+ \sum_{\bar{A},\bar{B}=\bar{1}}^{\bar{14}} c_{\bar{A},\bar{B}}DQ^{\bar{A}}DQ^{\bar{B}}(z)+\sum_{A=1}^{14} c_A\partial\overline{D}Q^A(z)+\sum_{\bar{A}=\bar{1}}^{\bar{14}} c_{\bar{A}}\partial DQ^{\bar{A}}(z).
\end{aligned}
\tag{3.2}
$$

The above quartic, cubic, quadratic and linear terms have the equal number of unbarred quantity and barred quantity. We would like to determine the coefficient functions appearing in (3.2) explicitly. Because the regularity conditions between the spin-1 current $U(z)$ (and the spin-$\frac{1}{2}$ currents $F_a(z)$) and the spin-2 current $T^{(2)}(w)$ are preserved in this extended nonlinear algebra, the following relations, together with (2.20), (2.16), (2.14), (2.12), (2.17) and (3.2), should be satisfied

$$
U(z)\,T^{(2)}(w) = +\cdots, \qquad F_a(z)\,T^{(2)}(w)=+\cdots, \qquad a=11,12,21,22. \tag{3.3}
$$

Then after (3.3) the remaining undetermined coefficient functions can be fixed by the following primary field condition under the stress tensor (2.28)

$$\hat{T}(z)\,T^{(2)}(w) \;=\; \frac{1}{(z-w)^2}\,2T^{(2)}(w) + \frac{1}{(z-w)}\,\partial T^{(2)}(w) + \cdots . \tag{3.4}$$

All the first-order singular terms between the above spin-2 current and other six spin-1 currents can be obtained from the defining equations (2.30), (2.32) and (3.2) and by requiring that the commutators between the zero mode $T_0^{(2)}$ and six spin-1 currents should vanish [1]

$$T^{(2)}(z)\,\hat{A}_i(w)|_{\frac{1}{(z-w)}} \;=\; 0, \qquad T^{(2)}(z)\,\hat{B}_i(w)|_{\frac{1}{(z-w)}} = 0. \tag{3.5}$$

All the remaining coefficient functions, after using (3.4) and (3.5), are completely determined except overall constants [21]. It turns out that the lowest higher spin-2 current (3.2) can be obtained as follows :

$$\begin{aligned} T^{(2)}(z) \;&=\; \frac{2i}{(6+k)}\Big[K^1J^8K^{\bar{1}}J^{\bar{8}} + K^1J^8K^{\bar{2}}J^{\bar{7}} + K^1J^{10}K^{\bar{1}}J^{\bar{10}} + K^1J^{12}K^{\bar{1}}J^{\bar{12}}\Big](z) \\ &+\; \text{other 341 terms}, \end{aligned} \tag{3.6}$$

where each term has vanishing $U(1)$ charge.

Let us calculate the OPE between the spin-$\frac{3}{2}$ current $\hat{G}_{21}(z)$ given in (2.36) and the spin-2 current $T^{(2)}(w)$ given in (3.6). This particular OPE is motivated by the fact that the OPE between the second component of $\mathcal{N}=2$ super stress tensor and the first component of any $\mathcal{N}=2$ super primary current provides its second component. It turns out that

$$\hat{G}_{21}(z)\,T^{(2)}(w) \;=\; \frac{1}{(z-w)}T_+^{(\frac{5}{2})}(w) + \cdots, \tag{3.7}$$

[21] After computing the OPE $T^{(2)}(z)\,T^{(2)}(w)$, one finds that three undetermined coefficients are fixed and, with an appropriate normalization constant, one has

$$\begin{aligned} T^{(2)}(z)\,T^{(2)}(w) \;&=\; -\frac{1}{(z-w)^4}\left[\frac{8k(6+k)^3(4+5k)(16+11k)}{(2+k)(8+k)}\right] \\ &-\; \frac{1}{(z-w)^2}\left[\frac{36ik(6+k)^2}{(8+k)}T^{(2)} + \frac{16(6+k)^4(4+5k)}{(8+k)}\hat{T} + \frac{16(6+k)^4(4+5k)}{(2+k)(8+k)}\hat{A}_3\hat{A}_3\right. \\ &+\; \frac{16(6+k)^4(4+5k)}{(2+k)(8+k)}\hat{A}_+\hat{A}_- + \frac{8(6+k)^4(4+5k)}{3(8+k)}\hat{B}_3\hat{B}_3 + \frac{8(6+k)^4(4+5k)}{3(8+k)}\hat{B}_+\hat{B}_- \\ &+\; \left.\frac{16i(6+k)^4(4+5k)}{(2+k)(8+k)}\partial\hat{A}_3 + \frac{8i(6+k)^4(4+5k)}{3(8+k)}\partial\hat{B}_3\right](w) + \frac{1}{(z-w)}\frac{1}{2}\partial\{T^{(2)}\,T^{(2)}\}_{-2}(w) + \cdots . \end{aligned}$$

There exist other six different solutions for $T^{(2)}(z)$. In other words, these six different expressions for $T^{(2)}(z)$ are independent each other. All the OPEs in Appendix C and D do not depend on the explicit seven different forms for $T^{(2)}(z)$.

where the spin-$\frac{5}{2}$ current is given by

$$\begin{aligned} T_{+}^{(\frac{5}{2})}(z) &= -\frac{2i\sqrt{2}(16+11k)}{(2+k)(6+k)(8+k)}\Big[K^1K^4J^8K^{\bar{4}}J^{\bar{11}}+K^1K^4J^{10}K^{\bar{4}}J^{\bar{13}} \\ &- K^1K^4J^{12}K^{\bar{4}}J^{\bar{7}}-K^1K^4J^{14}K^{\bar{4}}J^{\bar{9}}\Big](z)+\text{other 636 terms.} \end{aligned} \tag{3.8}$$

This is a primary field under the stress energy tensor (2.28). One can check that both sides in (3.7) have correct $U(1)$ charge defined in (2.39). It turns out that the corresponding $U(1)$ charge of $T^{(2)}(w)$ is equal to zero and the $U(1)$ charge of $T_{+}^{(\frac{5}{2})}(w)$ is given by the one for $\hat{G}_{21}(z)$, $\frac{(4+k)}{(6+k)}$. We will come to this issue in subsection 3.6 later. The OPE (3.7) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\,P^{(2)}(w)$ and there are many composite fields in the first-order pole, for example. According to Table 4, there are also other five composite fields satisfying the above $U(1)$ charge but they do not appear in the first order pole in (3.7).

Recall that the OPE between the third component of $\mathcal{N}=2$ super stress tensor and the first component of any $\mathcal{N}=2$ super primary current provides its third component. Then one describes the following OPE between the spin-$\frac{3}{2}$ current $\hat{G}_{12}(z)$ in (2.35) and the spin-2 current $T^{(2)}(w)$ in (3.6)

$$\hat{G}_{12}(z)\,T^{(2)}(w) = \frac{1}{(z-w)}T_{-}^{(\frac{5}{2})}(w)+\cdots. \tag{3.9}$$

In the right hand side of (3.9), the spin-$\frac{5}{2}$ current appears as follows:

$$\begin{aligned} T_{-}^{(\frac{5}{2})}(z) &= \frac{2i\sqrt{2}(16+11k)}{(2+k)(6+k)(8+k)}\Big[K^4J^7K^{\bar{1}}K^{\bar{4}}J^{\bar{12}}+K^4J^8K^{\bar{2}}K^{\bar{4}}J^{\bar{12}} \\ &+ K^4J^9K^{\bar{1}}K^{\bar{4}}J^{\bar{14}}-K^4J^{11}K^{\bar{1}}K^{\bar{4}}J^{\bar{8}}\Big](z)+\text{other 643 terms,} \end{aligned} \tag{3.10}$$

which transforms as a primary field under the stress energy tensor (2.28). The $U(1)$ charge conservation implies that the $U(1)$ charge of this spin-$\frac{5}{2}$ current is given by $-\frac{(4+k)}{(6+k)}$ which is the $U(1)$ charge of $\hat{G}_{12}(z)$ in the left hand side of (3.9). In this case also, the OPE (3.9) has simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{12}(z)\,P^{(2)}(w)$ as before. Moreover, there are no other five composite fields in (3.9).

Once again, the OPE between the second component of $\mathcal{N}=2$ super stress tensor and the third component of any $\mathcal{N}=2$ super primary current provides its first and fourth components. It turns out that the OPE between the spin-$\frac{3}{2}$ current $\hat{G}_{21}(z)$ and the spin-$\frac{5}{2}$ current $T_{-}^{(\frac{5}{2})}(w)$ (3.10) reads as

$$\hat{G}_{21}(z)\,T_{-}^{(\frac{5}{2})}(w) = \frac{1}{(z-w)^2}4T^{(2)}(w)+\frac{1}{(z-w)}\left[\frac{1}{4}4\partial T^{(2)}+T^{(3)}\right](w)+\cdots. \tag{3.11}$$

In the second-order pole of (3.11), one sees that the first component of the first $\mathcal{N}=2$ multiplet in (3.1) and, in the first-order pole, the following spin-3 current occurs

$$\begin{aligned} T^{(3)}(z) &= \frac{2i\sqrt{2}(16+11k)}{(2+k)(6+k)^2(8+k)}\Big[K^{1}K^{4}J^{8}K^{\bar{4}}K^{\bar{5}}J^{\bar{11}}-K^{1}K^{4}J^{10}K^{\bar{3}}K^{\bar{4}}J^{\bar{13}} \\ &+ K^{1}K^{4}J^{12}K^{\bar{3}}K^{\bar{4}}J^{\bar{7}}-K^{1}K^{4}J^{12}K^{\bar{4}}K^{\bar{5}}J^{\bar{7}}\Big](z)+\text{other 4941 terms.} \end{aligned} \tag{3.12}$$

This is a primary field under the stress energy tensor (2.28) and has a vanishing $U(1)$ charge. The relative coefficient $\frac{1}{4}$ above is fixed by the spins of the left hand side of (3.11) and the spin of $T^{(2)}(w)$. The OPE (3.11) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\,P_{-}^{(\frac{5}{2})}(w)$ and three kinds of quasiprimary fields due to the presence of spin-1 currents with vanishing $U(1)$ charges appear in the first-order pole. In other words, there are no other composite fields (appearing in Table 5) in the first order pole of (3.11).

In this subsection, the first $\mathcal{N}=2$ multiplet given in the first one in (3.1) is determined completely.

3.2 Construction of higher spin currents of spins $\left(\frac{5}{2},3,3,\frac{7}{2}\right)$

Now let us move on the other $\mathcal{N}=2$ multiplet in (3.1). According to the behavior in the unitary case [20], the spin-$\frac{3}{2}$ current $\hat{G}_{11}(z)$ provides the second $\mathcal{N}=2$ multiplet after calculating the OPE between $\hat{G}_{11}(z)$ and the first $\mathcal{N}=2$ multiplet. One obtains, from (2.33) and (3.6), the following OPE

$$\hat{G}_{11}(z)\,T^{(2)}(w) = \frac{1}{(z-w)}U^{(\frac{5}{2})}(w)+\cdots. \tag{3.13}$$

In the right hand side of (3.13), the following spin-$\frac{5}{2}$ current, which is the lowest component of the second $\mathcal{N}=2$ multiplet, occurs

$$\begin{aligned} U^{(\frac{5}{2})}(z) &= \frac{2(1-i)}{(2+k)(6+k)}\Big[K^{1}K^{6}J^{8}K^{\bar{5}}J^{\bar{11}}-K^{1}K^{6}J^{10}K^{\bar{5}}J^{\bar{13}} \\ &- K^{1}K^{6}J^{12}K^{\bar{5}}J^{\bar{7}}+K^{1}K^{6}J^{14}K^{\bar{5}}J^{\bar{9}}\Big](z)+\text{other 540 terms.} \end{aligned} \tag{3.14}$$

This is a primary field under the stress energy tensor (2.28) and the $U(1)$ charge is given by $\frac{(-4+k)}{(6+k)}$. There are no other five composite fields in (3.13).

Let us describe the following OPE between the above spin-$\frac{3}{2}$ current $\hat{G}_{11}(z)$ (2.33) and the

next component spin-$\frac{5}{2}$ current $T_+^{(\frac{5}{2})}(w)$ (3.8) of the second $\mathcal{N}=2$ multiplet [22]

$$\hat{G}_{11}(z)\, T_+^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}\left[-U_+^{(3)}\right](w)+\cdots. \tag{3.16}$$

Here the explicit form for the spin-3 current is given by

$$\begin{aligned} U_+^{(3)}(z) \;&=\; \frac{8(16+11k)}{(2+k)(6+k)^2(8+k)}\Big[K^1K^4J^8J^{12}K^{\bar{3}}K^{\bar{5}}+K^1K^4J^8J^{13}K^{\bar{1}}K^{\bar{4}} \\ &-\; K^1K^4J^9J^{12}K^{\bar{1}}K^{\bar{4}}+K^1K^4J^{10}J^{11}K^{\bar{1}}K^{\bar{4}}\Big](z)+\text{other 2086 terms}. \end{aligned} \tag{3.17}$$

According to Table 5, there are also other composite fields with the correct $U(1)$ charge, but they do not appear in the first order pole of (3.16).

Let us describe the following OPE between the above spin-$\frac{3}{2}$ current $\hat{G}_{11}(z)$ (2.33) and the third component spin-$\frac{5}{2}$ current $T_-^{(\frac{5}{2})}(w)$ (3.10) of the second $\mathcal{N}=2$ multiplet [23]

$$\hat{G}_{11}(z)\, T_-^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}\left[-U_-^{(3)}\right](w)+\cdots, \tag{3.19}$$

where the spin-3 current in the first-order pole is given by

$$\begin{aligned} U_-^{(3)}(z) \;&=\; -\frac{2(1-i)\sqrt{2}}{(2+k)(6+k)^2}\Big[K^1K^6J^{10}K^{\bar{1}}K^{\bar{5}}J^{\bar{10}}-K^1K^6J^{10}K^{\bar{4}}K^{\bar{5}}J^{\bar{7}} \\ &+\; K^1K^6J^{12}K^{\bar{1}}K^{\bar{5}}J^{\bar{12}}+K^1K^6J^{14}K^{\bar{1}}K^{\bar{5}}J^{\bar{14}}\Big](z)+\text{other 2068 terms}. \end{aligned} \tag{3.20}$$

There are no other composite fields (with the correct $U(1)$ charge) in the first order pole of (3.19).

[22] The normalization with sign for the spin-3 current comes from the following OPE result

$$\hat{G}_{21}(z)\, U^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}U_+^{(3)}(w)+\cdots. \tag{3.15}$$

The $U(1)$ charge for the spin-3 current $U_+^{(3)}(w)$ in (3.15) is given by the sum of the $U(1)$ charge of $\hat{G}_{21}(z)$ and the one of $U^{(\frac{5}{2})}(w)$:$\frac{2k}{(6+k)}$. This spin-3 current appears in (3.16). The OPE (3.15) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\, Q^{(\frac{5}{2})}(w)$ and the first-order pole contains the quasiprimary field due to the spin-1 current.

[23] In this case the normalization for the spin-3 current comes from the following OPE

$$\hat{G}_{12}(z)\, U^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}U_-^{(3)}(w)+\cdots. \tag{3.18}$$

The $U(1)$ charge for the spin-3 current $U_-^{(3)}(w)$ in (3.18) is given by the sum of the $U(1)$ charge of $\hat{G}_{12}(z)$ and the one of $U^{(\frac{5}{2})}(w)$:$-\frac{8}{(6+k)}$. This spin-3 current appears in the OPE (3.19). The OPE (3.18) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{12}(z)\, Q^{(\frac{5}{2})}(w)$ and the first-order pole contains the quasiprimary field due to the spin-1 current.

Let us consider the following OPE between (2.36) and (3.20), as in (3.11)

$$\begin{aligned}\hat{G}_{21}(z)\,U^{(3)}_{-}(w) &= \frac{1}{(z-w)^2}\frac{2(11+3k)}{(6+k)}U^{(\frac{5}{2})}(w) \\ &+ \frac{1}{(z-w)}\left[\frac{1}{5}\frac{2(11+3k)}{(6+k)}\partial U^{(\frac{5}{2})}+U^{(\frac{7}{2})}\right](w)+\cdots. \qquad (3.21)\end{aligned}$$

The relative coefficient $\frac{1}{5}$ is fixed by the spins of $\hat{G}_{21}(z)$, $U^{(3)}_{-}(w)$ and $U^{(\frac{5}{2})}(w)$ in (3.21). The spin-$\frac{7}{2}$ current is given by

$$\begin{aligned}U^{(\frac{7}{2})}(z) &= -\frac{8\sqrt{2}(16+11k)}{(2+k)(6+k)^3(8+k)}\Big[K^1K^4K^6J^7K^{\bar{1}}K^{\bar{4}}J^{\bar{7}}+K^1K^4K^6J^8K^{\bar{3}}K^{\bar{5}}J^{\bar{9}} \\ &- K^1K^4K^6J^9K^{\bar{1}}K^{\bar{4}}J^{\bar{9}}+K^1K^4K^6J^{10}K^{\bar{1}}K^{\bar{4}}J^{\bar{10}}\Big](z)+\ 13191\ \text{terms}. \qquad (3.22)\end{aligned}$$

The $U(1)$ charge of this current is the same as the one of the spin-$\frac{5}{2}$ current $U^{(\frac{5}{2})}(z)$. The OPE (3.21) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\,Q^{(3)}_{-}(w)$ and the first-order pole contains the quasiprimary fields due to the spin-$\frac{3}{2}$ currents. In this case also, there are no other possible composite fields (appearing in Table 6) in the first order pole of (3.21).

In this subsection, the second $\mathcal{N}=2$ multiplet in (3.1) is determined completely.

3.3 Construction of higher spin currents of spins $\left(\frac{5}{2},3,3,\frac{7}{2}\right)$

Now let us move on the other $\mathcal{N}=2$ multiplet in (3.1). According to the behavior in the unitary case [20], the spin-$\frac{3}{2}$ current $\hat{G}_{22}(z)$ provides the third $\mathcal{N}=2$ multiplet after calculating the OPE between $\hat{G}_{22}(z)$ and the first $\mathcal{N}=2$ multiplet. One obtains, from (2.34) and (3.6), the following OPE

$$\hat{G}_{22}(z)\,T^{(2)}(w) = \frac{1}{(z-w)}V^{(\frac{5}{2})}(w)+\cdots. \qquad (3.23)$$

In the right hand side of (3.23), the following spin-$\frac{5}{2}$ current, which is the lowest component of the third $\mathcal{N}=2$ multiplet, occurs

$$\begin{aligned}V^{(\frac{5}{2})}(z) &= \frac{\sqrt{2}(16+11k)}{(2+k)(6+k)}\Big[K^1J^7J^{\bar{7}}J^{\bar{1}\bar{1}}J^{\bar{1}\bar{3}}-K^1J^9J^{\bar{9}}J^{\bar{1}\bar{1}}J^{\bar{1}\bar{3}} \\ &- K^1J^{11}J^{\bar{7}}J^{\bar{9}}J^{\bar{1}\bar{1}}+K^1J^{13}J^{\bar{7}}J^{\bar{9}}J^{\bar{1}\bar{3}}\Big](z)+\text{other 542 terms}. \qquad (3.24)\end{aligned}$$

This primary spin-$\frac{5}{2}$ current has the $U(1)$ charge of $\frac{(4-k)}{(6+k)}$.

Let us describe the following OPE between the above spin-$\frac{3}{2}$ current $\hat{G}_{22}(z)$ (2.34) and the third component spin-$\frac{5}{2}$ current $T_{+}^{(\frac{5}{2})}(w)$ (3.8) of the first $\mathcal{N}=2$ multiplet [24]

$$\hat{G}_{22}(z)\, T_{+}^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}\left[-V_{+}^{(3)}\right](w)+\cdots, \tag{3.26}$$

where the spin-3 current in the first-order pole of (3.26) is given by

$$\begin{aligned} V_{+}^{(3)}(z) \;&=\; \frac{2(16+11k)}{(2+k)(6+k)^2}\Big[K^1K^2J^7J^{\bar{7}}J^{\bar{9}}J^{\bar{1}\bar{1}}-K^1K^2J^8J^{\bar{8}}J^{\bar{9}}J^{\bar{1}\bar{1}} \\ &+\; K^1K^2J^{13}J^{\bar{9}}J^{\bar{1}\bar{1}}J^{\bar{1}\bar{3}}-K^1K^2J^{14}J^{\bar{9}}J^{\bar{1}\bar{1}}J^{\bar{1}\bar{4}}\Big](z)+\text{other 2114 terms}. \end{aligned} \tag{3.27}$$

Let us describe the following OPE between the above spin-$\frac{3}{2}$ current $\hat{G}_{22}(z)$ (2.34) and the next component spin-$\frac{5}{2}$ current $T_{-}^{(\frac{5}{2})}(w)$ (3.10) of the first $\mathcal{N}=2$ multiplet [25]

$$\hat{G}_{22}(z)\, T_{-}^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}\left[-V_{-}^{(3)}\right](w)+\cdots. \tag{3.29}$$

In this case, the spin-3 current is given by

$$\begin{aligned} V_{-}^{(3)}(z) \;&=\; \frac{8(16+11k)}{(2+k)(6+k)^2(8+k)}\Big[K^1K^4K^{\bar{1}}K^{\bar{4}}J^{\bar{7}}J^{\bar{1}\bar{4}}-K^1K^4K^{\bar{1}}K^{\bar{4}}J^{\bar{8}}J^{\bar{1}\bar{3}} \\ &+\; K^1K^4K^{\bar{1}}K^{\bar{4}}J^{\bar{9}}J^{\bar{1}\bar{2}}-K^1K^4K^{\bar{1}}K^{\bar{4}}J^{\bar{1}\bar{0}}J^{\bar{1}\bar{1}}\Big](z)+\text{other 2204 terms}. \end{aligned} \tag{3.30}$$

Let us consider the following OPE between (2.36) and (3.30), as in (3.11)

$$\begin{aligned} \hat{G}_{21}(z)\, V_{-}^{(3)}(w) \;&=\; \frac{1}{(z-w)^2}\frac{(30+4k)}{(6+k)}V^{(\frac{5}{2})}(w) \\ &+\; \frac{1}{(z-w)}\left[\frac{1}{5}\frac{(30+4k)}{(6+k)}\partial V^{(\frac{5}{2})}+V^{(\frac{7}{2})}\right](w)+\cdots. \end{aligned} \tag{3.31}$$

[24] The normalization with sign for the spin-3 current comes from the following OPE result

$$\hat{G}_{21}(z)\, V^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}V_{+}^{(3)}(w)+\cdots. \tag{3.25}$$

The $U(1)$ charge for the spin-3 current $V_{+}^{(3)}(w)$ in (3.25) is given by the sum of the $U(1)$ charge of $\hat{G}_{21}(z)$ and the one of $V^{(\frac{5}{2})}(w)$:$\frac{8}{(6+k)}$. The OPE (3.25) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\, R^{(\frac{5}{2})}(w)$ and the first-order pole contains the quasiprimary field due to the spin-1 current.

[25] The normalization with sign for the spin-3 current comes from the following OPE result

$$\hat{G}_{12}(z)\, V^{(\frac{5}{2})}(w) \;=\; \frac{1}{(z-w)}V_{-}^{(3)}(w)+\cdots. \tag{3.28}$$

The $U(1)$ charge for the spin-3 current $V_{-}^{(3)}(w)$ in (3.28) is given by the sum of the $U(1)$ charge of $\hat{G}_{12}(z)$ and the one of $V^{(\frac{5}{2})}(w)$:$-\frac{2k}{(6+k)}$. This spin-3 current appears in (3.29). The OPE (3.28) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{12}(z)\, R^{(\frac{5}{2})}(w)$ and the first-order pole contains the quasiprimary field due to the spin-1 current.

The relative coefficient $\frac{1}{5}$ is fixed by the spins of $\hat{G}_{21}(z)$, $V_{-}^{(3)}(w)$ and $V^{(\frac{5}{2})}(w)$ in (3.31). The spin-$\frac{7}{2}$ current is given by

$$\begin{aligned} V^{(\frac{7}{2})}(z) &= -\frac{2\sqrt{2}(16+11k)}{(2+k)(6+k)^3}\Big[K^1K^2J^7K^{\bar{1}}J^{\bar{7}}J^{\bar{8}}J^{\bar{9}}+K^1K^2J^7K^{\bar{1}}J^{\bar{7}}J^{\bar{11}}J^{\bar{14}} \\ &+ K^1K^2J^7K^{\bar{1}}J^{\bar{9}}J^{\bar{11}}J^{\bar{12}}+K^1K^2J^7K^{\bar{2}}J^{\bar{7}}J^{\bar{11}}J^{\bar{13}}\Big](z) \\ &+ \text{other 13109 terms.} \end{aligned} \tag{3.32}$$

The $U(1)$ charge of this current is the same as the one of the spin-$\frac{5}{2}$ current $V^{(\frac{5}{2})}(z)$. The OPE (3.31) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\,R_{-}^{(3)}(w)$ and the first-order pole contains the quasiprimary fields due to the spin-$\frac{3}{2}$ currents.

In this subsection, the third $\mathcal{N}=2$ multiplet in (3.1) is determined completely.

3.4 Construction of higher spin currents of spins $\left(3,\frac{7}{2},\frac{7}{2},4\right)$

Let us consider the following OPE between (2.34) and (3.14)

$$\hat{G}_{22}(z)\,U^{(\frac{5}{2})}(w) = \frac{1}{(z-w)^2}4T^{(2)}(w)+\frac{1}{(z-w)}\left[\frac{1}{4}4\partial T^{(2)}+W^{(3)}\right](w)+\cdots. \tag{3.33}$$

The spin-3 current in (3.33) is given by

$$\begin{aligned} W^{(3)}(z) &= \frac{8i(16+11k)}{(2+k)(6+k)^2(8+k)}\Big[-K^1K^4K^6K^{\bar{4}}J^{\bar{7}}J^{\bar{9}}+K^1K^4K^6K^{\bar{4}}J^{\bar{11}}J^{\bar{13}} \\ &+ K^1K^4J^9K^{\bar{1}}K^{\bar{4}}J^{\bar{9}}-K^1K^4J^{11}K^{\bar{1}}K^{\bar{4}}J^{\bar{11}}\Big](z)+\text{other 5367 terms.} \end{aligned} \tag{3.34}$$

The OPE (3.33) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{22}(z)\,Q^{(\frac{5}{2})}(w)$ and the first-order pole contains the quasiprimary fields due to the spin-1 currents. In other words, there are no other possible composite fields in (3.33).

The following OPE between (2.36) and (3.34) provides the spin-$\frac{7}{2}$ current

$$\begin{aligned} \hat{G}_{21}(z)\,W^{(3)}(w) &= \frac{1}{(z-w)^2}\frac{(4-k)}{(6+k)}T_{+}^{(\frac{5}{2})}(w) \\ &+ \frac{1}{(z-w)}\left[\frac{1}{5}\frac{(4-k)}{(6+k)}\partial T_{+}^{(\frac{5}{2})}+W_{+}^{(\frac{7}{2})}\right](w)+\cdots, \end{aligned} \tag{3.35}$$

where the spin-$\frac{7}{2}$ current is

$$\begin{aligned} W_{+}^{(\frac{7}{2})}(z) &= \frac{8i\sqrt{2}(16+11k)}{(2+k)(6+k)^3(8+k)}\Big[K^1K^2K^4J^7K^{\bar{1}}K^{\bar{4}}J^{\bar{11}}-K^1K^2K^4J^8K^{\bar{2}}K^{\bar{4}}J^{\bar{11}} \\ &- K^1K^2K^4J^{10}K^{\bar{1}}K^{\bar{4}}J^{\bar{14}}+K^1K^2K^4J^{10}K^{\bar{2}}K^{\bar{4}}J^{\bar{13}}\Big](z)+16536\text{ terms.} \end{aligned} \tag{3.36}$$

Note that there is the $(k-4)$ factor in the $T_{+}^{(\frac{5}{2})}(w)$ and its descendant field in (3.35). When $k=4$ (or $\hat{k}^{+}=\hat{k}^{-}$), these terms vanish. Also the large $\mathcal{N}=4$ nonlinear superonformal algebra with $\hat{k}^{+}=\hat{k}^{-}$ becomes the $SO(\mathcal{N}=4)$ Knizhnik-Bershadsky algebra [60, 61]. The OPE (3.35) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\,S^{(3)}(w)$ and the first-order pole contains the quasiprimary fields due to the spin-$\frac{3}{2}$ currents.

Let us describe the following OPE between (2.15) and (3.34)

$$\begin{aligned}\hat{G}_{12}(z)\,W^{(3)}(w) &= \frac{1}{(z-w)^2}\frac{(-4+k)}{(6+k)}T_{-}^{(\frac{5}{2})}(w)\\ &+ \frac{1}{(z-w)}\left[\frac{1}{5}\frac{(-4+k)}{(6+k)}\partial T_{-}^{(\frac{5}{2})}+W_{-}^{(\frac{7}{2})}\right](w)+\cdots. \end{aligned}\tag{3.37}$$

The spin-$\frac{7}{2}$ current is given by

$$\begin{aligned}W_{-}^{(\frac{7}{2})}(z) &= \frac{16i\sqrt{2}(16+11k)}{(2+k)(6+k)^3(8+k)}\Big[K^{1}K^{4}K^{6}K^{\bar{1}}K^{\bar{4}}J^{\bar{7}}J^{\bar{14}}-K^{1}K^{4}K^{6}K^{\bar{1}}K^{\bar{4}}J^{\bar{8}}J^{\bar{13}}\\ &+ K^{1}K^{4}K^{6}K^{\bar{1}}K^{\bar{4}}J^{\bar{9}}J^{\bar{12}}-K^{1}K^{4}K^{6}K^{\bar{1}}K^{\bar{4}}J^{\bar{10}}J^{\bar{11}}\Big](z)+\ 16662\text{ terms}. \end{aligned}\tag{3.38}$$

There is the $(k-4)$ factor in the $T_{-}^{(\frac{5}{2})}(w)$ and its descendant field in (3.37). The OPE (3.37) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{12}(z)\,S^{(3)}(w)$ and the first-order pole contains the quasiprimary fields due to the spin-$\frac{3}{2}$ currents.

From the following OPE between (2.36) and (3.38)

$$\begin{aligned}\hat{G}_{21}(z)\,W_{-}^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^3}\left[-\frac{48(-4+k)}{5(6+k)}T^{(2)}\right](w)\\ &+ \frac{1}{(z-w)^2}\left[-\frac{6(-4+k)}{5(6+k)}T^{(3)}+\frac{2(16+3k)}{(6+k)}W^{(3)}+\frac{16i}{(6+k)}(\hat{A}_3-\hat{B}_3)T^{(2)}\right](w)\\ &+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{21}\,W_{-}^{(\frac{7}{2})}\}_{-2}\right.\\ &- \left.\frac{144(-4+k)}{(324+179k)}(\hat{T}T^{(2)}-\frac{3}{10}\partial^2T^{(2)})+W^{(4)}\right](w)+\cdots, \end{aligned}\tag{3.39}$$

one obtains, from (3.39), the following spin-4 current

$$\begin{aligned}W^{(4)}(z) &= \frac{16i(16+11k)}{(2+k)(6+k)^4(8+k)}\Big[K^{1}K^{2}K^{4}J^{7}K^{\bar{1}}K^{\bar{2}}K^{\bar{4}}J^{\bar{7}}+K^{1}K^{2}K^{4}J^{8}K^{\bar{1}}K^{\bar{2}}K^{\bar{4}}J^{\bar{8}}\\ &+ 2K^{1}K^{2}K^{4}J^{10}K^{\bar{1}}K^{\bar{2}}K^{\bar{4}}J^{\bar{10}}-2K^{1}K^{2}K^{4}J^{12}K^{\bar{1}}K^{\bar{2}}K^{\bar{4}}J^{\bar{12}}\Big](z)\\ &+ \text{other 98358 terms}. \end{aligned}\tag{3.40}$$

There is no descendant field for the spin-2 field (appearing in the third order pole) in the second order pole in (3.39) because the spin difference of the left hand side is given by $\frac{3}{2} - \frac{7}{2} = -2$ which is equal to the minus spin of the above spin-2 current. Each term in the second order pole of (3.39) is a primary field under the stress energy tensor. The spin-4 current appears in the OPEs, (D.20), (D.23), (D.31) and (D.36). Note that the quasiprimary field in the first-order pole in (3.39) has the factor $(k-4)$. The OPE (3.39) has very simple form compared to the unitary case in [32] where the corresponding OPE is given by $\hat{G}_{21}(z)\, S_{-}^{(\frac{7}{2})}(w)$ and the first-order pole contains many quasiprimary fields.

Therefore, the 16 higher spin currents are found explicitly. We have also checked that these 16 currents as well as the 11 currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra have regular OPEs with the spin $\frac{1}{2}$ currents $K^{m}(w)$ and $K^{\bar{m}}(w)$ with $m = 1, 2, 3, \bar{m} = \bar{1}, \bar{2}, \bar{3}$ (and their superpartner spin-1 currents $\overline{D}K^{m}(w)$ and $DK^{\bar{m}}(w)$). These six spin-$\frac{1}{2}$ currents (and six spin-1 currents) live in the subgroup $SO(4)$ of Wolf space coset. Furthermore, the OPEs between the above $(11+16)$ currents and the spin $\frac{1}{2}$ currents $K^{4}(w)$ and $K^{\bar{4}}(w)$ (and their superpartner spin-1 currents $\overline{D}K^{4}(w)$ and $DK^{\bar{4}}(w)$) have regular terms. These two spin-$\frac{1}{2}$ currents (and two spin-1 currents) live in the first subgroup $SU(2)$ of Wolf space coset. By construction, the remaining currents, spin-$\frac{1}{2}$ current $\frac{(1+i)}{2}(K^{5}+iK^{\bar{5}})(z)$ which is $(F_{12}-F_{21})(z)$ and its superpartner spin-1 current $\frac{(1+i)}{2\sqrt{2}}(\overline{D}K^{5}+iDK^{\bar{5}})(z)$ which is $U(z)$ (both living in the first subgroup $SU(2)$ of Wolf space coset) do not have any singular terms with the above (11+16) currents. Furthermore, the three spin-$\frac{1}{2}$ currents $\frac{(1+i)}{2}(K^{5}-iK^{\bar{5}})(z)(=(F_{12}+F_{21})(z))$, $F_{11}(z) = \frac{i}{\sqrt{2}}K^{6}(z)$ and $F_{22}(z) = -\frac{i}{\sqrt{2}}K^{\bar{6}}(z)$, which live in the second $SU(2)$ subgroup of the Wolf space coset, do not have any singular terms with the above $(11+16)$ currents [26].

3.5 The OPEs between the 11 currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra and 16 higher spin currents

In section 2, the 11 currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra were constructed and in section 3, the 16 higher spin currents were found. For the extension of large $\mathcal{N} = 4$ nonlinear superconformal algebra, one should calculate the OPEs between the 11 currents in section 2 and the 16 currents in section 3 as follows:

$$\begin{pmatrix} \hat{T} \\ \hat{A}_1, \hat{A}_2, \hat{A}_3 \\ \hat{B}_1, \hat{B}_2, \hat{B}_3 \\ \hat{G}_{11}, \hat{G}_{12}, \hat{G}_{21}, \hat{G}_{22} \end{pmatrix}(z) \begin{pmatrix} T^{(2)}, T_{+}^{(\frac{5}{2})}, T_{-}^{(\frac{5}{2})}, T^{(3)} \\ U^{(\frac{5}{2})}, U_{+}^{(3)}, U_{-}^{(3)}, U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, V_{+}^{(3)}, V_{-}^{(3)}, V^{(\frac{7}{2})} \\ W^{(3)}, W_{+}^{(\frac{7}{2})}, W_{-}^{(\frac{7}{2})}, W^{(4)} \end{pmatrix}(w). \tag{3.41}$$

[26] The three spin-1 currents, $(\overline{D}K^{5} - iDK^{\bar{5}})(z)$, $\overline{D}K^{6}(z)$ and $DK^{\bar{6}}(z)$, do have the singular terms with the above $(11+16)$ currents.

In other words, the OPEs between 11 currents in the left hand side of (3.41) and the 16 currents in the right hand side of (3.41) are needed.

3.6 $U(1)$ charges of higher spin currents

It is straightforward to calculate the various $U(1)$ charges under the $U(1)$ current introduced in the previous section. The result is as follows with (2.38):

$$\begin{aligned}
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\,T^{(2)}(w)|_{\frac{1}{(z-w)}} &= 0,\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\,T^{(\frac{5}{2})}_{\pm}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\pm\frac{(4+k)}{(6+k)}\right]T^{(\frac{5}{2})}_{\pm}(w),\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\,T^{(3)}(w)|_{\frac{1}{(z-w)}} &= 0,\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\begin{pmatrix} U^{(\frac{5}{2})}\\ V^{(\frac{5}{2})}\end{pmatrix}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\pm\frac{(-4+k)}{(6+k)}\right]\begin{pmatrix} U^{(\frac{5}{2})}\\ V^{(\frac{5}{2})}\end{pmatrix}(w),\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\begin{pmatrix} U^{(3)}_{+}\\ V^{(3)}_{-}\end{pmatrix}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\pm\frac{2k}{(6+k)}\right]\begin{pmatrix} U^{(3)}_{+}\\ V^{(3)}_{-}\end{pmatrix}(w),\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\begin{pmatrix} U^{(3)}_{-}\\ V^{(3)}_{+}\end{pmatrix}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\mp\frac{8}{(6+k)}\right]\begin{pmatrix} U^{(3)}_{-}\\ V^{(3)}_{+}\end{pmatrix}(w),\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\begin{pmatrix} U^{(\frac{7}{2})}\\ V^{(\frac{7}{2})}\end{pmatrix}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\pm\frac{(-4+k)}{(6+k)}\right]\begin{pmatrix} U^{(\frac{7}{2})}\\ V^{(\frac{7}{2})}\end{pmatrix}(w),\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\,W^{(3)}(w)|_{\frac{1}{(z-w)}} &= 0,\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\,W^{(\frac{7}{2})}_{\pm}(w)|_{\frac{1}{(z-w)}} &= \frac{1}{(z-w)}\left[\pm\frac{(4+k)}{(6+k)}\right]W^{(\frac{7}{2})}_{\pm}(w),\\
\left(-2i\gamma_A\hat{A}_3-2i\gamma_B\hat{B}_3\right)(z)\,W^{(4)}(w)|_{\frac{1}{(z-w)}} &= 0.
\end{aligned}\tag{3.42}$$

We present these $U(1)$ charges in the Table 2 explicitly [27]. Then the $U(1)$ charges of all the composite fields coming from the $(11+16)$ currents can be determined by the assignments in Table 1 and Table 2. The necessary $U(1)$ assignments in this paper are presented in Table 3-Table 10.

3.7 Structure of OPEs between the 11 currents and 16 currents

The 16 higher spin currents are primary fields under the stress energy tensor $\hat{T}(z)$ (2.28). Then the remaining nontrivial OPEs between the 11 currents and the 16 currents are given by 1) the OPEs between the six spin-1 currents and the 16 higher spin currents described in

[27] Compared to the unitary case in [20], the $U(1)$ charges behave similarly in the sense that for given $U(1)$ charge each current among 16 currents shares their $U(1)$ charges with each current among 11 currents.

Appendix C and 2) the OPEs between the four spin-$\frac{3}{2}$ currents and the 16 higher spin currents described in Appendix D. One should write down the correct terms, which preserve the right $U(1)$ charge described before via Tables 1-10, with arbitrary coefficients at the specific pole (with fixed spin) in the given OPE.

For example, in the OPEs between the spin-1 currents of large $\mathcal{N} = 4$ nonlinear algebra and the largest spin-4 higher spin current, the first-order pole contains the spin-4 fields in Appendices (C.40)-(C.45). The $U(1)$ charges of composite spin-4 fields are presented in Tables $7, 8$. The $U(1)$ charge of the left hand side of Appendix (C.40) is the sum of the $U(1)$ charge of $\hat{A}_+(z)$ and the one of $W^{(4)}(w)$. The former is given by $-\frac{8}{(6+k)}$ and the latter is given by 0 from Table 2. Therefore, the total $U(1)$ charge, $-\frac{8}{(6+k)}$, should appear in the right hand side of Appendix (C.40). Then from the first row of Table 8, there exist 50 possible composite spin-4 fields with the $U(1)$ charge $-\frac{8}{(6+k)}$. Because the left hand side of Appendix (C.40) is written in terms of $\mathcal{N} = 2$ WZW affine currents explicitly, let us subtract the above 50 terms with arbitrary k-dependent coefficients from the left hand side of Appendix (C.40). In order for these quantities to vanish, one should solve this equation. It turns out that the unknown coefficient functions can be obtained completely and they are given in Appendix (C.40) [28].

The OPEs between the spin-$\frac{3}{2}$ currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra and the above spin-4 higher spin current are presented in Appendix (D.37)-(D.40). One can have the composite spin-$\frac{9}{2}$ fields in the first-order pole of Appendix (D.37). The total $U(1)$ charge of this OPE should be preserved. The $U(1)$ charge of the left hand side is given by $\frac{(-4+k)}{(6+k)}$ from the Table 2, and the possible composite fields with this $U(1)$ charge are presented in the Table 9. The unknown coefficient functions can be obtained completely and are given in Appendix (D.37) [29].

The result from Appendices C and D shows that the right hand sides of all the OPEs contains the composite fields which can be obtained from the known $(11 + 16)$ currents. One expects that the new primary fields will appear in the OPEs between the 16 currents and themselves.

[28] Compared to the convention in [35], because the third order pole of the OPE between the OPE $\hat{A}_3(z)$ and $W^{(4)}(w)$ (C.42) (similarly the OPE between the OPE $\hat{B}_3(z)$ and $W^{(4)}(w)$ (C.45)) does not vanish, one can find other basis where these higher order poles should vanish. Their $W^{(4)}(z)$ in [35] is not a primary field (but a quasiprimary field) under the stress energy tensor. Note that the other OPEs appearing in Appendix C do not contain the third order (and higher order) poles. Then we do not need to change the basis for these remaining 15 higher spin currents.

[29] The highest order pole in Appendix D appears in the third order pole. As in [20], one can find the new basis where these singular terms vanish.

$U(1)$ charge	16 higher spin currents	11 currents in Table 1
$\frac{2k}{(6+k)}$	$U_+^{(3)}$	$\hat{B}_-$
$\frac{(4+k)}{(6+k)}$	$T_+^{(\frac{5}{2})}, \quad W_+^{(\frac{7}{2})}$	$\hat{G}_{21}$
$\frac{(-4+k)}{(6+k)}$	$U^{(\frac{5}{2})}, \quad U^{(\frac{7}{2})}$	$\hat{G}_{11}$
$\frac{8}{(6+k)}$	$V_+^{(3)}$	$\hat{A}_-$
0	$T^{(2)}, \quad T^{(3)}, \quad W^{(3)}, \quad W^{(4)}$	$\hat{A}_3, \quad \hat{B}_3, \quad \hat{T}$
$-\frac{8}{(6+k)}$	$U_-^{(3)}$	$\hat{A}_+$
$-\frac{(-4+k)}{(6+k)}$	$V^{(\frac{5}{2})}, \quad V^{(\frac{7}{2})}$	$\hat{G}_{22}$
$-\frac{(4+k)}{(6+k)}$	$T_-^{(\frac{5}{2})}, \quad W_-^{(\frac{7}{2})}$	$\hat{G}_{12}$
$-\frac{2k}{(6+k)}$	$V_-^{(3)}$	$\hat{B}_+$

Table 2: The $U(1)$ charges for the 16 currents from the equation (3.42). For convenience, the $U(1)$ charges for the 11 currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra in Table 1 are presented also.

4 Conclusions and outlook

The 16 higher spin currents given in (1.5) and 11 currents of large $\mathcal{N} = 4$ nonlinear algebra are determined explicitly in the Wolf space coset model $\frac{SO(8)}{SO(4)\times SU(2)\times SU(2)}$. Some OPEs of the extended large $\mathcal{N} = 4$ nonlinear algebra are described in Appendices C and D. As in the unitary case [20], the $(11+16)$ currents commute with the $\mathcal{N} = 2$ WZW affine currents living in the $\mathcal{N} = 1$ subgroup $SO(4) \times SU(2)$ and the bosonic second subgroup $SU(2)$. Three of four spin-$\frac{1}{2}$ currents live in the above bosonic second subgroup $SU(2)$ while one of them and spin-1 current live in the $\mathcal{N} = 1$ first subgroup $SU(2)$. The two $SU(2)$ affine algebras of large $\mathcal{N} = 4$ nonlinear algebra are embedded in the $\mathcal{N} = 4$ coset theory in a nontrivial way. That is, the $SU(2)_4$ affine algebra is realized by the Wolf space coset currents (2.32) while the $SU(2)_k$ affine algebra is realized by the currents (2.31) from the Wolf space coset subgroup.

One should determine the higher spin currents for general N [30]. This is because one should have the general N-dependence for the higher spin currents in order to calculate the various three point functions (for example, [62, 63, 64, 65]). Based on the present results for $N = 4$ case, one should find their N-generalization. For the large $\mathcal{N} = 4$ nonlinear superconformal

[30]In order to do this job, it is useful to obtain the higher spin currents for low N values $N = 5, 8, 9, 12, 13$ where the corresponding $SO(N + 4)$ group has even dimensions and one does not introduce the extra $U(1)$ factor in order to describe the $\mathcal{N} = 2$ WZW affine currents in the complex basis. One expects that the structure constants appearing all the OPEs we have discussed in this paper depend on N-dependence explicitly and one determines the structure constants for general N from the above low N values results. Furthermore, one should calculate each pole term by hand rather than the mathematica package.

$U(1)$ charge	Composite fields of spin-2
$\frac{2k}{(6+k)}$	$\hat{B}_-\hat{A}_3, \quad \hat{B}_-\hat{B}_3, \quad \partial\hat{B}_-$
$\frac{4k}{(6+k)}$	$\hat{B}_-\hat{B}_-$
$\frac{2(4+k)}{(6+k)}$	$\hat{B}_-\hat{A}_-$
$\frac{2(-4+k)}{(6+k)}$	$\hat{B}_-\hat{A}_+$
$\frac{16}{(6+k)}$	$\hat{A}_-\hat{A}_-$
$\frac{8}{(6+k)}$	$\hat{A}_-\hat{A}_3, \quad \hat{A}_-\hat{B}_3, \quad \partial\hat{A}_-$
0	$\hat{T},\ T^{(2)},\ \hat{A}_3\hat{A}_3,\ \hat{A}_3\hat{B}_3,\ \hat{A}_-\hat{A}_+,\ \hat{B}_3\hat{B}_3,\ \hat{B}_-\hat{B}_+,\ \partial\hat{A}_3,\ \partial\hat{B}_3$
$-\frac{8}{(6+k)}$	$\hat{A}_3\hat{A}_+, \quad \hat{B}_3\hat{A}_+, \quad \partial\hat{A}_+$
$-\frac{16}{(6+k)}$	$\hat{A}_+\hat{A}_+$
$-\frac{2(-4+k)}{(6+k)}$	$\hat{A}_-\hat{B}_+$
$-\frac{2(4+k)}{(6+k)}$	$\hat{A}_+\hat{B}_+$
$-\frac{4k}{(6+k)}$	$\hat{B}_+\hat{B}_+$
$-\frac{2k}{(6+k)}$	$\hat{A}_3\hat{B}_+, \quad \hat{B}_3\hat{B}_+, \quad \partial\hat{B}_+$

Table 3: The $U(1)$ charges for the spin-2 composite fields.

algebra, one can find their general N expressions in [29]. It is an open problem to construct the higher spin currents in terms of $\mathcal{N} = 2$ WZW affine currents with general N. For the unitary case [20], the spin of lowest higher spin current is given by one. On the other hands, for the orthogonal case, the lowest spin of the higher spin currents is given by two. The immediate question is why the spin-1 current (can be written in terms of spin-$\frac{1}{2}$ currents and its superpartner spin-1 currents with arbitrary tensorial coefficients) for the orthogonal Wolf space coset (for general N) is not allowed in the spin contents. Somehow the various geometrical, tensorial coefficient functions appearing in this spin-1 current satisfy nontrivial identities and it will turn out that there exists no spin-1 current because all the coefficient functions vanish. Similarly, for the candidate of spin-$\frac{3}{2}$ Casimir current one should show that the coefficient functions in front of this current vanish. However, for the unitary Wolf space coset [20], one should expect that there exists a nontrivial spin-1 current (and all the 15 higher spin currents) for general N with nontrivial tensorial coefficient functions. Once the spin-1 current for general N is found, it is straightforward to determine other higher spin currents.

As in the unitary case [20], it is an open problem to construct the next higher spin currents given in

$$\left(4, \frac{9}{2}, \frac{9}{2}, 5\right) \quad : \quad (P^{(4)}, P_+^{(\frac{9}{2})}, P_-^{(\frac{9}{2})}, P^{(5)}),$$

$U(1)$ charge	Composite fields of spin-$\frac{5}{2}$
$\frac{(4+3k)}{(6+k)}$	$\hat{B}_-\hat{G}_{21}$
$\frac{(-4+3k)}{(6+k)}$	$\hat{B}_-\hat{G}_{11}$
$\frac{(12+k)}{(6+k)}$	$\hat{G}_{21}\hat{A}_-$
$\frac{(-12+k)}{(6+k)}$	$\hat{G}_{11}\hat{A}_+$
$\frac{(4+k)}{(6+k)}$	$T_+^{(\frac{5}{2})}$, $\hat{B}_-\hat{G}_{22}$, $\hat{G}_{11}\hat{A}_-$, $\hat{G}_{21}\hat{A}_3$, $\hat{G}_{21}\hat{B}_3$, $\partial\hat{G}_{21}$
$\frac{(-4+k)}{(6+k)}$	$U^{(\frac{5}{2})}$, $\hat{B}_-\hat{G}_{12}$, $\hat{G}_{11}\hat{A}_3$, $\hat{G}_{11}\hat{B}_3$, $\hat{G}_{21}\hat{A}_+$, $\partial\hat{G}_{11}$
$-\frac{(-4+k)}{(6+k)}$	$V^{(\frac{5}{2})}$, $\hat{A}_3\hat{G}_{22}$, $\hat{A}_-\hat{G}_{12}$, $\hat{B}_3\hat{G}_{22}$, $\hat{G}_{21}\hat{B}_+$, $\partial\hat{G}_{22}$
$-\frac{(4+k)}{(6+k)}$	$T_-^{(\frac{5}{2})}$, $\hat{A}_3\hat{G}_{12}$, $\hat{A}_+\hat{G}_{22}$, $\hat{B}_3\hat{G}_{12}$, $\hat{G}_{11}\hat{B}_+$, $\partial\hat{G}_{12}$
$-\frac{(-12+k)}{(6+k)}$	$\hat{A}_-\hat{G}_{22}$
$-\frac{(12+k)}{(6+k)}$	$\hat{A}_+\hat{G}_{12}$
$-\frac{(-4+3k)}{(6+k)}$	$\hat{G}_{22}\hat{B}_+$
$-\frac{(4+3k)}{(6+k)}$	$\hat{G}_{12}\hat{B}_+$

Table 4: The $U(1)$ charges for the spin-$\frac{5}{2}$ composite fields.

$$\begin{aligned}
\left(\frac{9}{2},5,5,\frac{11}{2}\right) &: (Q^{(\frac{9}{2})},Q_+^{(5)},Q_-^{(5)},Q^{(\frac{11}{2})}),\\
\left(\frac{9}{2},5,5,\frac{11}{2}\right) &: (R^{(\frac{9}{2})},R_+^{(5)},R_-^{(5)},R^{(\frac{11}{2})}),\\
\left(5,\frac{11}{2},\frac{11}{2},6\right) &: (S^{(5)},S_+^{(\frac{11}{2})},S_-^{(\frac{11}{2})},S^{(6)}).
\end{aligned} \tag{4.1}$$

In order to see this structure, one should consider the OPEs between the higher spin currents in (3.1) and themselves. One expects that the explicit calculations on these constructions will be rather complicated because the spins become larger and the number of corresponding WZW affine currents become bigger and bigger. Even in the unitary case [32], the particular OPE between the higher spin currents has very complicated expression. For the OPE between $W^{(4)}(z)\,W^{(4)}(w)$, one should determine the possible composite fields of spin $s=7$. In this paper, we have considered the composite fields up to the spin $s=\frac{9}{2}$ and the possible composite fields with $s=5,\frac{11}{2},6,\frac{13}{2}$ and 7 should be classified according to their $U(1)$ charges.

It would be interesting to how one can obtain the higher spin algebra in AdS_3 bulk theory side. From the higher spin contents in (3.1) and (4.1), one expects that there are

$$\begin{aligned}
&\text{Six fields of spin } s=1,3,5,\cdots,(2n-1),\cdots,\\
&\text{Four fields of spin } s=\frac{3}{2},\frac{5}{2},\frac{7}{2},\cdots,(n+\frac{1}{2}),\cdots,\\
&\text{Two fields of spin } s=2,4,6,\cdots,2n,\cdots.
\end{aligned} \tag{4.2}$$

$U(1)$ charge	Composite fields of spin-3
$\frac{2k}{(6+k)}$	$U_+^{(3)}$, $\hat{A}_3\partial\hat{B}_-$, $\hat{B}_3\partial\hat{B}_-$, $\hat{B}_-T^{(2)}$, $\hat{B}_-\hat{T}$, $\hat{B}_-\hat{A}_3\hat{A}_3$, $\hat{B}_-\hat{A}_3\hat{B}_3$, $\hat{B}_-\hat{A}_-\hat{A}_+$, $\hat{B}_-\hat{B}_3\hat{B}_3$, $\hat{B}_-\hat{B}_-\hat{B}_+$, $\hat{B}_-\partial\hat{A}_3$, $\hat{B}_-\partial\hat{B}_3$, $\hat{G}_{21}\hat{G}_{11}$, $\partial^2\hat{B}_-$
$\frac{4k}{(6+k)}$	$\hat{B}_-\hat{B}_-\hat{A}_3$, $\hat{B}_-\hat{B}_-\hat{B}_3$, $\hat{B}_-\partial\hat{B}_-$
$\frac{2(4+k)}{(6+k)}$	$\hat{A}_-\partial\hat{B}_-$, $\hat{B}_-\hat{A}_-\hat{A}_3$, $\hat{B}_-\hat{A}_-\hat{B}_3$, $\hat{B}_-\partial\hat{A}_-$, $\hat{G}_{21}\hat{G}_{21}$
$\frac{2(-4+k)}{(6+k)}$	$\hat{A}_+\partial\hat{B}_-$, $\hat{B}_-\hat{A}_3\hat{A}_+$, $\hat{B}_-\hat{B}_3\hat{A}_+$, $\hat{B}_-\partial\hat{A}_+$, $\hat{G}_{11}\hat{G}_{11}$
$\frac{16}{(6+k)}$	$\hat{A}_-\hat{A}_-\hat{A}_3$, $\hat{A}_-\hat{A}_-\hat{B}_3$, $\hat{A}_-\partial\hat{A}_-$
$\frac{8}{(6+k)}$	$V_+^{(3)}$, $\hat{A}_3\partial\hat{A}_-$, $\hat{A}_-T^{(2)}$, $\hat{A}_-\hat{T}$, $\hat{A}_-\hat{A}_3\hat{A}_3$, $\hat{A}_-\hat{A}_3\hat{B}_3$, $\hat{A}_-\hat{A}_-\hat{A}_+$, $\hat{A}_-\hat{B}_3\hat{B}_3$, $\hat{A}_-\partial\hat{A}_3$, $\hat{A}_-\partial\hat{B}_3$, $\hat{B}_3\partial\hat{A}_-$, $\hat{B}_-\hat{A}_-\hat{B}_+$, $\hat{G}_{21}\hat{G}_{22}$, $\partial^2\hat{A}_-$
0	$T^{(3)}$, $W^{(3)}$, $\hat{A}_3T^{(2)}$, $\hat{A}_3\hat{T}$, $\hat{A}_3\hat{A}_3\hat{A}_3$, $\hat{A}_3\hat{A}_3\hat{B}_3$, $\hat{A}_3\hat{B}_3\hat{B}_3$, $\hat{A}_3\partial\hat{A}_3$, $\hat{A}_3\partial\hat{B}_3$, $\hat{A}_-\hat{A}_3\hat{A}_+$, $\hat{A}_-\hat{B}_3\hat{A}_+$, $\hat{A}_-\partial\hat{A}_+$, $\hat{A}_+\partial\hat{A}_-$, $\hat{B}_3T^{(2)}$, $\hat{B}_3\hat{T}$, $\hat{B}_3\hat{B}_3\hat{B}_3$, $\hat{B}_3\partial\hat{A}_3$, $\hat{B}_3\partial\hat{B}_3$, $\hat{B}_-\hat{A}_3\hat{B}_+$, $\hat{B}_-\hat{B}_3\hat{B}_+$, $\hat{B}_-\partial\hat{B}_+$, $\hat{B}_+\partial\hat{B}_-$, $\hat{G}_{11}\hat{G}_{22}$, $\hat{G}_{21}\hat{G}_{12}$, $\partial T^{(2)}$, $\partial\hat{T}$, $\partial^2\hat{A}_3$, $\partial^2\hat{B}_3$
$-\frac{8}{(6+k)}$	$U_-^{(3)}$, $\hat{A}_3\hat{A}_3\hat{A}_+$, $\hat{A}_3\hat{B}_3\hat{A}_+$, $\hat{A}_3\partial\hat{A}_+$, $\hat{A}_-\hat{A}_+\hat{A}_+$, $\hat{A}_+\partial\hat{A}_3$, $\hat{A}_+\partial\hat{B}_3$, $\hat{B}_3\hat{B}_3\hat{A}_+$, $\hat{B}_3\partial\hat{A}_+$, $\hat{B}_-\hat{A}_+\hat{B}_+$, $\hat{G}_{11}\hat{G}_{12}$, $T^{(2)}\hat{A}_+$, $\hat{T}\hat{A}_+$, $\partial^2\hat{A}_+$
$-\frac{16}{(6+k)}$	$\hat{A}_3\hat{A}_+\hat{A}_+$, $\hat{A}_+\partial\hat{A}_+$, $\hat{B}_3\hat{A}_+\hat{A}_+$
$-\frac{2(-4+k)}{(6+k)}$	$\hat{A}_-\hat{A}_3\hat{B}_+$, $\hat{A}_-\hat{B}_3\hat{B}_+$, $\hat{A}_-\partial\hat{B}_+$, $\hat{B}_+\partial\hat{A}_-$, $\hat{G}_{22}\hat{G}_{22}$
$-\frac{2(4+k)}{(6+k)}$	$\hat{A}_3\hat{A}_+\hat{B}_+$, $\hat{A}_+\partial\hat{B}_+$, $\hat{B}_3\hat{A}_+\hat{B}_+$, $\hat{B}_+\partial\hat{A}_+$, $\hat{G}_{12}\hat{G}_{12}$
$-\frac{4k}{(6+k)}$	$\hat{A}_3\hat{B}_+\hat{B}_+$, $\hat{B}_3\hat{B}_+\hat{B}_+$, $\hat{B}_+\partial\hat{B}_+$
$-\frac{2k}{(6+k)}$	$V_-^{(3)}$, $\hat{A}_3\hat{A}_3\hat{B}_+$, $\hat{A}_3\hat{B}_3\hat{B}_+$, $\hat{A}_3\partial\hat{B}_+$, $\hat{A}_-\hat{A}_+\hat{B}_+$, $\hat{B}_3\hat{B}_3\hat{B}_+$, $\hat{B}_3\partial\hat{B}_+$, $\hat{B}_-\hat{B}_+\hat{B}_+$, $\hat{B}_+\partial\hat{A}_3$, $\hat{B}_+\partial\hat{B}_3$, $\hat{G}_{22}\hat{G}_{12}$, $T^{(2)}\hat{B}_+$, $\hat{T}\hat{B}_+$, $\partial^2\hat{B}_+$

Table 5: The $U(1)$ charges for the spin-3 composite fields.

In [1], the explicit construction using the oscillator formalism [66, 67] for spin-1 and four spin-$\frac{3}{2}$ currents living in the lowest $\mathcal{N}=4$ multiplet (as well as the currents of large $\mathcal{N}=4$ superconformal algebra) was found. In the present case, it is nontrivial to obtain the spin-2 current and four spin-$\frac{5}{2}$ currents (living in (3.1)) using the method of higher spin algebra in the bulk. Furthermore, it is an open problem to construct the 16 higher spin currents we have constructed in this paper using the oscillator formalism in the context of higher spin algebra. One expects that it is nontrivial to express them in the primary basis where all the higher spin fields (4.2) are primary under the stress energy tensor.

So far, we described an extension of large $\mathcal{N}=4$ nonlinear superconformal algebra. By incorporating both the four spin-$\frac{1}{2}$ currents and spin-1 current again, it is interesting to obtain

an extension of large $\mathcal{N}=4$ linear superconformal algebra. One of the main results in [42] is that the small $\mathcal{N}=4$ superconformal algebra (spin-2 stress energy tensor, four spin-$\frac{3}{2}$ currents and three spin-1 currents) plays the important role in CFT side and AdS_3 bulk side. Then it is natural to consider the extension of small $\mathcal{N}=4$ superconformal algebra. It is known that the small $\mathcal{N}=4$ superconformal algebra can be obtained from the large $\mathcal{N}=4$ linear superconformal algebra in several ways rather than the large $\mathcal{N}=4$ nonlinear superconformal algebra. In other words, it is not straightforward to decouple the half of spin-1 currents from the OPE in the two fermionic spin-$\frac{3}{2}$ currents with nonlinear terms in the spin-1 currents.

In the classification of [4], there exists the following coset

$$\text{Wolf} \times SU(2) \times U(1) = \frac{Sp(N+2)}{Sp(N)} \times U(1). \tag{4.3}$$

It would be interesting to obtain the higher spin currents in this symplectic Wolf space coset model. Note that $Sp(2)$ is isomorphic to $SU(2)$. One expects that the spin-$\frac{1}{2}$ current and spin-1 current should live in $U(1)$ inside of $Sp(N)$ and the three spin-$\frac{1}{2}$ currents should be in $SU(2)$ subgroup of Wolf space coset. Also this $SU(2)$ group plays the role of $SU(2)_k$ affine algebra of large $\mathcal{N}=4$ nonlinear superconformal algebra. This coset (4.3) looks similar to the orthogonal coset but the $SU(2)$ in the denominator is missing. This is the reason why the $\mathcal{N}=1$ subgroup corresponding to one spin-1 current and three spin-$\frac{1}{2}$ currents should be present inside of $Sp(N)$.

Acknowledgments

We would like to thank H. Kim for discussions. This work was supported by the Mid-career Researcher Program through the National Research Foundation of Korea (NRF) grant funded by the Korean government (MEST) (No. 2012-045385/2013-056327/2014-051185). CA would like to thank the participants of the focus program of Asia Pacific Center for Theoretical Physics (APCTP) on "Liouville, Integrability and Branes (10) Focus Program at Asia-Pacific Center for Theoretical Physics", Sept. 03-14, 2014 for their feedbacks. We appreciate APCTP for its hospitality during completion of this work. CA acknowledges warm hospitality from the School of Liberal Arts (and Institute of Convergence Fundamental Studies), Seoul National University of Science and Technology.

Appendix A The $SO(8)$ generators in the complex basis (or Cartan-Weyl basis)

A Lie algebra is described by a set of generators $\{X_i\}$ and their commutation relations

$$[X_i, X_j] \;=\; f_{ij}{}^{k} X_k.$$

The number of generators is the dimension of the algebra and the constants $f_{ij}{}^{k}$ are structure constants. In the standard Cartan-Weyl basis, the commutation relations between generators are summarized as follows [68, 69, 70]:

$$\begin{aligned}
[H_i,\ H_j] &= 0, \quad H_i = H_i^{\dagger} \qquad i = 1, 2, 3 \cdots, l, \\
[H_i,\ E_\alpha] &= \alpha_i E_\alpha, \\
[E_\alpha,\ E_{-\alpha}] &= \alpha^i H_i, \\
[E_\alpha,\ E_\beta] &= N_{\alpha\beta} E_{\alpha+\beta} \text{ if } \alpha+\beta \text{ is a non zero root}, \\
&= 0 \qquad\qquad \text{if } \alpha+\beta \text{ is not a root.}
\end{aligned}$$

The H_i are elements of the Cartan generators of the Lie group (l is the rank of the algebra) and can be diagonalized simultaneously. They also satisfy $\mathrm{Tr}(H_i H_j) = k_D \delta_{ij}$, where k_D is some constant that depends on the representation and on the normalization of generators [68]. The vector $\alpha = (\alpha_1, \alpha_2, \cdots, \alpha_l)$ in the l-dimensional space is called a root. Root components can be extracted from the nonzero eigenvalues of the H_i in the adjoint representation [69]. The $E_{\pm\alpha}$ are corresponding raising and lowering operators for the weights and satisfy $E_\alpha = (E_{-\alpha})^{\dagger}$. To find the complex basis of a Lie algebra with even dimension, one just needs to construct generators $\widetilde{H}_i$ which are linear combinations of H_i and satisfy $\left[\widetilde{H}_i, \widetilde{H}_j\right] = 0$ where $i = 1, 2, 3 \cdots, l$ and one can find its complex conjugated one for each $\widetilde{H}_i$ where $i = 1, 2, \cdots, \frac{l}{2}$.

The $SO(N)$ generator can be realized as $N \times N$ matrix with the $(i,\ j)$ components as follows:

$$X_A^{ij} = X_{(ab)}^{ij} = i(\delta_a^i \delta_b^j - \delta_a^j \delta_b^i), \tag{A.1}$$

where the generators satisfy the usual commutator relation $[X_A,\ X_B] = f_{AB}^{C} X_C$ and $\mathrm{Tr}(X_A X_B) =$

$2\delta_{AB}$. From (A.1), we find the 28 generators of $SO(8)$ as follows:

$$X_1 = \begin{pmatrix} 0 & i & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_2 = \begin{pmatrix} 0 & 0 & i & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_3 = \begin{pmatrix} 0 & 0 & 0 & i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_4 = \begin{pmatrix} 0 & 0 & 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_5 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_6 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_7 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_8 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & i & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_9 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_{10} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_{11} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_{12} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_{13} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_{14} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_{15} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_{16} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_{17} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_{18} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$X_{19} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, X_{20} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$
X_{21} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&i&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&-i&0&0&0&0\\ 0&0&0&0&0&0&0&0 \end{pmatrix}, X_{22} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&i\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&-i&0&0&0&0 \end{pmatrix},
$$

$$
X_{23} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&i&0&0\\ 0&0&0&0&-i&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0 \end{pmatrix}, X_{24} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&i&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&-i&0&0&0\\ 0&0&0&0&0&0&0&0 \end{pmatrix},
$$

$$
X_{25} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&i\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&-i&0&0&0 \end{pmatrix}, X_{26} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&i&0\\ 0&0&0&0&0&-i&0&0\\ 0&0&0&0&0&0&0&0 \end{pmatrix},
$$

$$
X_{27} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&i\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&-i&0&0 \end{pmatrix}, X_{28} = \begin{pmatrix} 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&i\\ 0&0&0&0&0&0&-i&0 \end{pmatrix}. \quad \text{(A.2)}
$$

The $SO(8)$ generators in (A.2) are not expressed in the complex basis. To express them in the complex basis, one should find the Cartan-Weyl basis of the $SO(8)$. We choose X_1, X_{14}, X_{23} and X_{28} to form the Cartan subalgebra.[31] We rearrange the $SO(8)$ generators in (A.2) as follows:

$$
\widetilde{\mathbf{X}} = \{X_2, X_3, \cdots, X_{12}, X_{13}, X_{15}, \cdots, X_{21}, X_{22}, X_{24}, X_{25}, X_{26}, X_{27}, X_1, X_{14}, X_{23}, X_{28}\}.
$$

The last four components of $\widetilde{\mathbf{X}}$ form the Cartan subspace. Then we find the transformation

[31] There are 104 other possibilities to form the Cartan subspace. For example, one can use $X_1, X_{14}, X_{24}, X_{27}$ or $X_1, X_{14}, X_{25}, X_{26}$, or $X_1, X_{15}, X_{20}, X_{28}$ and so on.

matrix $S^{'}$ that diagonalizes X_1, X_{14}, X_{23} and X_{28} simultaneously. It is expressed as

$$S^{'} \;=\; \mathrm{diag}(C, C, C, C),$$

where

$$C \;=\; \frac{1}{\sqrt{2}} \begin{pmatrix} i & -i \\ 1 & 1 \end{pmatrix}.$$

Then we transform $\widetilde{\mathbf{X}}_i$ by the transformation $\widetilde{\mathbf{X}}_i \to Y_i = (S^{'})^{\dagger} \widetilde{\mathbf{X}}_i S^{'}$ and find the structure constants with Y_i. Next we find the adjoint representation of Y_i. Then we write the most general vector in the Cartan subspace as follows:

$$Z = a Y_{25}^{adj} + b Y_{26}^{adj} + c Y_{27}^{adj} + d Y_{28}^{adj}.$$

Then we find the matrix S that makes Z diagonalized by the transformation $Z^{'} = S^{\dagger} Z S$. The 28×28 matrix S is given by

$$S \;=\; \begin{pmatrix} A_1 & A_2 & 0 & 0 & 0 & 0 & 0 \\ A_3 & 0 & A_4 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & A_5 & A_6 & 0 & 0 \\ 0 & A_7 & A_8 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & A_9 & A_{10} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & A_{11} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & A_{12} \end{pmatrix}, \tag{A.3}$$

where the zeros 0 stand for 4×4 zero matrices and the nonzero 4×4 matrix elements are given by

$$A_1 \;=\; \begin{pmatrix} -\frac{i}{2} & -\frac{1}{2} & 0 & 0 \\ \frac{i}{2} & -\frac{1}{2} & 0 & 0 \\ \frac{i}{2} & \frac{1}{2} & 0 & 0 \\ -\frac{i}{2} & \frac{1}{2} & 0 & 0 \end{pmatrix} = A_5 = A_7, \quad A_2 = \begin{pmatrix} 0 & 0 & -\frac{1}{2} & \frac{i}{2} \\ 0 & 0 & -\frac{1}{2} & -\frac{i}{2} \\ 0 & 0 & -\frac{1}{2} & \frac{i}{2} \\ 0 & 0 & -\frac{1}{2} & -\frac{i}{2} \end{pmatrix} = A_8 = A_{10},$$

$$A_3 \;=\; \begin{pmatrix} 0 & 0 & -\frac{i}{2} & -\frac{1}{2} \\ 0 & 0 & \frac{i}{2} & -\frac{1}{2} \\ 0 & 0 & \frac{i}{2} & \frac{1}{2} \\ 0 & 0 & -\frac{i}{2} & \frac{1}{2} \end{pmatrix} = A_9, \qquad A_4 = \begin{pmatrix} -\frac{1}{2} & \frac{i}{2} & 0 & 0 \\ -\frac{1}{2} & -\frac{i}{2} & 0 & 0 \\ -\frac{1}{2} & \frac{i}{2} & 0 & 0 \\ -\frac{1}{2} & -\frac{i}{2} & 0 & 0 \end{pmatrix} = A_6,$$

$$A_{11} \;=\; \begin{pmatrix} -\frac{i}{2} & -\frac{1}{2} & -\frac{1}{2} & \frac{i}{2} \\ \frac{i}{2} & -\frac{1}{2} & -\frac{1}{2} & -\frac{i}{2} \\ \frac{i}{2} & \frac{1}{2} & -\frac{1}{2} & \frac{i}{2} \\ -\frac{i}{2} & \frac{1}{2} & -\frac{1}{2} & -\frac{i}{2} \end{pmatrix}, \qquad A_{12} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

The diagonalized matrix $Z^{'}$ with (A.3) is expressed as

$$Z^{'} \;=\; S^{\dagger} Z S = \mathrm{diag}(B_1, B_2, B_3, B_4, B_5, B_6, B_7),$$

where each matrix element is given by

$$\begin{aligned}
B_1 &= \mathrm{diag}(a+b, -(a+b), -a+b, -(-a+b)), \\
B_2 &= \mathrm{diag}(a+c, -(a+c), -a+c, -(-a+c)), \\
B_3 &= \mathrm{diag}(b+c, -(b+c), -b+c, -(-b+c)), \\
B_4 &= \mathrm{diag}(a+d, -(a+d), -a+d, -(-a+d)), \\
B_5 &= \mathrm{diag}(b+d, -(b+d), -b+d, -(-b+d)), \\
B_6 &= \mathrm{diag}(c+d, -(c+d), -c+d, -(-c+d)), \\
B_7 &= \mathrm{diag}(0,0,0,0).
\end{aligned} \tag{A.4}$$

From (A.4), one can find all the roots (eigenvalues) as follows [32]:

$$\begin{aligned}
\alpha_1 &= a+b \equiv (1,1,0,0), & -\alpha_1 = -a+b \equiv (-1,-1,0,0), \\
\alpha_2 &= -a+b \equiv (-1,1,0,0), & -\alpha_2 = -(-a+b) \equiv (1,-1,0,0), \\
\alpha_3 &= a+c \equiv (1,0,1,0), & -\alpha_3 = -(a+c) \equiv (-1,0,-1,0), \\
\alpha_4 &= -a+c \equiv (-1,0,1,0), & -\alpha_4 = -(-a+c) \equiv (1,0,-1,0), \\
\alpha_5 &= b+c \equiv (0,1,1,0), & -\alpha_5 = -(b+c) \equiv (0,-1,-1,0), \\
\alpha_6 &= -b+c \equiv (0,-1,1,0), & -\alpha_6 = -(-b+c) \equiv (0,1,-1,0), \\
\alpha_7 &= a+d \equiv (1,0,0,1), & -\alpha_7 = -(a+d) \equiv (-1,0,0,-1), \\
\alpha_8 &= -a+d \equiv (-1,0,0,1), & -\alpha_8 = -(-a+d) \equiv (1,0,0,-1), \\
\alpha_9 &= b+d \equiv (0,1,0,1), & -\alpha_9 = -(b+d) \equiv (0,-1,0,-1), \\
\alpha_{10} &= -b+d \equiv (0,-1,0,1), & -\alpha_{10} = -(-b+d) \equiv (0,1,0,-1), \\
\alpha_{11} &= c+d \equiv (0,0,1,1), & -\alpha_{11} = -(c+d) \equiv (0,0,-1,-1), \\
\alpha_{12} &= -c+d \equiv (0,0,-1,1), & -\alpha_{12} = -(-c+d) \equiv (0,0,1,-1), \\
\alpha_{13} &= -\alpha_{13} = \alpha_{14} = -\alpha_{14} = 0 \equiv (0,0,0,0).
\end{aligned} \tag{A.5}$$

Finally we find $E_{\pm\alpha}$ and H_i by the following linear transformation:

$$\mathbf{E} = S\,\widetilde{\mathbf{X}},$$

where the left hand side is

$$\mathbf{E} = \{E_{\alpha_1}, E_{-\alpha_1}, E_{\alpha_2}, E_{-\alpha_2}, \cdots, E_{\alpha_{12}}, E_{-\alpha_{12}}, H_1, H_2, H_3, H_4\}.$$

[32] Usually, positive roots are roots whose the first non-vanishing element is positive. In our case, positive roots are roots whose the last non-vanishing element is positive.

Then we construct the $SO(8)$ generators, for our purpose, in the complex basis as follows:

$$\begin{aligned} T_A &= \{E_{\alpha_1}, E_{\alpha_2}, \frac{1}{\sqrt{2}}(H_1 + iH_2), E_{\alpha_{12}}, \frac{1}{\sqrt{2}}(H_3 + iH_4), E_{\alpha_{11}}, E_{\alpha_3}, E_{\alpha_4} \\ & \quad E_{\alpha_9}, E_{\alpha_{10}}, E_{\alpha_5}, E_{\alpha_6}, E_{\alpha_7}, E_{\alpha_8}\}, \\ T_{\bar{A}} &= T_A^\dagger. \end{aligned}$$

Note that $T_3 = \frac{1}{\sqrt{2}}(H_1 + iH_2)$ and $T_5 = \frac{1}{\sqrt{2}}(H_3 + iH_4)$ where the four Cartan generators are given [68]

$$\begin{aligned} H_1 &= \text{diag}(1, -1, 0, 0, 0, 0, 0, 0), \\ H_2 &= \text{diag}(0, 0, 1, -1, 0, 0, 0, 0), \\ H_3 &= \text{diag}(0, 0, 0, 0, 1, -1, 0, 0), \\ H_4 &= \text{diag}(0, 0, 0, 0, 0, 0, 1, -1). \end{aligned}$$

We present the 14 generators T_A of $SO(8)$ (the other 14 generators are given by $T_{\bar{A}} = T_A^\dagger$) as follows:

$$T_1 = \left(\begin{array}{cccc|cccc} 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array}\right), T_2 = \left(\begin{array}{cccc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array}\right),$$

$$T_3 = \left(\begin{array}{cccc|cccc} \frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{i}{\sqrt{2}} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{i}{\sqrt{2}} & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array}\right), T_4 = \left(\begin{array}{cccc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array}\right),$$

$$T_5 = \left(\begin{array}{cccc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & \frac{1}{\sqrt{2}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{i}{\sqrt{2}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\frac{i}{\sqrt{2}} \end{array}\right), T_6 = \left(\begin{array}{cccc|cccc} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array}\right),$$

$$
T_7 = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&-1&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
\hline
0&1&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right), T_8 = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&0&0&0\\
0&0&0&0&0&-1&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
\hline
1&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right),
$$

$$
T_9 = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&-1\\
0&0&0&0&0&0&0&0\\
\hline
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&1&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right), T_{10} = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&-1\\
\hline
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&1&0&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right),
$$

$$
T_{11} = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&-1&0&0\\
0&0&0&0&0&0&0&0\\
\hline
0&0&0&1&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right), T_{12} = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&-1&0&0\\
\hline
0&0&1&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right),
$$

$$
T_{13} = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&0&0&-1\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
\hline
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
0&1&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right), T_{14} = \left(\begin{array}{cccc|cccc}
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&-1\\
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
\hline
0&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0\\
1&0&0&0&0&0&0&0\\
0&0&0&0&0&0&0&0
\end{array}\right). \tag{A.6}
$$

The 6 generators T_m where $m = 1, 2, \cdots, 6$ (and their conjugated ones) belong to the Wolf space subgroup $SO(4) \times SU(2) \times SU(2)$ of the $SO(8)$. Among them, the ones with $1, 2, 3$ (and conjugated ones) are located at the first 4×4 matrices and the remaining ones with $4, 5, 6$ (and conjugated ones) are in the last 4×4 matrices inside 8×8 matrices. The 8 generators T_a where $a = 7, 8, \cdots, 14$ belong to the Wolf space coset $\frac{SO(8)}{SO(4) \times SU(2) \times SU(2)}$ and are located off diagonal 4×4 matrices. Then the 8×8 matrices are classified as the first 4×4 matrices, the last 4×4 matrices and the remaining two off diagonal 4×4 matrices. For general N, the subgroup $SO(N)$ lives in the first $N \times N$ matrices and the remaining $SU(2) \times SU(2)$ lives in

the last 4×4 matrices.

The commutators between T_A, and $T_{\bar{A}}$ are given by [33]

$$
\begin{aligned}
&[T_1,\ T_{\bar{1}}] = [E_{\alpha_1},\ E_{-\alpha_1}] = H_1 + H_2 = \frac{(1-i)}{\sqrt{2}}(T_3 + iT_{\bar{3}}),\\
&[T_2,\ T_{\bar{2}}] = [E_{\alpha_2},\ E_{-\alpha_2}] = -H_1 + H_2 = -\frac{(1+i)}{\sqrt{2}}(T_3 - iT_{\bar{3}}),\\
&[T_4,\ T_{\bar{4}}] = [E_{\alpha_{12}},\ E_{-\alpha_{12}}] = -H_3 + H_4 = -\frac{(1+i)}{\sqrt{2}}(T_5 - iT_{\bar{5}}),\\
&[T_6,\ T_{\bar{6}}] = [E_{\alpha_{11}},\ E_{-\alpha_{11}}] = H_3 + H_4 = \frac{(1-i)}{\sqrt{2}}(T_5 + iT_{\bar{5}}),\\
&[T_7,\ T_{\bar{7}}] = [E_{\alpha_3},\ E_{-\alpha_3}] = H_1 + H_3 = \frac{1}{\sqrt{2}}(T_3 + T_{\bar{3}} + T_5 + T_{\bar{5}}),\\
&[T_8,\ T_{\bar{8}}] = [E_{\alpha_4},\ E_{-\alpha_4}] = -H_1 + H_3 = \frac{1}{\sqrt{2}}(-T_3 - T_{\bar{3}} + T_5 + T_{\bar{5}}),\\
&[T_9,\ T_{\bar{9}}] = [E_{\alpha_9},\ E_{-\alpha_9}] = H_2 + H_4 = -\frac{i}{\sqrt{2}}(T_3 - T_{\bar{3}} + T_5 - T_{\bar{5}}),\\
&[T_{10},\ T_{\bar{10}}] = [E_{\alpha_{10}},\ E_{-\alpha_{10}}] = -H_2 + H_4 = \frac{i}{\sqrt{2}}(T_3 - T_{\bar{3}} - T_5 + T_{\bar{5}}),\\
&[T_{11},\ T_{\bar{11}}] = [E_{\alpha_5},\ E_{-\alpha_5}] = H_2 + H_3 = \frac{1}{\sqrt{2}}(-iT_3 + iT_{\bar{3}} + T_5 + T_{\bar{5}}),\\
&[T_{12},\ T_{\bar{12}}] = [E_{\alpha_6},\ E_{-\alpha_6}] = -H_2 + H_3 = \frac{1}{\sqrt{2}}(iT_3 - iT_{\bar{3}} + T_5 + T_{\bar{5}}),\\
&[T_{13},\ T_{\bar{13}}] = [E_{\alpha_7},\ E_{-\alpha_7}] = H_1 + H_4 = \frac{1}{\sqrt{2}}(T_3 + T_{\bar{3}} - iT_5 + iT_{\bar{5}}),\\
&[T_{14},\ T_{\bar{14}}] = [E_{\alpha_8},\ E_{-\alpha_8}] = -H_1 + H_4 = \frac{1}{\sqrt{2}}(-T_3 - T_{\bar{3}} - iT_5 + iT_{\bar{5}}).
\end{aligned}
\tag{A.7}
$$

Each commutator in (A.7) represents a $SU(2)$ subgroup of $SO(8)$ and the coefficients of H_i in each algebra gives the corresponding root vector (A.5) [68]. Moreover, $T_1, T_{\bar{1}}, T_2, T_{\bar{2}}$ with $\frac{(1-i)}{\sqrt{2}}(T_3 + iT_{\bar{3}})$, and $-\frac{(1+i)}{\sqrt{2}}(T_3 - iT_{\bar{3}})$ forms the $SO(4)$ subgroup and $T_4, T_{\bar{4}}, T_6, T_{\bar{6}}$ with $-\frac{(1+i)}{\sqrt{2}}(T_5 - iT_{\bar{5}})$ and $\frac{(1-i)}{\sqrt{2}}(T_5 + iT_{\bar{5}})$ forms the $SU(2) \times SU(2)$ subgroup in the coset.

We present the nonzero structure constants of $SO(8)$ in Table 11 as follows.

[33] Other commutators can be obtained from (A.6).

Appendix B The current algebra for the $\mathcal{N}=2$ supersymmetric WZW model on a group $SO(N+4)$ in terms of components of spin $s=\frac{1}{2}$ and 1

Most of the OPEs between the component fields for the $\mathcal{N}=2$ supersymmetric WZW model on a group $SO(N+4)$ take the same forms for those OPEs of [11]. Let us describe the following OPEs which have the same behaviors to the ones in [11]

$$
\begin{aligned}
K^m(z)\,\overline{D}K^n(w) &= -\frac{1}{(z-w)} f_{\bar{m}\bar{n}}{}^{\bar{p}} K^p(w) + \cdots, \\
\overline{D}K^m(z)\,\overline{D}K^n(w) &= -\frac{1}{(z-w)} f_{\bar{m}\bar{n}}{}^{\bar{p}} \overline{D}K^p(w) + \cdots, \\
K^m(z)\,\overline{D}J^a(w) &= -\frac{1}{(z-w)} f_{\bar{m}\bar{a}}{}^{\bar{b}} J^b(w) + \cdots, \\
\overline{D}K^m(z)\,J^a(w) &= -\frac{1}{(z-w)} f_{\bar{m}\bar{a}}{}^{\bar{b}} J^b(w) + \cdots, \\
\overline{D}K^m(z)\,\overline{D}J^a(w) &= -\frac{1}{(z-w)} f_{\bar{m}\bar{a}}{}^{\bar{b}} \overline{D}J^b(w) + \cdots, \\
K^{\bar{m}}(z)\,DK^{\bar{n}}(w) &= -\frac{1}{(z-w)} f_{mn}{}^{p} K^{\bar{p}}(w) + \cdots, \\
DK^{\bar{m}}(z)\,DK^{\bar{n}}(w) &= -\frac{1}{(z-w)} f_{mn}{}^{p} DK^{\bar{p}}(w) + \cdots, \\
K^{\bar{m}}(z)\,DJ^{\bar{a}}(w) &= -\frac{1}{(z-w)} f_{ma}{}^{b} J^{\bar{b}}(w) + \cdots, \\
DK^{\bar{m}}(z)\,J^{\bar{a}}(w) &= -\frac{1}{(z-w)} f_{ma}{}^{b} J^{\bar{b}}(w) + \cdots, \\
DK^{\bar{m}}(z)\,DJ^{\bar{a}}(w) &= -\frac{1}{(z-w)} f_{ma}{}^{b} DJ^{\bar{b}}(w) + \cdots, \\
K^m(z)\,K^{\bar{n}}(w) &= -\frac{1}{(z-w)} (k+N+2)\delta^{m\bar{n}} + \cdots, \\
K^m(z)\,DK^{\bar{n}}(w) &= -\frac{1}{(z-w)} f_{\bar{m}n}{}^{\bar{p}} K^p(w) + \cdots, \\
\overline{D}K^m(z)\,K^{\bar{n}}(w) &= -\frac{1}{(z-w)} f_{\bar{m}n}{}^{p} K^{\bar{p}}(w) + \cdots, \\
\overline{D}J^a(z)\,DJ^{\bar{b}}(w) &= \frac{1}{(z-w)^2}\frac{1}{2}\left[2(k+N+2)\delta^{a\bar{b}} + f_{\bar{a}m}{}^{\bar{c}} f_{b\bar{m}}{}^{c} + f_{\bar{a}c}{}^{\bar{m}} f_{b\bar{c}}{}^{m}\right] \\
&- \frac{1}{(z-w)}\left[f_{\bar{a}b}{}^{\bar{m}}\overline{D}K^m + f_{\bar{a}b}{}^{m} DK^{\bar{m}}\right. \\
&+ \left.\frac{1}{(k+N+2)}\left(f_{\bar{a}m}{}^{\bar{c}} f_{b\bar{m}}{}^{d} J^c J^{\bar{d}} + f_{\bar{a}c}{}^{\bar{m}} f_{b\bar{c}}{}^{n} K^m K^{\bar{n}}\right)\right](w) + \cdots,
\end{aligned}
$$

$$
\begin{aligned}
J^a(z)\, J^{\bar b}(w) &= -\frac{1}{(z-w)}(k+N+2)\delta^{a\bar b}+\cdots,\\
J^a(z)\, DJ^{\bar b}(w) &= -\frac{1}{(z-w)} f_{\bar a b}{}^{\bar m} K^m(w)+\cdots,\\
\overline{D}J^a(z)\, J^{\bar b}(w) &= -\frac{1}{(z-w)} f_{\bar a b}{}^{m} K^{\bar m}(w)+\cdots,\\
\overline{D}K^m(z)\, J^{\bar a}(w) &= -\frac{1}{(z-w)} f_{\bar m a}{}^{b} J^{\bar b}(w)+\cdots,\\
J^a(z)\, DK^{\bar m}(w) &= -\frac{1}{(z-w)} f_{\bar a m}{}^{\bar b} J^{b}(w)+\cdots.
\end{aligned}
\tag{B.1}
$$

The following OPEs have different behaviors from the ones in [11] because, for the present case, the nonzero structure constants $f_{ab}{}^{m}$ and $f_{\bar a\bar b}{}^{\bar m}$ occur

$$
\begin{aligned}
\overline{D}K^m(z)\, DK^{\bar n}(w) &= \frac{1}{(z-w)^2}\frac{1}{2}\left[2(k+N+2)\delta^{m\bar n}+f_{\bar m p}{}^{\bar q} f_{n\bar p}{}^{q}+f_{\bar m a}{}^{\bar b} f_{n\bar a}{}^{b}\right]\\
&- \frac{1}{(z-w)}\Big[f_{\bar m n}{}^{p} DK^{\bar p}+f_{\bar m n}{}^{\bar p}\overline{D}K^{p}\\
&+ \frac{1}{(k+N+2)}(f_{\bar m p}{}^{\bar q} f_{n\bar p}{}^{r} K^q K^{\bar r}+f_{\bar m a}{}^{\bar b} f_{n\bar a}{}^{c} J^b J^{\bar c})\Big](w)+\cdots,\\
\overline{D}K^m(z)\, DJ^{\bar a}(w) &= -\frac{1}{(z-w)}\Big[f_{\bar m a}{}^{\bar b}\overline{D}J^{b}+f_{\bar m a}{}^{b} DJ^{\bar b}\\
&+ \frac{1}{(k+N+2)}(f_{\bar m p}{}^{\bar q} f_{a\bar p}{}^{b} K^q J^{\bar b}+f_{\bar m b}{}^{\bar c} f_{a\bar b}{}^{p} J^c K^{\bar p})\Big](w)+\cdots,\\
\overline{D}J^a(z)\, DK^{\bar m}(w) &= -\frac{1}{(z-w)}\Big[f_{\bar a m}{}^{\bar b}\overline{D}J^{b}+f_{\bar a m}{}^{b} DJ^{\bar b}\\
&+ \frac{1}{(k+N+2)}(f_{\bar a b}{}^{\bar n} f_{m\bar b}{}^{c} K^n J^{\bar c}+f_{\bar a p}{}^{\bar b} f_{m\bar p}{}^{n} J^b K^{\bar n})\Big](w)+\cdots,\\
K^m(z)\, DJ^{\bar a}(w) &= -\frac{1}{(z-w)} f_{\bar m a}{}^{\bar b} J^{b}(w)+\cdots,\\
\overline{D}J^a(z)\, \overline{D}J^{b}(w) &= -\frac{1}{(z-w)} f_{\bar a\bar b}{}^{\bar m}\overline{D}K^{m}(w)+\cdots,\\
\overline{D}J^a(z)\, K^{\bar m}(w) &= -\frac{1}{(z-w)} f_{\bar a m}{}^{b} J^{\bar b}(w)+\cdots,\\
J^{\bar a}(z)\, DJ^{\bar b}(w) &= -\frac{1}{(z-w)} f_{ab}{}^{m} K^{\bar m}(w)+\cdots,\\
DJ^{\bar a}(z)\, DJ^{\bar b}(w) &= -\frac{1}{(z-w)} f_{ab}{}^{m} DK^{\bar m}(w)+\cdots.
\end{aligned}
\tag{B.2}
$$

For simplicity, the notation $|_{\theta=\bar\theta=0}$ acting on the $\mathcal{N}=2$ superfields [13] are ignored in (B.1) and (B.2). That is, $DK^{\bar m}|_{\theta=\bar\theta=0}(z)\equiv DK^{\bar m}(z), \overline{D}K^m|_{\theta=\bar\theta=0}(z)\equiv \overline{D}K^m(z), DJ^{\bar a}|_{\theta=\bar\theta=0}(z)\equiv DJ^{\bar a}(z)$, and $\overline{D}J^a|_{\theta=\bar\theta=0}(z)\equiv\overline{D}J^a(z)$.

Therefore, the complete OPEs between the WZW affine currents with $s = \frac{1}{2}, 1$ are summarized by (B.1) and (B.2) [34].

Appendix C The OPEs between the six spin-1 currents and the higher spin currents

In this section, we present the OPEs between the six spin-1 currents of large $\mathcal{N} = 4$ nonlinear superconformal algebra in section 2 and the 16 higher spin currents in section 3.

Appendix C.1 The OPEs between six spin-1 currents and the higher spin current of spins $(2, \frac{5}{2}, \frac{5}{2}, 3)$

The nontrivial OPEs between six spin-1 currents (2.30) and (2.32) and the higher spin current $T_+^{(\frac{5}{2})}(w)$ (3.8) are

$$\begin{pmatrix} \hat{A}_+ \\ \hat{B}_+ \end{pmatrix}(z)\, T_+^{(\frac{5}{2})}(w) = -\frac{1}{(z-w)}\, i \begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(w) + \cdots, \tag{C.1}$$

$$\hat{A}_3(z)\, T_+^{(\frac{5}{2})}(w) = \frac{1}{(z-w)}\, \frac{i}{2}\, T_+^{(\frac{5}{2})}(w) + \cdots = \hat{B}_3(z)\, T_+^{(\frac{5}{2})}(w). \tag{C.2}$$

One can check the $U(1)$ charge conservation in these OPEs from Table 2 or 4. There are no nonlinear terms in (C.1) or (C.2).

The nontrivial OPEs between six spin-1 currents (2.30) and (2.32) and the higher spin current $T_-^{(\frac{5}{2})}(w)$ (3.10) as follows:

$$\begin{pmatrix} \hat{A}_- \\ \hat{B}_- \end{pmatrix}(z)\, T_-^{(\frac{5}{2})}(w) = \frac{1}{(z-w)}\, i \begin{pmatrix} V^{(\frac{5}{2})} \\ U^{(\frac{5}{2})} \end{pmatrix}(w) + \cdots, \tag{C.3}$$

[34] Other spin-1 currents are not independent because of the following constraints. The nonlinear constraints, by taking $\theta, \bar{\theta}$ independent terms [17, 11], are given by

$$\begin{aligned}
DK^m|_{\theta=\bar{\theta}=0}(z) &= -\frac{1}{2(k+N+2)}(f_{\bar{m}n}{}^{\bar{p}} K^n K^p + f_{\bar{m}a}{}^{\bar{b}} J^a J^b)|_{\theta=\bar{\theta}=0}(z), \\
DJ^a|_{\theta=\bar{\theta}=0}(z) &= -\frac{1}{(k+N+2)} f_{\bar{a}b}{}^{\bar{m}} (J^b K^m)|_{\theta=\bar{\theta}=0}(z), \\
\overline{D}K^{\bar{m}}|_{\theta=\bar{\theta}=0}(z) &= -\frac{1}{2(k+N+2)}(f_{m\bar{n}}{}^{p} K^{\bar{n}} K^{\bar{p}} + f_{m\bar{a}}{}^{b} J^{\bar{a}} J^{\bar{b}})|_{\theta=\bar{\theta}=0}(z), \\
\overline{D}J^{\bar{a}}|_{\theta=\bar{\theta}=0}(z) &= -\frac{1}{(k+N+2)} f_{a\bar{b}}{}^{m} (J^{\bar{b}} K^{\bar{m}})|_{\theta=\bar{\theta}=0}(z).
\end{aligned} \tag{B.3}$$

Compared to the unitary case, there exist the extra terms in the right hand side of first and third terms of (B.3) because the structure constants $f_{ab}{}^{m}$ and $f_{\bar{a}\bar{b}}{}^{\bar{m}}$ are nonvanishing from Appendix A. Furthermore, the spin-$\frac{3}{2}$ currents, the last components, can be expressed in terms of known independent spin-$\frac{1}{2}$ currents (and their spin-1 superpartners) as in footnote 12 of [20].

$$\hat{A}_3(z)\,T_-^{(\frac{5}{2})}(w) \;=\; -\frac{1}{(z-w)}\,\frac{i}{2}\,T_-^{(\frac{5}{2})}(w)+\cdots=\hat{B}_3(z)\,T_-^{(\frac{5}{2})}(w). \tag{C.4}$$

The $U(1)$ charge conservation in these OPEs can be checked from Table 2 or 4 also. No nonlinear terms in (C.3) or (C.4).

The OPEs between six spin-1 currents (2.30) and (2.32) and the higher spin current $T^{(3)}(w)$ (3.12) are as follows:

$$\hat{A}_\pm(z)\,T^{(3)}(w) \;=\; \frac{1}{(z-w)}\,i\begin{pmatrix} U_-^{(3)} \\ V_+^{(3)} \end{pmatrix}(w)+\cdots, \tag{C.5}$$

$$\hat{A}_3(z)\,T^{(3)}(w) \;=\; \frac{1}{(z-w)^2}\,2iT^{(2)}(w)+\cdots=\hat{B}_3(z)\,T^{(3)}(w), \tag{C.6}$$

$$\hat{B}_\pm(z)\,T^{(3)}(w) \;=\; \frac{1}{(z-w)}\,i\begin{pmatrix} V_-^{(3)} \\ U_+^{(3)} \end{pmatrix}(w)+\cdots. \tag{C.7}$$

The $U(1)$ charge conservation in these OPEs are checked by Table 2 or 5. There are no nonlinear terms in (C.5), (C.6) or (C.7). There is no descendant field for the spin-2 field in the first order pole of (C.6).

Therefore, one obtains the OPEs between six spin-1 currents and the higher spin current of spins $(2,\frac{5}{2},\frac{5}{2},3)$ explicitly. There are no nonlinear terms in the right hand side of these OPEs. Some of the higher spin currents corresponding to three $\mathcal{N}=2$ multiplets (3.1) occur in the right hand sides of the OPEs.

Appendix C.2 The OPEs between six spin 1 currents and two higher spin currents of spins $(\frac{5}{2},3,3,\frac{7}{2})$

One has the following nontrivial OPEs between the spin-1 currents and the spin-$\frac{5}{2}$ currents $U^{(\frac{5}{2})}(w)$ (3.14) and $V^{(\frac{5}{2})}(w)$ (3.24):

$$\hat{A}_\mp(z)\begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(w) \;=\; \frac{1}{(z-w)}\,i\begin{pmatrix} -T_+^{(\frac{5}{2})} \\ T_-^{(\frac{5}{2})} \end{pmatrix}(w)+\cdots, \tag{C.8}$$

$$\hat{A}_3(z)\begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(w) \;=\; \frac{1}{(z-w)}\,\frac{i}{2}\begin{pmatrix} -U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(w)+\cdots=-\hat{B}_3(z)\begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(w), \tag{C.9}$$

$$\hat{B}_\pm(z)\begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(w) \;=\; \frac{1}{(z-w)}\,i\begin{pmatrix} T_-^{(\frac{5}{2})} \\ -T_+^{(\frac{5}{2})} \end{pmatrix}(w)+\cdots. \tag{C.10}$$

The $U(1)$ charge conservation in these OPEs (C.8), (C.9) and (C.10) can be checked from Table 2 or 4.

As performed before, one has the nontrivial OPEs between the spin-1 currents and the spin-3 currents, $U_{\pm}^{(3)}(w)$ and $V_{\pm}^{(3)}(w)$ in (3.17), (3.20), (3.27) and (3.30) as follows:

$$\hat{B}_{\pm}(z)\begin{pmatrix} U_{+}^{(3)} \\ V_{-}^{(3)} \end{pmatrix}(w) = \mp\frac{1}{(z-w)^2}\,4i\,T^{(2)}(w) + \frac{1}{(z-w)}\,i(T^{(3)}-W^{(3)})(w) + \cdots, \quad \text{(C.11)}$$

$$\hat{B}_{3}(z)\begin{pmatrix} U_{+}^{(3)} \\ V_{-}^{(3)} \end{pmatrix}(w) = \frac{1}{(z-w)}\,i\begin{pmatrix} U_{+}^{(3)} \\ -V_{-}^{(3)} \end{pmatrix}(w) + \cdots, \quad \text{(C.12)}$$

$$\hat{A}_{\mp}(z)\begin{pmatrix} U_{-}^{(3)} \\ V_{+}^{(3)} \end{pmatrix}(w) = \pm\frac{1}{(z-w)^2}\,4i\,T^{(2)}(w) + \frac{1}{(z-w)}\,i(T^{(3)}+W^{(3)})(w)\cdots, \quad \text{(C.13)}$$

$$\hat{A}_{3}(z)\begin{pmatrix} U_{-}^{(3)} \\ V_{+}^{(3)} \end{pmatrix}(w) = \frac{1}{(z-w)}\,i\begin{pmatrix} -U_{-}^{(3)} \\ V_{+}^{(3)} \end{pmatrix}(w) + \cdots. \quad \text{(C.14)}$$

One can check the $U(1)$ charge conservation in these OPEs (C.11), (C.12), (C.13) and (C.14) from Table 2 or 5. There is no descendant field for the spin-2 field in the first order pole of (C.11) or (C.13).

The following nontrivial OPEs between the spin-1 currents and the spin-$\frac{7}{2}$ current $U^{(\frac{7}{2})}(w)$ (3.22) can be obtained as

$$\hat{A}_{+}(z)\,U^{(\frac{7}{2})}(w) = \frac{1}{(z-w)}\left[\frac{2}{(6+k)}\hat{A}_{+}U^{(\frac{5}{2})}\right](w) + \cdots, \quad \text{(C.15)}$$

$$\hat{A}_{-}(z)\,U^{(\frac{7}{2})}(w) = \frac{1}{(z-w)^2}\left[\frac{2i(71+13k)}{5(6+k)}T_{+}^{(\frac{5}{2})}\right](w) \quad \text{(C.16)}$$

$$+ \frac{1}{(z-w)}\left[iW_{+}^{(\frac{7}{2})} - \frac{2}{(6+k)}\left(\hat{B}_{-}V^{(\frac{5}{2})} + U^{(\frac{5}{2})}\hat{A}_{-} - \frac{3i}{5}\partial T_{+}^{(\frac{5}{2})}\right)\right](w) + \cdots,$$

$$\hat{A}_{3}(z)\,U^{(\frac{7}{2})}(w) = \frac{1}{(z-w)^2}\left[\frac{6i(11+3k)}{5(6+k)}U^{(\frac{5}{2})}\right](w) - \frac{1}{(z-w)}\,\frac{i}{2}\,U^{(\frac{7}{2})}(w) + \cdots, \quad \text{(C.17)}$$

$$\hat{B}_{+}(z)\,U^{(\frac{7}{2})}(w) = \frac{1}{(z-w)^2}\left[\frac{8i(11+3k)}{5(6+k)}T_{-}^{(\frac{5}{2})}\right](w) \quad \text{(C.18)}$$

$$+ \frac{1}{(z-w)}\left[iW_{-}^{(\frac{7}{2})} + \frac{2}{(6+k)}\left(2\hat{A}_{3}T_{-}^{(\frac{5}{2})} - U^{(\frac{5}{2})}\hat{B}_{+} - \frac{3i}{5}\partial T_{-}^{(\frac{5}{2})}\right)\right](w) + \cdots,$$

$$\hat{B}_{3}(z)\,U^{(\frac{7}{2})}(w) = \frac{1}{(z-w)^2}\left[\frac{4i(11+3k)}{5(6+k)}U^{(\frac{5}{2})}\right](w) + \frac{1}{(z-w)}\,\frac{i}{2}\,U^{(\frac{7}{2})}(w) + \cdots. \quad \text{(C.19)}$$

One can check the $U(1)$ charge conservation in these OPEs from Table 2 or 6. Nonlinear terms in (C.15), (C.16), (C.18) (linear in the higher spin currents) appear. In (C.17) and (C.19), only linear terms appear. There is no descendant field for the spin-$\frac{5}{2}$ field in the first order pole of (C.16), (C.17), (C.18) or (C.19). One can check in (C.16) or (C.18) whether the last terms (containing the composite fields) in the first order poles are (quasi)primary fields.

The nontrivial OPEs between the spin-1 currents and the spin-$\frac{7}{2}$ current $V^{(\frac{7}{2})}(w)$ (3.32) are given by

$$\begin{aligned}
\hat{A}_+(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{8i(15+2k)}{5(6+k)}T_-^{(\frac{5}{2})}\right](w) && \text{(C.20)}\\
&+ \frac{1}{(z-w)}\left[-iW_-^{(\frac{7}{2})}+\frac{2}{(6+k)}\left(2\hat{B}_3T_-^{(\frac{5}{2})}-\hat{A}_+V^{(\frac{5}{2})}+\frac{2i}{5}\partial T_-^{(\frac{5}{2})}\right)\right](w)+\cdots,\\
\hat{A}_3(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{4i(15+2k)}{5(6+k)}V^{(\frac{5}{2})}\right](w)+\frac{1}{(z-w)}\,\frac{i}{2}\,V^{(\frac{7}{2})}(w)+\cdots, && \text{(C.21)}\\
\hat{B}_+(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)}\left[\frac{2}{(6+k)}V^{(\frac{5}{2})}\hat{B}_+\right](w)+\cdots, && \text{(C.22)}\\
\hat{B}_-(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{6i(25+4k)}{5(6+k)}T_+^{(\frac{5}{2})}\right](w) && \text{(C.23)}\\
&- \frac{1}{(z-w)}\left[iW_+^{(\frac{7}{2})}+\frac{2}{(6+k)}\left(\hat{B}_-V^{(\frac{5}{2})}+U^{(\frac{5}{2})}\hat{A}_--\frac{3i}{5}\partial T_+^{(\frac{5}{2})}\right)\right](w)+\cdots,\\
\hat{B}_3(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{6i(15+2k)}{5(6+k)}V^{(\frac{5}{2})}\right](w)-\frac{1}{(z-w)}\,\frac{i}{2}\,V^{(\frac{7}{2})}(w)+\cdots. && \text{(C.24)}
\end{aligned}$$

One can check the $U(1)$ charge conservation in these OPEs from Table 2 or 6. In (C.20), (C.22) and (C.23), the nonlinear terms (still linear in higher spin currents) occur while, in (C.21) and (C.24), the linear terms occur. There is no descendant field for the spin-$\frac{5}{2}$ field in the first order pole of (C.20), (C.21), (C.23) or (C.24). One can check in (C.20) or (C.23) whether the last terms (containing the composite fields) in the first order poles are (quasi)primary fields.

Therefore, one obtains the OPEs between six spin-1 currents and the higher spin current of spins $(\frac{5}{2},3,3,\frac{7}{2})$ explicitly. Most of the higher spin currents corresponding to four $\mathcal{N}=2$ multiplets (3.1) occur in the right hand sides of the OPEs.

Appendix C.3 The OPEs between six spin 1 currents and the higher spin current of spins $(3,\frac{7}{2},\frac{7}{2},4)$

The OPEs between the spin-1 currents and the spin-3 current $W^{(3)}(w)$ (3.34) can be described as

$$\begin{aligned}
\hat{A}_\pm(z)\,W^{(3)}(w) &= \frac{1}{(z-w)}i\begin{pmatrix}U_-^{(3)}\\ V_+^{(3)}\end{pmatrix}(w)+\cdots, && \text{(C.25)}\\
\hat{A}_3(z)\,W^{(3)}(w) &= \frac{1}{(z-w)^2}\,2iT^{(2)}(w)+\cdots=-\hat{B}_3(z)\,W^{(3)}(w), && \text{(C.26)}\\
\hat{B}_\pm(z)\,W^{(3)}(w) &= -\frac{1}{(z-w)}i\begin{pmatrix}V_-^{(3)}\\ U_+^{(3)}\end{pmatrix}(w)+\cdots. && \text{(C.27)}
\end{aligned}$$

From Table 2 or 5, one can check the $U(1)$ charges in (C.25), (C.26) and (C.27).

The OPEs between the spin-1 currents and the spin-$\frac{7}{2}$ current $W_+^{(\frac{7}{2})}(w)$ (3.36) are as follows:

$$\begin{aligned}
\hat{A}_+(z)\,W_+^{(\frac{7}{2})}(w) &= -\frac{1}{(z-w)^2}\left[\frac{2i(63+13k)}{5(6+k)}U^{(\frac{5}{2})}\right](w) \qquad \text{(C.28)}\\
&+ \frac{1}{(z-w)}\left[iU^{(\frac{7}{2})}+\frac{2}{(6+k)}\left(\hat{B}_-T_-^{(\frac{5}{2})}-T_+^{(\frac{5}{2})}\hat{A}_+ +\frac{3i}{5}\partial U^{(\frac{5}{2})}\right)\right](w)+\cdots,\\
\hat{A}_-(z)\,W_+^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)}\left[\frac{2}{(6+k)}T_+^{(\frac{5}{2})}\hat{A}_-\right](w)+\cdots, \qquad \text{(C.29)}\\
\hat{A}_3(z)\,W_+^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{4i(17+2k)}{5(6+k)}T_+^{(\frac{5}{2})}\right](w)+\frac{1}{(z-w)}\frac{i}{2}W_+^{(\frac{7}{2})}(w)+\cdots, \qquad \text{(C.30)}\\
\hat{B}_+(z)\,W_+^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{2i(67+12k)}{5(6+k)}V^{(\frac{5}{2})}\right](w) \qquad \text{(C.31)}\\
&+ \frac{1}{(z-w)}\left[-iV^{(\frac{7}{2})}+\frac{2}{(6+k)}\left(T_+^{(\frac{5}{2})}\hat{B}_+ -\hat{A}_-T_-^{(\frac{5}{2})}-\frac{3i}{5}\partial V^{(\frac{5}{2})}\right)\right](w)+\cdots,\\
\hat{B}_-(z)\,W_+^{(\frac{7}{2})}(w) &= -\frac{1}{(z-w)}\left[\frac{2}{(6+k)}\hat{B}_-T_+^{(\frac{5}{2})}\right](w)+\cdots, \qquad \text{(C.32)}\\
\hat{B}_3(z)\,W_+^{(\frac{7}{2})}(w) &= -\frac{1}{(z-w)^2}\left[\frac{4i(13+3k)}{5(6+k)}T_+^{(\frac{5}{2})}\right](w)+\frac{1}{(z-w)}\frac{i}{2}W_+^{(\frac{7}{2})}(w)+\cdots. \qquad \text{(C.33)}
\end{aligned}$$

The $U(1)$ charge conservation in these OPEs can be checked by Table 2 or 6. The nonlinear terms (still linear in higher spin currents) appear in (C.28), (C.29), (C.31) and (C.32) while the linear terms appear in (C.30) and (C.33). There is no descendant field for the spin-$\frac{5}{2}$ field in the first order pole of (C.28), (C.30), (C.31) or (C.33). One can check in (C.28) or (C.31) whether the last terms (containing the composite fields) in the first order poles are (quasi)primary fields.

The OPEs between the spin-1 currents and the spin-$\frac{7}{2}$ current $W_-^{(\frac{7}{2})}(w)$ (3.38) are expressed as

$$\begin{aligned}
\hat{A}_+(z)\,W_-^{(\frac{7}{2})}(w) &= -\frac{1}{(z-w)}\left[\frac{2}{(6+k)}\hat{A}_-T_-^{(\frac{5}{2})}\right](w)+\cdots, \qquad \text{(C.34)}\\
\hat{A}_-(z)\,W_-^{(\frac{7}{2})}(w) &= -\frac{1}{(z-w)^2}\left[\frac{2i(63+13k)}{5(6+k)}V^{(\frac{5}{2})}\right](w) \qquad \text{(C.35)}\\
&+ \frac{1}{(z-w)}\left[-iV^{(\frac{7}{2})}+\frac{4}{(6+k)}\left(\hat{B}_3V^{(\frac{5}{2})}-\hat{A}_3V^{(\frac{5}{2})}+\frac{i}{5}\partial V^{(\frac{5}{2})}\right)\right](w)+\cdots,\\
\hat{A}_3(z)\,W_-^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{4i(17+2k)}{5(6+k)}T_-^{(\frac{5}{2})}\right](w)-\frac{1}{(z-w)}\frac{i}{2}W_-^{(\frac{7}{2})}(w)+\cdots, \qquad \text{(C.36)}\\
\hat{B}_+(z)\,W_-^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)}\left[\frac{2}{(6+k)}T_-^{(\frac{5}{2})}\hat{B}_+\right](w)+\cdots, \qquad \text{(C.37)}
\end{aligned}$$

$$\hat{B}_-(z)\,W_-^{(\frac{7}{2})}(w) = \frac{1}{(z-w)^2}\left[\frac{2i(67+12k)}{5(6+k)}U^{(\frac{5}{2})}\right](w) \tag{C.38}$$
$$+ \frac{1}{(z-w)}\left[iU^{(\frac{7}{2})}+\frac{4}{(6+k)}\left(U^{(\frac{5}{2})}\hat{B}_3-U^{(\frac{5}{2})}\hat{A}_3+\frac{4i}{5}\partial U^{(\frac{5}{2})}\right)\right](w)+\cdots,$$
$$\hat{B}_3(z)\,W_-^{(\frac{7}{2})}(w) = -\frac{1}{(z-w)^2}\left[\frac{4i(13+3k)}{5(6+k)}T_-^{(\frac{5}{2})}\right](w)-\frac{1}{(z-w)}\frac{i}{2}W_-^{(\frac{7}{2})}(w)+\cdots. \tag{C.39}$$

The $U(1)$ charge conservation in these OPEs can be checked by Table 2 or 6. The linear terms appear in (C.36) and (C.39) and the nonlinear terms (linear in higher spin currents) appear in (C.34), (C.35), (C.37) and (C.38). There is no descendant field for the spin-$\frac{5}{2}$ field in the first order pole of (C.35), (C.36), (C.38) or (C.39). One can check in (C.35) or (C.39) whether the last terms (containing the composite fields) in the first order poles are (quasi)primary fields.

Finally the six nontrivial OPEs between the spin-1 currents and the spin-4 current $W^{(4)}(w)$ (3.40) can be described as

$$\hat{A}_+(z)\,W^{(4)}(w) = \frac{1}{(z-w)^2}\left[\frac{2i(43+6k)}{3(6+k)}U_-^{(3)}+\frac{16(-972+k(-125+27k))}{3(324+179k)(6+k)}T^{(2)}\hat{A}_+\right](w)$$
$$+ \frac{1}{(z-w)}\frac{1}{(6+k)}\left[\frac{8}{3}\hat{A}_+\partial T^{(2)}+4\hat{B}_3U_-^{(3)}-\frac{16}{3}T^{(2)}\partial\hat{A}_+\right.$$
$$- \left.4T^{(3)}\hat{A}_+-\frac{10i}{3}\partial U_-^{(3)}\right](w)+\cdots, \tag{C.40}$$
$$\hat{A}_-(z)\,W^{(4)}(w) = \frac{1}{(z-w)^2}\left[-\frac{2i(47+9k)}{3(6+k)}V_+^{(3)}+\frac{8(-1620-71k+54k^2)}{3(324+179k)(6+k)}\hat{A}_-T^{(2)}\right](w)$$
$$+ \frac{1}{(z-w)}\frac{1}{(6+k)}\left[2\hat{A}_-T^{(3)}-2\hat{A}_-W^{(3)}-\frac{8}{3}\hat{A}_-\partial T^{(2)}+\frac{16}{3}T^{(2)}\partial\hat{A}_-\right.$$
$$- \left.4V_+^{(3)}\hat{A}_3+4V_+^{(3)}\hat{B}_3-\frac{10i}{3}\partial V_+^{(3)}\right](w)+\cdots, \tag{C.41}$$
$$\hat{A}_3(z)\,W^{(4)}(w) = -\frac{1}{(z-w)^3}\left[\frac{16i(1+k)}{3(6+k)}T^{(2)}\right](w)$$
$$+ \frac{1}{(z-w)^2}\left[\frac{i(16+k)}{(6+k)}T^{(3)}+\frac{i(16+3k)}{(6+k)}W^{(3)}+\frac{8}{(6+k)}\hat{B}_3T^{(2)}\right.$$
$$+ \left.\frac{8(-756-143k+18k^2)}{(324+179k)(6+k)}\hat{A}_3T^{(2)}+\frac{4i(1+k)}{3(6+k)}\partial T^{(2)}\right](w)+\cdots, \tag{C.42}$$
$$\hat{B}_+(z)\,W^{(4)}(w) = \frac{1}{(z-w)^2}\left[-\frac{2i(31+9k)}{3(6+k)}V_-^{(3)}+\frac{16(-324+233k+27k^2)}{3(324+179k)(6+k)}T^{(2)}\hat{B}_+\right](w)$$
$$+ \frac{1}{(z-w)}\frac{1}{(6+k)}\left[-4\hat{A}_3V_-^{(3)}-\frac{8}{3}\hat{B}_+\partial T^{(2)}+\frac{16}{3}T^{(2)}\partial\hat{B}_+\right.$$
$$+ \left.4T^{(3)}\hat{B}_++\frac{10i}{3}\partial V_-^{(3)}\right](w)+\cdots, \tag{C.43}$$

$$\begin{aligned}
\hat{B}_-(z)\,W^{(4)}(w) &= \frac{1}{(z-w)^2}\left[\frac{2i(47+9k)}{3(6+k)}U_+^{(3)}+\frac{8(-972+287k+54k^2)}{3(324+179k)(6+k)}\hat{B}_-T^{(2)}\right](w)\\
&+ \frac{1}{(z-w)}\frac{1}{(6+k)}\left[-2\hat{B}_-T^{(3)}-2\hat{B}_-W^{(3)}+\frac{8}{3}\hat{B}_-\partial T^{(2)}+\frac{16}{3}T^{(2)}\partial\hat{B}_-\right.\\
&- \left. 4U_+^{(3)}\hat{A}_3+4U_+^{(3)}\hat{B}_3+\frac{10i}{3}\partial U_+^{(3)}\right](w)+\cdots, \qquad \text{(C.44)}\\
\hat{B}_3(z)\,W^{(4)}(w) &= \frac{1}{(z-w)^3}\left[\frac{80i}{3(6+k)}iT^{(2)}\right](w)\\
&+ \frac{1}{(z-w)^2}\left[-\frac{i(8+3k)}{(6+k)}T^{(3)}+\frac{i(16+3k)}{(6+k)}W^{(3)}-\frac{8}{(6+k)}\hat{A}_3T^{(2)}\right.\\
&+ \left.\frac{8(-108+215k+18k^2)}{(324+179k)(6+k)}\hat{B}_3T^{(2)}-\frac{20i}{3(6+k)}\partial T^{(2)}\right](w)+\cdots. \qquad \text{(C.45)}
\end{aligned}$$

The $U(1)$ charge conservation in these OPEs (C.40), (C.41), (C.42), (C.43), (C.44) and (C.45) can be checked by Table 2 or 5. There is no descendant field for the spin-3 field in the first order pole of (C.40), (C.41), (C.42), (C.43), (C.44) or (C.45). In (C.42) or (C.45), each term (except the last term) in the second order pole is a primary field.

Therefore, one obtains the OPEs between six spin-1 currents and the higher spin current of spins $(3,\frac{7}{2},\frac{7}{2},4)$ explicitly. Most of the higher spin currents corresponding to four $\mathcal{N}=2$ multiplets (3.1) occur in the right hand sides of the OPEs. The nonlinear terms in this Appendix are quadratic terms. Note that, for unitary case [20], there are also cubic terms in their corresponding OPEs.

One obtains the following results between the spin-1 currents and the 16 higher spin currents by focusing on the linear terms of higher spin currents in the right hand side of the OPEs as follows:

$$\hat{A}_+ \times \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix} \to \begin{pmatrix} 0, & U^{(\frac{5}{2})}, & 0, & U_-^{(3)} \\ 0, & 0, & 0, & 0 \\ T_-^{(\frac{5}{2})}, & T^{(2)},T^{(3)},W^{(3)}, & 0, & T_-^{(\frac{5}{2})},W_-^{(\frac{7}{2})} \\ U_-^{(3)}, & U^{(\frac{5}{2})},U^{(\frac{7}{2})}, & 0, & U_-^{(3)} \end{pmatrix},$$

$$\hat{A}_- \times \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix} \to \begin{pmatrix} 0, & 0, & V^{(\frac{5}{2})}, & V_+^{(3)} \\ T_+^{(\frac{5}{2})}, & 0, & T^{(2)},T^{(3)},W^{(3)}, & T_+^{(\frac{5}{2})},W_+^{(\frac{7}{2})} \\ 0, & 0, & 0, & 0 \\ V_+^{(3)}, & 0, & V^{(\frac{5}{2})},V^{(\frac{7}{2})}, & V_+^{(3)} \end{pmatrix},$$

$$\hat{A}_3 \times \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix} \to \begin{pmatrix} 0, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(2)} \\ U^{(\frac{5}{2})}, & 0, & U_-^{(3)}, & U^{(\frac{5}{2})},U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})} & V_+^{(3)}, & 0, & V^{(\frac{5}{2})},V^{(\frac{7}{2})} \\ T^{(2)} & T_+^{(\frac{5}{2})},W_+^{(\frac{7}{2})}, & T_-^{(\frac{5}{2})},W_-^{(\frac{7}{2})}, & T^{(2)},T^{(3)},W^{(3)} \end{pmatrix},$$

$$\hat{B}_+ \times \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix} \to \begin{pmatrix} 0, & V^{(\frac{5}{2})}, & 0, & V_-^{(3)} \\ T_-^{(\frac{5}{2})}, & T^{(2)}, T^{(3)}, W^{(3)}, & 0, & T_-^{(\frac{5}{2})}, W_-^{(\frac{7}{2})} \\ 0 & 0, & 0, & 0 \\ V_-^{(3)}, & V^{(\frac{5}{2})}, V^{(\frac{7}{2})}, & 0, & V_-^{(3)} \end{pmatrix},$$

$$\hat{B}_- \times \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix} \to \begin{pmatrix} 0, & 0, & U^{(\frac{5}{2})}, & U_+^{(3)} \\ 0, & 0, & 0, & 0 \\ T_+^{(\frac{5}{2})}, & 0, & T^{(2)}, T^{(3)}, W^{(3)}, & T_+^{(\frac{5}{2})}, W_+^{(\frac{7}{2})} \\ U_+^{(3)}, & 0, & U^{(\frac{5}{2})}, U^{(\frac{7}{2})}, & U_+^{(3)} \end{pmatrix},$$

$$\hat{B}_3 \times \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix} \to \begin{pmatrix} 0 & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(2)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & 0, & U^{(\frac{5}{2})}, U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & 0, & V_-^{(3)}, & V^{(\frac{5}{2})}, V^{(\frac{7}{2})} \\ T^{(2)} & T_+^{(\frac{5}{2})}, W_+^{(\frac{7}{2})}, & T_-^{(\frac{5}{2})}, W_-^{(\frac{7}{2})}, & T^{(2)}, T^{(3)}, W^{(3)} \end{pmatrix}.$$

Note that the zeros of the right hand sides for the OPEs between $\hat{A}_\pm$ and $\hat{B}_\pm$ acting on the 16 higher spin currents are located at the row and column containing the corresponding spin-1 current in (2.44) of [20]. The zeros from the OPEs between these four spin-1 currents and spin-2 current $T^{(2)}(w)$ also occur.

Appendix D The OPEs between the four spin-$\frac{3}{2}$ currents and the higher spin currents

In this Appendix, we present the OPEs between the four spin-$\frac{3}{2}$ currents in section 2 and the 16 higher spin currents in section 3.

Appendix D.1 The OPEs between four spin $\frac{3}{2}$ currents and the higher spin current of spins $(2, \frac{5}{2}, \frac{5}{2}, 3)$

The OPEs between the spin-$\frac{3}{2}$ currents (2.33), (2.34), (2.35) and (2.36) and the spin-2 current $T^{(2)}(w)$ are derived as follows:

$$\begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{22} \end{pmatrix}(z)\, T^{(2)}(w) = \frac{1}{(z-w)} \begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(w) + \cdots, \tag{D.1}$$

$$\begin{pmatrix} \hat{G}_{12} \\ \hat{G}_{21} \end{pmatrix}(z)\, T^{(2)}(w) = \frac{1}{(z-w)} \begin{pmatrix} T_-^{(\frac{5}{2})} \\ T_+^{(\frac{5}{2})} \end{pmatrix}(w) + \cdots. \tag{D.2}$$

One can check the $U(1)$ charge conservation in these OPEs (D.1) and (D.2) from Table 2 or 4.

The following nontrivial OPEs between the spin-$\frac{3}{2}$ currents and the spin-$\frac{5}{2}$ currents $T_{\pm}^{(\frac{5}{2})}(w)$ in (3.8) and (3.10) are as follows:

$$\begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{22} \end{pmatrix}(z)\, T_{+}^{(\frac{5}{2})}(w) = -\frac{1}{(z-w)} \begin{pmatrix} U_{+}^{(3)} \\ V_{+}^{(3)} \end{pmatrix}(w) + \cdots, \tag{D.3}$$

$$\begin{pmatrix} \hat{G}_{12} \\ \hat{G}_{21} \end{pmatrix}(z)\, T_{\pm}^{(\frac{5}{2})}(w) = \frac{1}{(z-w)} 4T^{(2)}(w) + \frac{1}{(z-w)^2}(\partial T^{(2)} \mp T^{(3)})(w) + \cdots, \tag{D.4}$$

$$\begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{22} \end{pmatrix}(z)\, T_{-}^{(\frac{5}{2})}(w) = -\frac{1}{(z-w)} \begin{pmatrix} U_{-}^{(3)} \\ V_{-}^{(3)} \end{pmatrix}(w) + \cdots. \tag{D.5}$$

The $U(1)$ charge conservation in these OPEs can be checked from Tables 2 or 5. No nonlinear terms in (D.3), (D.4) and (D.5) occur.

One calculates the following OPEs between the spin-$\frac{3}{2}$ currents and the spin-3 current $T^{(3)}(w)$ (3.12) as follows:

$$\begin{aligned} \begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{22} \end{pmatrix}(z)\, T^{(3)}(w) &= \frac{1}{(z-w)^2} \frac{(-4+k)}{(6+k)} \begin{pmatrix} U^{(\frac{5}{2})} \\ -V^{(\frac{5}{2})} \end{pmatrix} \\ &+ \frac{1}{(z-w)} \left[\frac{(-4+k)}{5(6+k)} \partial \begin{pmatrix} U^{(\frac{5}{2})} \\ -V^{(\frac{5}{2})} \end{pmatrix} + \begin{pmatrix} U^{(\frac{7}{2})} \\ V^{(\frac{7}{2})} \end{pmatrix} \right. \\ &+ \left. \frac{2}{(6+k)} \begin{pmatrix} 2iU^{(\frac{5}{2})} \hat{A}_3 - i\hat{B}_- T_{-}^{(\frac{5}{2})} + \frac{3}{5}\partial U^{(\frac{5}{2})} \\ 2i\hat{B}_3 V^{(\frac{5}{2})} - i\hat{A}_- T_{-}^{(\frac{5}{2})} - \frac{2}{5}\partial V^{(\frac{5}{2})} \end{pmatrix} \right](w) + \cdots, \end{aligned} \tag{D.6}$$

$$\begin{aligned} \begin{pmatrix} \hat{G}_{12} \\ \hat{G}_{21} \end{pmatrix}(z)\, T^{(3)}(w) &= \frac{1}{(z-w)^2} \frac{(26+5k)}{(6+k)} \begin{pmatrix} T_{-}^{(\frac{5}{2})} \\ -T_{+}^{(\frac{5}{2})} \end{pmatrix} \\ &+ \frac{1}{(z-w)} \left[\frac{(26+5k)}{5(6+k)} \partial \begin{pmatrix} T_{-}^{(\frac{5}{2})} \\ -T_{+}^{(\frac{5}{2})} \end{pmatrix} \right. \\ &+ \left. \frac{2}{(6+k)} \begin{pmatrix} i\hat{A}_+ V^{(\frac{5}{2})} + iU^{(\frac{5}{2})} \hat{B}_+ - \frac{3}{5}\partial T_{-}^{(\frac{5}{2})} \\ i\hat{B}_- V^{(\frac{5}{2})} + iU^{(\frac{5}{2})} \hat{A}_- + \frac{3}{5}\partial T_{+}^{(\frac{5}{2})} \end{pmatrix} \right](w) + \cdots. \end{aligned} \tag{D.7}$$

The $U(1)$ charge conservation in these OPEs can be checked from Tables 2 or 6. The quadratic terms (still linear in higher spin currents) in (D.6) and (D.7) occur. The factor $(k-4)$ appears in (D.6).

Therefore, one obtains the OPEs between six spin-1 currents and the higher spin current of spins $(2, \frac{5}{2}, \frac{5}{2}, 3)$ explicitly. The higher spin currents corresponding to three $\mathcal{N} = 2$ multiplets (3.1) occur in the right hand sides of the OPEs.

Appendix D.2 The OPEs between four spin $\frac{3}{2}$ currents and two higher spin currents of spins $(\frac{5}{2}, 3, 3, \frac{7}{2})$

The nontrivial OPEs between the spin-$\frac{3}{2}$ currents and the spin-$\frac{5}{2}$ currents, $U^{(\frac{5}{2})}(z)$ (3.14) and $V^{(\frac{5}{2})}(z)$ (3.24), can be obtained as follows:

$$\begin{pmatrix} \hat{G}_{22} \\ \hat{G}_{11} \end{pmatrix}(z) \begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(z) = \frac{1}{(z-w)^2} 4T^{(2)}(w) + \frac{1}{(z-w)} (\partial T^{(2)} \pm W^{(3)})(w) + \cdots, \tag{D.8}$$

$$\hat{G}_{12}(z) \begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(z) = \frac{1}{(z-w)} \begin{pmatrix} U^{(3)}_{-} \\ V^{(3)}_{-} \end{pmatrix}(w) + \cdots, \tag{D.9}$$

$$\hat{G}_{21}(z) \begin{pmatrix} U^{(\frac{5}{2})} \\ V^{(\frac{5}{2})} \end{pmatrix}(z) = \frac{1}{(z-w)} \begin{pmatrix} U^{(3)}_{+} \\ V^{(3)}_{+} \end{pmatrix}(w) + \cdots. \tag{D.10}$$

The $U(1)$ charge conservation in these OPEs (D.8), (D.9) and (D.10) can be checked from Table 2 or 5.

The OPEs between the spin-$\frac{3}{2}$ currents and the spin-3 currents $U^{(3)}_{\pm}(w)$ in (3.17) and (3.20) are as follows:

$$\begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{21} \end{pmatrix}(z)\, U^{(3)}_{+}(w) = -\frac{1}{(z-w)} \frac{2i}{(6+k)} \hat{B}_{-} \begin{pmatrix} U^{(\frac{5}{2})} \\ T^{(\frac{5}{2})}_{+} \end{pmatrix}(w) + \cdots, \tag{D.11}$$

$$\begin{aligned} \begin{pmatrix} \hat{G}_{22} \\ \hat{G}_{12} \end{pmatrix}(z)\, U^{(3)}_{+}(w) &= \frac{1}{(z-w)^2} \frac{(30+4k)}{(6+k)} \begin{pmatrix} -T^{(\frac{5}{2})}_{+} \\ U^{(\frac{5}{2})} \end{pmatrix} \\ &+ \frac{1}{(z-w)} \left[\frac{(30+4k)}{5(6+k)} \partial \begin{pmatrix} -T^{(\frac{5}{2})}_{+} \\ U^{(\frac{5}{2})} \end{pmatrix} - \begin{pmatrix} W^{(\frac{7}{2})}_{+} \\ U^{(\frac{7}{2})} \end{pmatrix} \right. \\ &+ \left. \frac{2}{(6+k)} \begin{pmatrix} iU^{(\frac{5}{2})} \hat{A}_{-} - 2iT^{(\frac{5}{2})}_{+} \hat{B}_3 + \frac{8}{5} \partial T^{(\frac{5}{2})}_{+} \\ i\hat{B}_{-} T^{(\frac{5}{2})}_{-} - iT^{(\frac{5}{2})}_{+} \hat{A}_{+} - 2iU^{(\frac{5}{2})} (\hat{A}_3 - \hat{B}_3) - \frac{11}{5} \partial U^{(\frac{5}{2})} \end{pmatrix} \right](w) \\ &+ \cdots, \end{aligned} \tag{D.12}$$

$$\begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{12} \end{pmatrix}(z)\, U^{(3)}_{-}(w) = \frac{1}{(z-w)} \frac{2i}{(6+k)} \begin{pmatrix} U^{(\frac{5}{2})} \hat{A}_{+} \\ \hat{A}_{+} T^{(\frac{5}{2})}_{-} \end{pmatrix}(w) + \cdots, \tag{D.13}$$

$$\begin{aligned} \begin{pmatrix} \hat{G}_{22} \\ \hat{G}_{21} \end{pmatrix}(z)\, U^{(3)}_{-}(w) &= \frac{1}{(z-w)^2} \frac{(22+6k)}{(6+k)} \begin{pmatrix} -T^{(\frac{5}{2})}_{-} \\ U^{(\frac{5}{2})} \end{pmatrix} \\ &+ \frac{1}{(z-w)} \left[\frac{(22+6k)}{5(6+k)} \partial \begin{pmatrix} -T^{(\frac{5}{2})}_{-} \\ U^{(\frac{5}{2})} \end{pmatrix} + \begin{pmatrix} -W^{(\frac{7}{2})}_{-} \\ U^{(\frac{7}{2})} \end{pmatrix} \right. \\ &+ \left. \frac{2}{(6+k)} \begin{pmatrix} 2i\hat{A}_3 T^{(\frac{5}{2})}_{-} - iU^{(\frac{5}{2})} \hat{B}_{+} + \frac{3}{5} \partial T^{(\frac{5}{2})}_{-} \\ 0 \end{pmatrix} \right](w) + \cdots. \end{aligned} \tag{D.14}$$

The $U(1)$ charge conservation in these OPEs can be checked from Table 2 or 6. The nonlinear terms (linear in higher spin currents) occur in (D.11), (D.12), (D.13) and (D.14).

The OPEs between the spin-$\frac{3}{2}$ currents and the spin-3 currents $V^{(3)}_{\pm}(w)$ in (3.27) and (3.30) are expressed as

$$\begin{aligned}
\begin{pmatrix} \hat{G}_{22} \\ \hat{G}_{21} \end{pmatrix}(z)\, V^{(3)}_{+}(w) &= -\frac{1}{(z-w)}\frac{2i}{(6+k)} \begin{pmatrix} \hat{A}_{-}V^{(\frac{5}{2})} \\ T^{(\frac{5}{2})}_{+}\hat{A}_{-} \end{pmatrix}(w) + \cdots, && \text{(D.15)} \\
\begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{12} \end{pmatrix}(z)\, V^{(3)}_{+}(w) &= \frac{1}{(z-w)^2}\frac{(22+6k)}{(6+k)} \begin{pmatrix} -T^{(\frac{5}{2})}_{+} \\ V^{(\frac{5}{2})} \end{pmatrix} && \text{(D.16)} \\
&+ \frac{1}{(z-w)}\left[\frac{(22+6k)}{5(6+k)}\partial \begin{pmatrix} -T^{(\frac{5}{2})}_{+} \\ V^{(\frac{5}{2})} \end{pmatrix} + \begin{pmatrix} W^{(\frac{7}{2})}_{+} \\ -V^{(\frac{7}{2})} \end{pmatrix}\right. \\
&+ \left.\frac{2}{(6+k)} \begin{pmatrix} i\hat{B}_{-}V^{(\frac{5}{2})} - 2iT^{(\frac{5}{2})}_{+}\hat{A}_{3} + \frac{3}{5}\partial T^{(\frac{5}{2})}_{+} \\ i\hat{A}_{-}T^{(\frac{5}{2})}_{-} + iT^{(\frac{5}{2})}_{+}\hat{B}_{+} + 2i(\hat{A}_{3}-\hat{B}_{3})V^{(\frac{5}{2})} - \frac{1}{5}\partial V^{(\frac{5}{2})} \end{pmatrix}\right](w) \\
&+ \cdots, \\
\begin{pmatrix} \hat{G}_{22} \\ \hat{G}_{12} \end{pmatrix}(z)\, V^{(3)}_{-}(w) &= \frac{1}{(z-w)}\frac{2i}{(6+k)} \begin{pmatrix} V^{(\frac{5}{2})} \\ T^{(\frac{5}{2})}_{-} \end{pmatrix}\hat{B}_{+}(w) + \cdots, && \text{(D.17)} \\
\begin{pmatrix} \hat{G}_{11} \\ \hat{G}_{21} \end{pmatrix}(z)\, V^{(3)}_{-}(w) &= \frac{1}{(z-w)^2}\frac{(30+4k)}{(6+k)} \begin{pmatrix} -T^{(\frac{5}{2})}_{-} \\ V^{(\frac{5}{2})} \end{pmatrix} && \text{(D.18)} \\
&+ \frac{1}{(z-w)}\left[\frac{(30+4k)}{5(6+k)}\partial \begin{pmatrix} -T^{(\frac{5}{2})}_{-} \\ V^{(\frac{5}{2})} \end{pmatrix} + \begin{pmatrix} W^{(\frac{7}{2})}_{-} \\ V^{(\frac{7}{2})} \end{pmatrix}\right. \\
&+ \left.\frac{2}{(6+k)} \begin{pmatrix} -i\hat{A}_{+}V^{(\frac{5}{2})} + 2i\hat{B}_{3}T^{(\frac{5}{2})}_{-} - \frac{2}{5}\partial T^{(\frac{5}{2})}_{-} \\ 0 \end{pmatrix}\right](w) + \cdots.
\end{aligned}$$

The $U(1)$ charge conservation in these OPEs can be checked from Table 2 or 6. In this case also, the nonlinear terms (linear in higher spin currents) appear in (D.15), (D.16), (D.17) and (D.18).

The OPEs between the spin-$\frac{3}{2}$ currents and the spin-$\frac{7}{2}$ currents $U^{(\frac{7}{2})}(w)$ (3.22) are given by

$$\begin{aligned}
\hat{G}_{11}(z)\, U^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)}\left[-\frac{4i}{(6+k)}\hat{B}_{-}U^{(3)}_{-} + \frac{2}{(6+k)}\hat{G_{11}}U^{(\frac{5}{2})} - \frac{2i}{(6+k)}U^{(3)}_{+}\hat{A}_{+}\right](w) \\
&+ \cdots, && \text{(D.19)} \\
\hat{G}_{22}(z)\, U^{(\frac{7}{2})}(w) &= -\frac{1}{(z-w)^3}\left[\frac{48(-3+k)}{5(6+k)}T^{(2)}\right](w) && \text{(D.20)} \\
&+ \frac{1}{(z-w)^2}\left[\frac{6(5+k)}{(6+k)}T^{(3)} - \frac{6(-3+k)}{5(6+k)}W^{(3)} + \frac{16i}{(6+k)}\hat{B}_{3}T^{(2)}\right](w) \\
&+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{22}\, U^{(\frac{7}{2})}\}_{-2} + W^{(4)} - \frac{4i}{3(6+k)}(\hat{A}_{3}+4\hat{B}_{3})\partial T^{(2)}\right.
\end{aligned}$$

$$
\begin{aligned}
&+ \frac{4i}{(6+k)}\hat{B}_3(T^{(3)}+W^{(3)}) - \frac{4i}{(6+k)}\hat{A}_3T^{(3)} - \frac{2}{(6+k)}\hat{G}_{21}T_-^{(\frac{5}{2})} \\
&+ \frac{8i}{3(6+k)}T^{(2)}\partial(\hat{A}_3+4\hat{B}_3) - \frac{144(-4+k)}{(324+179k)}\hat{T}T^{(2)} + \frac{2i}{(6+k)}U_+^{(3)}\hat{B}_+ \\
&+ \frac{2}{(6+k)}U^{(\frac{5}{2})}\hat{G}_{22} + \frac{1}{3(6+k)}\partial(W^{(3)}-3T^{(3)}) \\
&+ \frac{6(-1188-107k+36k^2)}{5(324+179k)(6+k)}\partial^2T^{(2)}\Big](w)+\cdots,
\end{aligned}
$$

$$
\begin{aligned}
\hat{G}_{12}(z)\,U^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{2(89+12k)}{5(6+k)}U_-^{(3)} + \frac{8i}{(6+k)}T^{(2)}\hat{A}_+\right](w) + \\
&+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{12}\,U^{(\frac{7}{2})}\}_{-2} - \frac{4i}{(6+k)}(\hat{A}_3-\hat{B}_3)U_-^{(3)}\right. \\
&- \frac{10i}{3(6+k)}\hat{A}_+\partial T^{(2)} + \frac{2}{(6+k)}\hat{G}_{11}T_-^{(\frac{5}{2})} + \frac{2i}{(6+k)}W^{(3)}\hat{A}_+ \\
&+ \left.\frac{20i}{3(6+k)}T^{(2)}\partial\hat{A}_+ - \frac{1}{3(6+k)}\partial U_-^{(3)}\right](w)+\cdots,
\end{aligned}
\tag{D.21}
$$

$$
\begin{aligned}
\hat{G}_{21}(z)\,U^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[-\frac{12(11+3k)}{5(6+k)}U_+^{(3)} + \frac{16i}{(6+k)}\hat{B}_-T^{(2)}\right](w) \\
&+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{21}\,U^{(\frac{7}{2})}\}_{-2} + \frac{2i}{(6+k)}\hat{B}_-(T^{(3)}+W^{(3)})\right. \\
&- \left.\frac{8i}{3(6+k)}\hat{B}_-\partial T^{(2)} + \frac{16i}{3(6+k)}T^{(2)}\partial\hat{B}_-\right](w)+\cdots.
\end{aligned}
\tag{D.22}
$$

The $U(1)$ charge conservation in these OPEs (D.19), (D.20), (D.21) and (D.22) can be checked from Table 2, 7 or 8. Each term in the second order pole of (D.20), (D.21), or (D.22) is a primary field. There is no descendant field for the spin-2 field $T^{(2)}(w)$ in the second order pole of (D.20).

The OPEs between the spin-$\frac{3}{2}$ currents and the spin-$\frac{7}{2}$ current $V^{(\frac{7}{2})}(w)$ (3.32) are as follows:

$$
\begin{aligned}
\hat{G}_{11}(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^3}\left[\frac{48(-5+k)}{5(6+k)}T^{(2)}\right](w) \\
&+ \frac{1}{(z-w)^2}\left[\frac{6(5+k)}{(6+k)}T^{(3)} - \frac{6(-5+k)}{5(6+k)}W^{(3)} + \frac{16i}{(6+k)}\hat{A}_3T^{(2)}\right](w) \\
&+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{11}\,V^{(\frac{7}{2})}\}_{-2} - W^{(4)} + \frac{4i}{(6+k)}\hat{A}_3(T^{(3)}-W^{(3)})\right. \\
&- \frac{4i}{3(6+k)}(4\hat{A}_3+\hat{B}_3)\partial T^{(2)} - \frac{4i}{(6+k)}\hat{B}_3T^{(3)} - \frac{2}{(6+k)}\hat{G}_{11}V^{(\frac{5}{2})} \\
&- \left.\frac{2}{(6+k)}\hat{G}_{21}T_-^{(\frac{5}{2})} + \frac{8i}{3(6+k)}T^{(2)}\partial(4\hat{A}_3+\hat{B}_3) + \frac{144(-4+k)}{(324+179k)}\hat{T}T^{(2)}\right.
\end{aligned}
\tag{D.23}
$$

$$
\begin{aligned}
&+ \frac{2i}{(6+k)}V_+^{(3)}\hat{A}_+ - \frac{1}{3(6+k)}\partial(7W^{(3)}+3T^{(3)}) \\
&- \frac{8k(412+27k)}{5(324+179k)(6+k)}\partial^2 T^{(2)}\Big](w)+\cdots, \\
\hat{G}_{22}(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)}\Big[-\frac{4i}{(6+k)}\hat{A}_-V_-^{(3)} + \frac{2}{(6+k)}\hat{G_{22}}V^{(\frac{5}{2})} - \frac{2i}{(6+k)}V_+^{(3)}\hat{B}_+\Big](w) \\
&+ \cdots, \qquad \text{(D.24)} \\
\hat{G}_{12}(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\Big[\frac{2(65+18k)}{5(6+k)}V_-^{(3)} + \frac{8i}{(6+k)}T^{(2)}\hat{B}_+\Big](w) \qquad \text{(D.25)} \\
&+ \frac{1}{(z-w)}\Big[\frac{1}{6}\partial\{\hat{G}_{12}\,V^{(\frac{7}{2})}\}_{-2} + \frac{4i}{(6+k)}(\hat{A}_3-\hat{B}_3)V_-^{(3)} \\
&- \frac{10i}{3(6+k)}\hat{B}_+\partial T^{(2)} + \frac{2}{(6+k)}\hat{G}_{22}T_-^{(\frac{5}{2})} - \frac{2i}{(6+k)}W^{(3)}\hat{B}_+ \\
&+ \frac{20i}{3(6+k)}T^{(2)}\partial\hat{B}_+ - \frac{1}{3(6+k)}\partial V_-^{(3)}\Big](w)+\cdots, \\
\hat{G}_{21}(z)\,V^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\Big[-\frac{12(15+2k)}{5(6+k)}V_+^{(3)} + \frac{16i}{(6+k)}\hat{A}_-T^{(2)}\Big](w) \qquad \text{(D.26)} \\
&+ \frac{1}{(z-w)}\Big[\frac{1}{6}\partial\{\hat{G}_{21}\,V^{(\frac{7}{2})}\}_{-2} + \frac{2i}{(6+k)}\hat{A}_-(T^{(3)}-W^{(3)}) \\
&- \frac{8i}{3(6+k)}\hat{A}_-\partial T^{(2)} + \frac{16i}{3(6+k)}T^{(2)}\partial\hat{A}_-\Big](w)+\cdots.
\end{aligned}
$$

The $U(1)$ charge conservation in these OPEs (D.23), (D.24), (D.25) and (D.26) can be checked from Table 2, 7 or 8. Each term in the second order pole of (D.23), (D.25), or (D.26) is a primary field. There is no descendant field for the spin-2 field $T^{(2)}(w)$ in the second order pole of (D.23).

Therefore, one obtains the OPEs between six spin-1 currents and the higher spin current of spins $(\frac{5}{2},3,3,\frac{7}{2})$ explicitly. The higher spin currents corresponding to four $\mathcal{N}=2$ multiplets (3.1) occur in the right hand sides of the OPEs.

Appendix D.3 The OPEs between four spin $\frac{3}{2}$ currents and the higher spin current of spins $(3,\frac{7}{2},\frac{7}{2},4)$

The following four OPEs between the spin-$\frac{3}{2}$ currents and the spin-3 current $W^{(3)}(w)$ (3.34) can be obtained

$$
\begin{aligned}
\begin{pmatrix}\hat{G}_{11}\\ \hat{G}_{22}\end{pmatrix}(z)\,W^{(3)}(w) &= \frac{1}{(z-w)^2}\frac{(26+5k)}{(6+k)}\begin{pmatrix}U^{(\frac{5}{2})}\\ -V^{(\frac{5}{2})}\end{pmatrix}(w) \\
&+ \frac{1}{(z-w)}\frac{1}{(6+k)}\Big[(4+k)\partial\begin{pmatrix}U^{(\frac{5}{2})}\\ -V^{(\frac{5}{2})}\end{pmatrix}
\end{aligned}
$$

$$
\begin{aligned}
&+ \quad 2i \begin{pmatrix} \hat{B}_- T_-^{(\frac{5}{2})} - T_+^{(\frac{5}{2})} \hat{A}_+ \\ -\hat{A}_- T_-^{(\frac{5}{2})} + T_+^{(\frac{5}{2})} \hat{B}_+ \end{pmatrix} \Big] (w) + \cdots, \qquad \text{(D.27)} \\
\begin{pmatrix} \hat{G}_{12} \\ \hat{G}_{21} \end{pmatrix} (z)\, W^{(3)}(w) &= \frac{1}{(z-w)^2} \frac{(-4+k)}{(6+k)} \begin{pmatrix} T_-^{(\frac{5}{2})} \\ -T_+^{(\frac{5}{2})} \end{pmatrix} (w) \qquad \text{(D.28)} \\
&+ \quad \frac{1}{(z-w)} \left[\frac{(-4+k)}{5(6+k)} \partial \begin{pmatrix} T_-^{(\frac{5}{2})} \\ -T_+^{(\frac{5}{2})} \end{pmatrix} + \begin{pmatrix} W_-^{(\frac{7}{2})} \\ W_+^{(\frac{7}{2})} \end{pmatrix} \right] (w) + \cdots .
\end{aligned}
$$

The $U(1)$ charge conservation in these OPEs (D.27) and (D.28) can be checked from Table 2 or 6. The factor $(k-4)$ appears in (D.28).

The OPEs between the spin-$\frac{3}{2}$ currents and the spin-$\frac{7}{2}$ current $W_+^{(\frac{7}{2})}(w)$ (3.36) are expressed as

$$
\begin{aligned}
\hat{G}_{11}(z)\, W_+^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2} \left[-\frac{18(7+2k)}{5(6+k)} U_+^{(3)} + \frac{8i}{(6+k)} \hat{B}_-\, \hat{T}^{(2)} \right] (w) \\
&+ \quad \frac{1}{(z-w)} \left[\frac{1}{6} \partial \{ \hat{G}_{11}\, W_+^{(\frac{7}{2})} \}_{-2} - \frac{2i}{(6+k)} \hat{B}_-\, T^{(3)} - \frac{10i}{3(6+k)} \hat{B}_- \partial T^{(2)} \right. \\
&+ \quad \frac{20i}{3(6+k)} T^{(2)} \partial \hat{B}_- + \frac{2}{(6+k)} T_+^{(\frac{5}{2})} \hat{G}_{11} - \frac{4i}{(6+k)} U_+^{(3)} \hat{A}_3 \\
&+ \quad \left. \frac{1}{(6+k)} \partial U_+^{(3)} \right] (w) + \cdots, \qquad \text{(D.29)} \\
\hat{G}_{22}(z)\, W_+^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2} \left[\frac{6(29+4k)}{5(6+k)} V_+^{(3)} - \frac{8i}{(6+k)} \hat{A}_-\, \hat{T}^{(2)} \right] (w) \\
&+ \quad \frac{1}{(z-w)} \left[\frac{1}{6} \partial \{ \hat{G}_{22}\, W_+^{(\frac{7}{2})} \}_{-2} + \frac{2i}{(6+k)} \hat{A}_-\, T^{(3)} + \frac{10i}{3(6+k)} \hat{A}_- \partial T^{(2)} \right. \\
&- \quad \frac{20i}{3(6+k)} T^{(2)} \partial \hat{A}_- - \frac{2}{(6+k)} T_+^{(\frac{5}{2})} \hat{G}_{22} + \frac{4i}{(6+k)} V_+^{(3)} \hat{B}_3 \\
&- \quad \left. \frac{1}{(6+k)} \partial V_+^{(3)} \right] (w) + \cdots, \qquad \text{(D.30)} \\
\hat{G}_{12}(z)\, W_+^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^3} \left[\frac{48(-4+k)}{5(6+k)} T^{(2)} \right] (w) \\
&+ \quad \frac{1}{(z-w)^2} \left[-\frac{6(-4+k)}{5(6+k)} T^{(3)} + \frac{(32+6k)}{(6+k)} W^{(3)} + \frac{16i}{(6+k)} (\hat{A}_3 - \hat{B}_3) T^{(2)} \right] (w) \\
&+ \quad \frac{1}{(z-w)} \left[\frac{1}{6} \partial \{ \hat{G}_{12}\, W_+^{(\frac{7}{2})} \}_{-2} - W^{(4)} - \frac{16i}{3(6+k)} (\hat{A}_3 - \hat{B}_3) \partial T^{(2)} \right. \\
&+ \quad \frac{2i}{(6+k)} \hat{A}_- U_-^{(3)} - \frac{2i}{(6+k)} \hat{B}_- V_-^{(3)} + \frac{32i}{3(6+k)} T^{(2)} \partial (\hat{A}_3 - \hat{B}_3) \\
&+ \quad \left. \frac{144(-4+k)}{(324+179k)} \hat{T} T^{(2)} - \frac{2i}{(6+k)} U_+^{(3)} \hat{B}_+ + \frac{2i}{(6+k)} V_+^{(3)} \hat{A}_+ \right.
\end{aligned}
$$

$$
\begin{aligned}
&- \frac{8}{3(6+k)}\partial W^{(3)} - \frac{216(-4+k)}{(1620+895k)}\partial^2 T^{(2)}\Big](w) + \cdots, \qquad \text{(D.31)}\\
\hat{G}_{21}(z)\, W_+^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)}\left[\frac{2i}{(6+k)}\hat{B}_- V_+^{(3)} - \frac{2i}{(6+k)}U_+^{(3)}\hat{A}_-\right](w) + \cdots. \qquad \text{(D.32)}
\end{aligned}
$$

The $U(1)$ charge conservation in these OPEs can be checked from Table 2, 7 or 8. The nonlinear terms appear in (D.29), (D.30), (D.31) and (D.32). Each term in the second order pole of (D.29), (D.30), or (D.31) is a primary field. There is no descendant field for the spin-2 field $T^{(2)}(w)$ in the second order pole of (D.31).

The OPEs between the spin-$\frac{3}{2}$ currents and the spin-$\frac{7}{2}$ current $W_-^{(\frac{7}{2})}(w)$ (3.38) are given by

$$
\begin{aligned}
\hat{G}_{11}(z)\, W_-^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[-\frac{6(29+4k)}{5(6+k)}U_-^{(3)} - \frac{8i}{(6+k)}\hat{T}^{(2)}\hat{A}_+\right](w)\\
&+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{11}\, W_-^{(\frac{7}{2})}\}_{-2} + \frac{10i}{3(6+k)}\hat{A}_+\partial T^{(2)} + \frac{4i}{(6+k)}\hat{B}_3\, U_-^{(3)}\right.\\
&- \frac{2}{(6+k)}\hat{G}_{11}T_-^{(\frac{5}{2})} - \frac{20i}{3(6+k)}T^{(2)}\partial\hat{A}_+ - \frac{2i}{(6+k)}T^{(3)}\hat{A}_+\\
&+ \left.\frac{1}{(6+k)}\partial U_-^{(3)}\right](w) + \cdots, \qquad \text{(D.33)}\\
\hat{G}_{22}(z)\, W_-^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)^2}\left[\frac{18(7+2k)}{5(6+k)}V_-^{(3)} + \frac{8i}{(6+k)}\hat{T}^{(2)}\hat{B}_+\right](w)\\
&+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{22}\, W_-^{(\frac{7}{2})}\}_{-2} - \frac{4i}{(6+k)}\hat{A}_3\, V_-^{(3)} - \frac{10i}{3(6+k)}\hat{B}_+\partial T^{(2)}\right.\\
&+ \frac{2}{(6+k)}\hat{G}_{22}T_-^{(\frac{5}{2})} + \frac{20i}{3(6+k)}T^{(2)}\partial\hat{B}_+ + \frac{2i}{(6+k)}T^{(3)}\hat{B}_+\\
&- \left.\frac{1}{(6+k)}\partial V_-^{(3)}\right](w) + \cdots, \qquad \text{(D.34)}\\
\hat{G}_{12}(z)\, W_-^{(\frac{7}{2})}(w) &= \frac{1}{(z-w)}\left[-\frac{2i}{(6+k)}\hat{A}_+ V_-^{(3)} + \frac{2i}{(6+k)}U_-^{(3)}\hat{B}_+\right](w) + \cdots, \qquad \text{(D.35)}\\
\hat{G}_{21}(z)\, W_-^{(\frac{7}{2})}(w) &= -\frac{1}{(z-w)^3}\left[\frac{48(-4+k)}{5(6+k)}T^{(2)}\right](w)\\
&+ \frac{1}{(z-w)^2}\left[-\frac{6(-4+k)}{5(6+k)}T^{(3)} + \frac{(32+6k)}{(6+k)}W^{(3)} + \frac{16i}{(6+k)}(\hat{A}_3 - \hat{B}_3)T^{(2)}\right](w)\\
&+ \frac{1}{(z-w)}\left[\frac{1}{6}\partial\{\hat{G}_{21}\, W_+^{(\frac{7}{2})}\}_{-2} + W^{(4)}\right.\\
&- \left.\frac{144(-4+k)}{(324+179k)}\left(\hat{T}T^{(2)} - \frac{3}{10}\partial^2 T^{(2)}\right)\right](w) + \cdots. \qquad \text{(D.36)}
\end{aligned}
$$

The $U(1)$ charge conservation in these OPEs (D.33), (D.34), (D.35) and (D.36) can be checked

from Table 2, 7 or 8. The factor $(k-4)$ appears in (D.36). Each term in the second order pole of (D.33), (D.34), or (D.36) is a primary field. There is no descendant field for the spin-2 field $T^{(2)}(w)$ in the second order pole of (D.36).

The OPEs between the spin-$\frac{3}{2}$ currents and the spin-4 current $W^{(4)}(w)$ (3.40) is

$$
\begin{aligned}
\hat{G}_{11}(z)\,W^{(4)}(w) &= \frac{1}{(z-w)^3}\left[\frac{4(-87804+k(-32897+9k(549+166k)))}{3(6+k)^2(324+179k)}U^{(\frac{5}{2})}\right](w)\\
&+ \frac{1}{(z-w)^2}\left[\frac{(36+7k)}{(6+k)}U^{(\frac{7}{2})}+\frac{4i(34+3k)}{3(6+k)^2}\hat{B}_-T_-^{(\frac{5}{2})}\right.\\
&+ \frac{8(-2592-196k+81k^2)}{3(6+k)(324+179k)}\hat{G}_{11}T^{(2)}+\frac{2i(88+15k)}{3(6+k)^2}T_+^{(\frac{5}{2})}\hat{A}_+\\
&+ \frac{4i(72+19k)}{3(6+k)^2}U^{(\frac{5}{2})}\hat{A}_3-\frac{8i(-3+2k)}{3(6+k)^2}U^{(\frac{5}{2})}\hat{B}_3\\
&+ \left.\frac{4(57672+33962k+4327k^2-108k^3)}{5(6+k)^2(324+179k)}\partial U^{(\frac{5}{2})}\right](w)+\cdots,\\
&+ \frac{1}{(z-w)}\left[\frac{1}{7}\{\hat{G}_{11}\,W^{(4)}\}_{-2}-\frac{8i(53+24k)}{35(6+k)^2}\hat{A}_3\partial U^{(\frac{5}{2})}-\frac{4i(31+6k)}{7(6+k)^2}\hat{A}_+\partial T_+^{(\frac{5}{2})}\right.\\
&- \frac{8i(47+6k)}{35(6+k)^2}\hat{B}_3\partial U^{(\frac{5}{2})}-\frac{2i}{(6+k)}\hat{B}_-W_-^{(\frac{7}{2})}+\frac{8}{(6+k)^2}\hat{B}_-\hat{A}_3T_-^{(\frac{5}{2})}\\
&- \frac{4}{(6+k)^2}\hat{B}_-\hat{A}_+V^{(\frac{5}{2})}-\frac{8i(68+13k)}{35(6+k)^2}\hat{B}_-\partial T_-^{(\frac{5}{2})}-\frac{4}{(6+k)}\hat{G}_{11}T^{(3)}\\
&- \frac{24(-540-161k+9k^2)}{7(6+k)(324+179k)}\hat{G}_{11}\partial T^{(2)}-\frac{2}{(6+k)}\hat{G}_{21}U_-^{(3)}\\
&+ \frac{32(-540-161k+9k^2)}{7(6+k)(324+179k)}T^{(2)}\partial\hat{G}_{11}+\frac{4i(61+13k)}{7(6+k)^2}T_-^{(\frac{5}{2})}\partial\hat{B}_-\\
&+ \frac{4i(88+15k)}{7(6+k)^2}T_+^{(\frac{5}{2})}\partial\hat{A}_++\frac{144(-4+k)}{(324+179k)}U^{(\frac{5}{2})}\hat{T}-\frac{16}{(6+k)^2}U^{(\frac{5}{2})}\hat{A}_3\hat{A}_3\\
&+ \frac{16}{(6+k)^2}U^{(\frac{5}{2})}\hat{A}_3\hat{B}_3-\frac{4}{(6+k)^2}U^{(\frac{5}{2})}\hat{A}_-\hat{A}_++\frac{8i(79+12k)}{7(6+k)^2}U^{(\frac{5}{2})}\partial\hat{A}_3\\
&+ \frac{8i(13+3k)}{7(6+k)^2}U^{(\frac{5}{2})}\partial\hat{B}_3+\frac{4i}{(6+k)}U^{(\frac{7}{2})}(\hat{A}_3-\hat{B}_3)+\frac{34}{7(6+k)}\partial U^{(\frac{7}{2})}\\
&+ \left.\frac{(239760+90124k+5112k^2-972k^3)}{21(6+k)^2(324+179k)}\partial^2U^{(\frac{5}{2})}\right](w)+\cdots, \qquad \text{(D.37)}\\
\hat{G}_{22}(z)\,W^{(4)}(w) &= \frac{1}{(z-w)^3}\left[\frac{4(-71604+k(-21031+18k(364+83k)))}{3(6+k)^2(324+179k)}V^{(\frac{5}{2})}\right](w)\\
&+ \frac{1}{(z-w)^2}\left[-\frac{(36+7k)}{(6+k)}V^{(\frac{7}{2})}-\frac{8i(-9+k)}{3(6+k)^2}\hat{A}_3V^{(\frac{5}{2})}\right.\\
&- \left.\frac{8i(11+3k)}{3(6+k)^2}\hat{A}_-T_-^{(\frac{5}{2})}-\frac{4i(96+13k)}{3(6+k)^2}\hat{B}_3V^{(\frac{5}{2})}\right.
\end{aligned}
$$

$$
\begin{aligned}
&+ \frac{8(-1296+520k+81k^2)}{3(6+k)(324+179)}T^{(2)}\hat{G}_{22} - \frac{2i(88+15k)}{3(6+k)^2}T_+^{(\frac{5}{2})}\hat{B}_+ \\
&+ \frac{2(-184680-48354k+1153k^2+972k^3)}{15(6+k)^2(324+179k)}\partial V^{(\frac{5}{2})}\Big](w) \\
&+ \frac{1}{(z-w)}\Big[\frac{1}{7}\{\hat{G}_{22}\,W^{(4)}\}_{-2} + \frac{4i}{(6+k)}\hat{A}_3V^{(\frac{7}{2})} - \frac{16}{(6+k)^2}\hat{A}_3\hat{B}_3V^{(\frac{5}{2})} \\
&+ \frac{32i(5+k)}{35(6+k)^2}\hat{A}_3\partial V^{(\frac{5}{2})} - \frac{2i}{(6+k)}\hat{A}_-W_-^{(\frac{7}{2})} - \frac{8}{(6+k)^2}\hat{A}_-\hat{B}_3T_-^{(\frac{5}{2})} \\
&+ \frac{4i(109+24k)}{35(6+k)^2}\hat{A}_-\partial T_-^{(\frac{5}{2})} - \frac{4i}{(6+k)}\hat{B}_3V^{(\frac{7}{2})} + \frac{16}{(6+k)^2}\hat{B}_3\hat{B}_3V^{(\frac{5}{2})} \\
&+ \frac{16i(110+13k)}{35(6+k)^2}\hat{B}_3\partial V^{(\frac{5}{2})} + \frac{4}{(6+k)^2}\hat{B}_-V^{(\frac{5}{2})}\hat{B}_+ + \frac{4i(31+6k)}{7(6+k)^2}\hat{B}_+\partial T_+^{(\frac{5}{2})} \\
&+ \frac{2}{(6+k)}\hat{G}_{21}V_-^{(3)} - \frac{24(108+197k+9k^2)}{7(6+k)(324+179k)}\hat{G}_{22}\partial T^{(2)} \\
&+ \frac{4}{(6+k)}T^{(3)}\hat{G}_{22} + \frac{32(108+197k+9k^2)}{7(6+k)(324+179k)}T^{(2)}\partial\hat{G}_{22} - \frac{4i(79+12k)}{7(6+k)^2}T_-^{(\frac{5}{2})}\partial\hat{A}_- \\
&- \frac{4i(88+15k)}{7(6+k)^2}T_+^{(\frac{5}{2})}\partial\hat{B}_+ + \frac{144(-4+k)}{(324+179k)}\hat{T}V^{(\frac{5}{2})} + \frac{4}{(6+k)^2}U^{(\frac{5}{2})}\hat{A}_-\hat{B}_+ \\
&- \frac{8i(17+2k)}{7(6+k)^2}V^{(\frac{5}{2})}\partial\hat{A}_3 - \frac{8i(61+13k)}{7(6+k)^2}V^{(\frac{5}{2})}\partial\hat{B}_3 + \frac{22}{7(6+k)}\partial V^{(\frac{7}{2})} \\
&- \frac{4(27540+36371k+3771k^2+270k^3)}{105(6+k)^2(324+179k)}\partial^2V^{(\frac{5}{2})}\Big](w) + \cdots, \qquad \text{(D.38)}
\end{aligned}
$$

$$
\begin{aligned}
\hat{G}_{12}(z)\,W^{(4)}(w) &= \frac{1}{(z-w)^3}\left[\frac{8(-41472-13972k+3097k^2+747k^3)}{3(6+k)^2(324+179k)}T_-^{(\frac{5}{2})}\right](w) \\
&+ \frac{1}{(z-w)^2}\Big[\frac{(116+21k)}{3(6+k)}W_-^{(\frac{7}{2})} + \frac{8i(27+5k)}{3(6+k)^2}\hat{A}_3T_-^{(\frac{5}{2})} - \frac{2i(28+5k)}{(6+k)^2}\hat{A}_+V^{(\frac{5}{2})} \\
&- \frac{8i(27+5k)}{3(6+k)^2}\hat{B}_3T_-^{(\frac{5}{2})} + \frac{216(-4+k)}{(324+179k)}T^{(2)}\hat{G}_{12} + \frac{2i(28+5k)}{(6+k)^2}U^{(\frac{5}{2})}\hat{B}_+ \\
&+ \frac{2(-92016-37048k-1883k^2+324k^3)}{5(6+k)^2(324+179k)}\partial T_-^{(\frac{5}{2})}\Big](w) \\
&+ \frac{1}{(z-w)}\Big[\frac{1}{7}\{\hat{G}_{12}\,W^{(4)}\}_{-2} - \frac{8}{(6+k)^2}\hat{A}_3\hat{A}_+V^{(\frac{5}{2})} - \frac{8i(23+4k)}{7(6+k)^2}\hat{A}_3\partial T_-^{(\frac{5}{2})} \\
&- \frac{4}{(6+k)^2}\hat{A}_-\hat{A}_+T_-^{(\frac{5}{2})} + \frac{4i(35+6k)}{7(6+k)^2}\hat{A}_+\partial V^{(\frac{5}{2})} + \frac{8}{(6+k)^2}\hat{B}_3\hat{A}_+V^{(\frac{5}{2})} \\
&+ \frac{16i(15+2k)}{7(6+k)^2}\hat{B}_3\partial T_-^{(\frac{5}{2})} + \frac{4}{(6+k)^2}\hat{B}_-T_-^{(\frac{5}{2})}\hat{B}_+ - \frac{8i(14+3k)}{7(6+k)^2}\hat{B}_+\partial U^{(\frac{5}{2})} \\
&- \frac{2}{(6+k)}\hat{G}_{11}V_-^{(3)} - \frac{216(-4+k)}{7(324+179k)}\hat{G}_{12}\partial T^{(2)} + \frac{288(-4+k)}{7(324+179k)}T^{(2)}\partial\hat{G}_{12}
\end{aligned}
$$

$$
\begin{aligned}
&+ \frac{16i(34+5k)}{7(6+k)^2}T_-^{(\frac{5}{2})}\partial\hat{A}_3 - \frac{8i(61+10k)}{7(6+k)^2}T_-^{(\frac{5}{2})}\partial\hat{B}_3 + \frac{144(-4+k)}{(324+179k)}\hat{T}T_-^{(\frac{5}{2})} \\
&+ \frac{2}{(6+k)}U_-^{(3)}\hat{G}_{22} - \frac{8}{(6+k)^2}U^{(\frac{5}{2})}(\hat{A}_3-\hat{B}_3)\hat{B}_+ + \frac{12i(28+5k)}{7(6+k)^2}U^{(\frac{5}{2})}\partial\hat{B}_+ \\
&- \frac{4i(91+15k)}{7(6+k)^2}V^{(\frac{5}{2})}\partial\hat{A}_+ - \frac{6}{7(6+k)}\partial W_-^{(\frac{7}{2})} \\
&+ \frac{(804168+498326k+61528k^2-1080k^3)}{105(6+k)^2(324+179k)}\partial^2 T_-^{(\frac{5}{2})}\Big](w) + \cdots, \qquad \text{(D.39)}
\end{aligned}
$$

$$
\begin{aligned}
\hat{G}_{21}(z)\,W^{(4)}(w) &= \frac{1}{(z-w)^3}\left[\frac{4(-84240-28336k+6373k^2+1494k^3)}{3(6+k)^2(324+179k)}T_+^{(\frac{5}{2})}\right](w) \\
&+ \frac{1}{(z-w)^2}\Big[-\frac{(124+21k)}{3(6+k)}W_+^{(\frac{7}{2})} - \frac{6i(20+3k)}{(6+k)^2}\hat{B}_-V^{(\frac{5}{2})} \\
&+ \frac{216(-4+k)}{(324+179k)}\hat{G}_{21}T^{(2)} - \frac{8i(36+7k)}{3(6+k)^2}T_+^{(\frac{5}{2})}(\hat{A}_3-\hat{B}_3) + \frac{2i(52+11k)}{(6+k)^2}U^{(\frac{5}{2})}\hat{A}_- \\
&+ \frac{(235872+135472k+15518k^2-432k^3)}{5(6+k)^2(324+179k)}\partial T_+^{(\frac{5}{2})}\Big](w) \\
&+ \frac{1}{(z-w)}\Big[\frac{1}{7}\{\hat{G}_{21}\,W^{(4)}\}_{-2} + \frac{40i}{7(6+k)^2}\hat{A}_3\partial T_+^{(\frac{5}{2})} - \frac{12i(41+8k)}{35(6+k)^2}\hat{A}_-\partial U^{(\frac{5}{2})} \\
&- \frac{40i}{7(6+k)^2}\hat{B}_3\partial T_+^{(\frac{5}{2})} - \frac{2i}{(6+k)}\hat{B}_-V^{(\frac{7}{2})} - \frac{8}{(6+k)^2}\hat{B}_-(\hat{A}_3-\hat{B}_3)V^{(\frac{5}{2})} \\
&+ \frac{4i(115+26k)}{35(6+k)^2}\hat{B}_-\partial V^{(\frac{5}{2})} - \frac{216(-4+k)}{7(324+179k)}\hat{G}_{21}\partial T^{(2)} \\
&+ \frac{288(-4+k)}{7(324+179k)}T^{(2)}\partial\hat{G}_{21} + \frac{144(-4+k)}{(324+179k)}T_+^{(\frac{5}{2})}\hat{T} - \frac{16}{7(6+k)^2}T_+^{(\frac{5}{2})}\partial\hat{A}_3 \\
&+ \frac{16i}{7(6+k)^2}T_+^{(\frac{5}{2})}\partial\hat{B}_3 - \frac{8i}{(6+k)^2}U^{(\frac{5}{2})}\hat{A}_-(\hat{A}_3-\hat{B}_3) + \frac{8i(29+6k)}{7(6+k)^2}U^{(\frac{5}{2})}\partial\hat{A}_- \\
&+ \frac{2i}{(6+k)}U^{(\frac{7}{2})}\hat{A}_- - \frac{4i(68+13k)}{7(6+k)^2}V^{(\frac{5}{2})}\partial\hat{B}_- - \frac{10}{7(6+k)}\partial W_+^{(\frac{7}{2})} \\
&+ \frac{(1625832+736438k+53126k^2-4860k^3)}{105(6+k)^2(324+179k)}\partial^2 T_+^{(\frac{5}{2})}\Big](w) + \cdots. \qquad \text{(D.40)}
\end{aligned}
$$

One can check the $U(1)$ charges from Table 2, 9 or 10 and the nonlinear terms (linear in higher spin currents) appear in (D.37), (D.38), (D.39) and (D.40). There are no descendant fields for the spin-$\frac{5}{2}$ fields in the second order pole of (D.37), (D.38), (D.39) or (D.40). One can analyze each second-order pole in these OPEs whether it is written in terms of (quasi)primary fields.

Therefore, one obtains the OPEs between six spin-1 currents and the higher spin current of spins $(3, \frac{7}{2}, \frac{7}{2}, 4)$ explicitly. The higher spin currents corresponding to three $\mathcal{N}=2$ multiplets (3.1) occur in the right hand sides of the OPEs.

As described in the previous Appendix, one can analyze the above OPEs similarly. The

first result is as follows:

$$\hat{G}_{11} \quad \times \quad \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{5}{2})}, U^{(\frac{7}{2})} \\ 0, & 0, & 0, & 0 \\ T^{(2)}, W^{(3)}, & T_+^{(\frac{5}{2})}, W_+^{(\frac{7}{2})}, & T_-^{(\frac{5}{2})}, W_-^{(\frac{7}{2})}, & T^{(2)}, T^{(3)}, W^{(3)}, W^{(4)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{5}{2})}, U^{(\frac{7}{2})} \end{pmatrix}. \qquad \text{(D.41)}$$

There are no higher spin currents, denoted by zeros in (D.41) and the four currents living in the second multiplet of (3.1) appear in the first and last rows in (D.41). Also the current $U^{(\frac{5}{2})}(w)$ appears in the last columns of these rows. The first and last multiplets of (3.1) appear in the third row in (D.41). The currents $T^{(2)}(w)$ and $W^{(3)}(w)$ can appear in the last column of this row.

Similarly, it turns out that the following result for the spin-$\frac{3}{2}$ current and 16 currents is given by

$$\hat{G}_{12} \quad \times \quad \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} T_-^{(\frac{5}{2})}, & T^{(2)}, T^{(3)}, & 0, & T_-^{(\frac{5}{2})} \\ U_-^{(3)}, & U^{(\frac{5}{2})}, U^{(\frac{7}{2})}, & 0, & U_-^{(3)} \\ V_-^{(3)}, & V^{(\frac{5}{2})}, V^{(\frac{7}{2})}, & 0, & V_-^{(3)} \\ T_-^{(\frac{5}{2})}, W_-^{(\frac{7}{2})}, & T^{(2)}, T^{(3)}, W^{(3)}, W^{(4)}, & 0, & T_-^{(\frac{5}{2})}, W_-^{(\frac{7}{2})} \end{pmatrix}. \qquad \text{(D.42)}$$

The four currents living in the third component of each multiplet of (3.1) appear in the first and the last columns in (D.42). Also the current $T_-^{(\frac{5}{2})}(w)$ appears in the last rows of these columns. The first and last components in each multiplet of (3.1) appear in the second column in (D.42). The currents $T^{(2)}(w)$ and $W^{(3)}(w)$ can appear in the last row of this column.

One has the following result

$$\hat{G}_{21} \quad \times \quad \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} T_+^{(\frac{5}{2})}, & 0, & T^{(2)}, T^{(3)}, & T_+^{(\frac{5}{2})} \\ U_+^{(3)}, & 0, & U^{(\frac{5}{2})}, U^{(\frac{7}{2})}, & U_+^{(3)} \\ V_+^{(3)}, & 0, & V^{(\frac{5}{2})}, V^{(\frac{7}{2})}, & V_+^{(3)} \\ T_+^{(\frac{5}{2})}, W_+^{(\frac{7}{2})}, & 0, & T^{(2)}, T^{(3)}, W^{(3)}, W^{(4)}, & T_+^{(\frac{5}{2})}, W_+^{(\frac{7}{2})} \end{pmatrix}. \tag{D.43}$$

The four currents living in the second component of each multiplet of (3.1) appear in the first and the last columns in (D.43). Also the current $T_+^{(\frac{5}{2})}(w)$ appears in the last rows of these columns. The first and last components in each multiplet of (3.1) appear in the third column in (D.43). The currents $T^{(2)}(w)$ and $W^{(3)}(w)$ can appear in the last row of this column.

Finally, one obtains the following result

$$\begin{aligned} \hat{G}_{22} \quad \times & \begin{pmatrix} T^{(2)}, & T_+^{(\frac{5}{2})}, & T_-^{(\frac{5}{2})}, & T^{(3)} \\ U^{(\frac{5}{2})}, & U_+^{(3)}, & U_-^{(3)}, & U^{(\frac{7}{2})} \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{7}{2})} \\ W^{(3)}, & W_+^{(\frac{7}{2})}, & W_-^{(\frac{7}{2})}, & W^{(4)} \end{pmatrix} \\ \rightarrow & \begin{pmatrix} V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{5}{2})}, V^{(\frac{7}{2})} \\ T^{(2)}, W^{(3)}, & T_+^{(\frac{5}{2})}, W_+^{(\frac{7}{2})}, & T_-^{(\frac{5}{2})}, W_-^{(\frac{7}{2})}, & T^{(2)}, T^{(3)}, W^{(3)}, W^{(4)} \\ 0, & 0, & 0, & 0 \\ V^{(\frac{5}{2})}, & V_+^{(3)}, & V_-^{(3)}, & V^{(\frac{5}{2})}, V^{(\frac{7}{2})} \end{pmatrix}. \end{aligned} \tag{D.44}$$

The four currents living in the third multiplet of (3.1) appear in the first and last rows in (D.44). Also the current $V^{(\frac{5}{2})}(w)$ appears in the last columns of these rows. The first and last multiplets of (3.1) appear in the second row in (D.41). The currents $T^{(2)}(w)$ and $W^{(3)}(w)$ can appear in the last column of this row.

References

[1] M. R. Gaberdiel and R. Gopakumar, "Large $\mathcal{N} = 4$ Holography," JHEP **1309**, 036 (2013) [arXiv:1305.4181 [hep-th]].

[2] M. R. Gaberdiel and R. Gopakumar, "An AdS_3 Dual for Minimal Model CFTs," Phys. Rev. D **83**, 066007 (2011) [arXiv:1011.2986 [hep-th]].

[3] M. R. Gaberdiel and R. Gopakumar, "Triality in Minimal Model Holography," JHEP **1207**, 127 (2012) [arXiv:1205.2472 [hep-th]].

[4] A. Sevrin and G. Theodoridis, "N=4 Superconformal Coset Theories," Nucl. Phys. B **332**, 380 (1990).

[5] J. A. Wolf, "Complex Homogeneous Contact Manifolds and Quaternionic Symmetric Spaces," J. Math. Mech. **14**, 1033 (1965).

[6] D. V. Alekseevskii, "Classification of Quarternionic Spaces with a Transitive Solvable Group of Motions," Math. USSR Izv. **9**, 297 (1975).

[7] S. Salamon, "Quaternionic Kahler Manifolds," Invent. Math. **67**, 143 (1982).

[8] T. Creutzig, Y. Hikida and P. B. Ronne, "Higher spin AdS_3 supergravity and its dual CFT," JHEP **1202**, 109 (2012) [arXiv:1111.2139 [hep-th]].

[9] C. Candu and M. R. Gaberdiel, "Supersymmetric holography on AdS_3," JHEP **1309**, 071 (2013) [arXiv:1203.1939 [hep-th]].

[10] K. Hanaki and C. Peng, "Symmetries of Holographic Super-Minimal Models," JHEP **1308**, 030 (2013) [arXiv:1203.5768 [hep-th]].

[11] C. Ahn, "The Large N 't Hooft Limit of Kazama-Suzuki Model," JHEP **1208**, 047 (2012) [arXiv:1206.0054 [hep-th]].

[12] C. Candu and M. R. Gaberdiel, "Duality in N=2 Minimal Model Holography," JHEP **1302**, 070 (2013) [arXiv:1207.6646 [hep-th]].

[13] C. Ahn, "The Operator Product Expansion of the Lowest Higher Spin Current at Finite N," JHEP **1301**, 041 (2013) [arXiv:1208.0058 [hep-th]].

[14] Y. Hikida, "Conical defects and $N = 2$ higher spin holography," JHEP **1308**, 127 (2013) [arXiv:1212.4124].

[15] C. Candu and C. Vollenweider, "On the coset duals of extended higher spin theories," JHEP **1404**, 145 (2014) [arXiv:1312.5240 [hep-th]].

[16] C. Candu, C. Peng and C. Vollenweider, "Extended supersymmetry in AdS_3 higher spin theories," arXiv:1408.5144 [hep-th].

[17] C. M. Hull and B. J. Spence, "$N = 2$ Current Algebra and Coset Models," Phys. Lett. B **241**, 357 (1990).

[18] M. Rocek, C. Ahn, K. Schoutens and A. Sevrin, "Superspace WZW models and black holes," hep-th/9110035.

[19] C. Ahn, E. Ivanov and A. S. Sorin, "N=2 affine superalgebras and Hamiltonian reduction in N=2 superspace," Commun. Math. Phys. **183**, 205 (1997) [hep-th/9508005].

[20] C. Ahn, "Higher Spin Currents in Wolf Space. Part I," JHEP **1403**, 091 (2014) [arXiv:1311.6205 [hep-th]].

[21] A. Sevrin, W. Troost and A. Van Proeyen, "Superconformal Algebras in Two-Dimensions with N=4," Phys. Lett. B **208**, 447 (1988).

[22] A. Sevrin, W. Troost, A. Van Proeyen and P. Spindel, "EXTENDED SUPERSYMMETRIC sigma MODELS ON GROUP MANIFOLDS. 2. CURRENT ALGEBRAS," Nucl. Phys. B **311**, 465 (1988).

[23] K. Schoutens, "O(n) Extended Superconformal Field Theory in Superspace," Nucl. Phys. B **295**, 634 (1988).

[24] E. A. Ivanov and S. O. Krivonos, "N=4 Superliouville Equation. (in Russian)," J. Phys. A **17**, L671 (1984).

[25] E. A. Ivanov and S. O. Krivonos, "$N = 4$ Superextension of the Liouville Equation With Quaternionic Structure," Theor. Math. Phys. **63**, 477 (1985) [Teor. Mat. Fiz. **63**, 230 (1985)].

[26] E. A. Ivanov, S. O. Krivonos and V. M. Leviant, "A New Class of Superconformal σ Models With the Wess-Zumino Action," Nucl. Phys. B **304**, 601 (1988).

[27] E. A. Ivanov, S. O. Krivonos and V. M. Leviant, "Quantum N=3, N=4 Superconformal WZW Sigma Models," Phys. Lett. B **215**, 689 (1988) [Erratum-ibid. B **221**, 432 (1989)].

[28] P. Goddard and A. Schwimmer, "Factoring Out Free Fermions And Superconformal Algebras," Phys. Lett. B **214**, 209 (1988).

[29] A. Van Proeyen, "Realizations Of N=4 Superconformal Algebras On Wolf Spaces," Class. Quant. Grav. **6**, 1501 (1989).

[30] M. Gunaydin, J. L. Petersen, A. Taormina and A. Van Proeyen, "On The Unitary Representations Of A Class Of N=4 Superconformal Algebras," Nucl. Phys. B **322**, 402 (1989).

[31] S. J. Gates, Jr. and S. V. Ketov, "No N=4 strings on wolf spaces," Phys. Rev. D **52**, 2278 (1995) [hep-th/9501140].

[32] C. Ahn, "Higher Spin Currents in Wolf Space: Part II," arXiv:1408.0655 [hep-th].

[33] M. R. Gaberdiel and T. Hartman, "Symmetries of Holographic Minimal Models," JHEP **1105**, 031 (2011) [arXiv:1101.2910 [hep-th]].

[34] M. R. Gaberdiel and C. Peng, "The symmetry of large $\mathcal{N} = 4$ holography," JHEP **1405**, 152 (2014) [arXiv:1403.2396 [hep-th]].

[35] M. Beccaria, C. Candu and M. R. Gaberdiel, "The large N = 4 superconformal W_∞ algebra," JHEP **1406**, 117 (2014) [arXiv:1404.1694 [hep-th]].

[36] Y. Kazama and H. Suzuki, "New N=2 Superconformal Field Theories and Superstring Compactification," Nucl. Phys. B **321**, 232 (1989).

[37] Y. Kazama and H. Suzuki, "Characterization of N=2 Superconformal Models Generated by Coset Space Method," Phys. Lett. B **216**, 112 (1989).

[38] T. Creutzig, Y. Hikida and P. B. Ronne, "Higher spin AdS_3 holography with extended supersymmetry," arXiv:1406.1521 [hep-th].

[39] C. Ahn, "The Higher Spin Currents in the N=1 Stringy Coset Minimal Model," JHEP **1304**, 033 (2013) [arXiv:1211.2589 [hep-th]].

[40] C. Ahn, "Higher Spin Currents with Arbitrary N in the $\mathcal{N} = 1$ Stringy Coset Minimal Model," JHEP **1307**, 141 (2013) [arXiv:1305.5892 [hep-th]].

[41] M. Beccaria, C. Candu, M. R. Gaberdiel and M. Groher, "$\mathcal{N} = 1$ extension of minimal model holography," arXiv:1305.1048 [hep-th].

[42] M. R. Gaberdiel and R. Gopakumar, "Higher Spins & Strings," arXiv:1406.6103 [hep-th].

[43] T. Creutzig, Y. Hikida and P. B. Ronne, "Extended higher spin holography and Grassmannian models," JHEP **1311**, 038 (2013) [arXiv:1306.0466 [hep-th]].

[44] M. R. Gaberdiel and R. Gopakumar, "Minimal Model Holography," J. Phys. A **46**, 214002 (2013) [arXiv:1207.6697 [hep-th]].

[45] T. Creutzig, Y. Hikida and P. B. Ronne, "N=1 supersymmetric higher spin holography on AdS_3," JHEP **1302**, 019 (2013) [arXiv:1209.5404 [hep-th]].

[46] C. Ahn and J. Paeng, "Higher Spin Currents in the Holographic $\mathcal{N} = 1$ Coset Minimal Model," JHEP **1401**, 007 (2014) [arXiv:1310.6185 [hep-th]].

[47] C. Ahn, "Explicit construction of N=2 W(3) current in the N=2 coset SU(3) / Su(2) x U(1) model," Phys. Lett. B **348**, 77 (1995) [hep-th/9410170].

[48] L. Anguelova, M. Rocek and S. Vandoren, "Hyperkahler cones and orthogonal Wolf spaces," JHEP **0205**, 064 (2002) [hep-th/0202149].

[49] F. A. Bais, P. Bouwknegt, M. Surridge and K. Schoutens, "Extensions of the Virasoro Algebra Constructed from Kac-Moody Algebras Using Higher Order Casimir Invariants," Nucl. Phys. B **304**, 348 (1988).

[50] F. A. Bais, P. Bouwknegt, M. Surridge and K. Schoutens, "Coset Construction for Extended Virasoro Algebras," Nucl. Phys. B **304**, 371 (1988).

[51] P. Bouwknegt and K. Schoutens, "W symmetry in conformal field theory," Phys. Rept. **223**, 183 (1993) [hep-th/9210010].

[52] C. Ahn, "The Large N 't Hooft Limit of Coset Minimal Models," JHEP **1110**, 125 (2011) [arXiv:1106.0351 [hep-th]].

[53] M. R. Gaberdiel and C. Vollenweider, "Minimal Model Holography for SO(2N)," JHEP **1108**, 104 (2011) [arXiv:1106.2634 [hep-th]].

[54] C. Ahn, "The Primary Spin-4 Casimir Operators in the Holographic SO(N) Coset Minimal Models," JHEP **1205**, 040 (2012) [arXiv:1202.0074 [hep-th]].

[55] C. Candu, M. R. Gaberdiel, M. Kelm and C. Vollenweider, "Even spin minimal model holography," JHEP **1301**, 185 (2013) [arXiv:1211.3113 [hep-th]].

[56] C. Ahn and J. Paeng, "The OPEs of Spin-4 Casimir Currents in the Holographic $SO(N)$ Coset Minimal Models," Class. Quant. Grav. **30**, 175004 (2013) [arXiv:1301.0208].

[57] M. Henneaux, G. Lucena Gomez, J. Park and S. -J. Rey, "Super- W(infinity) Asymptotic Symmetry of Higher-Spin AdS_3 Supergravity," JHEP **1206**, 037 (2012) [arXiv:1203.5152 [hep-th]].

[58] K. Thielemans, "A Mathematica package for computing operator product expansions," Int. J. Mod. Phys. C **2**, 787 (1991).

[59] R. Slansky, Phys. Rep. 79 (1981) 1; J. Patera and D. Sankoff, Tables of Branch- ing rules for Representations of Simple Lie Algebras (LUniversite de Montreal, Montreal, 1973

); W. MacKay and J. Petera, Tables of Dimensions, Indices and Branching Rules for representations of Simple Algebras (Dekker, New York, 1981).

[60] V. G. Knizhnik, "Superconformal Algebras in Two-dimensions," Theor. Math. Phys. **66**, 68 (1986) [Teor. Mat. Fiz. **66**, 102 (1986)].

[61] M. A. Bershadsky, "Superconformal Algebras in Two-dimensions With Arbitrary N," Phys. Lett. B **174**, 285 (1986).

[62] C. Ahn, "The Coset Spin-4 Casimir Operator and Its Three-Point Functions with Scalars," JHEP **1202**, 027 (2012) [arXiv:1111.0091 [hep-th]].

[63] T. Creutzig, Y. Hikida and P. B. Ronne, "Three point functions in higher spin AdS_3 supergravity," JHEP **1301**, 171 (2013) [arXiv:1211.2237 [hep-th]].

[64] H. Moradi and K. Zoubos, "Three-Point Functions in N=2 Higher-Spin Holography," JHEP **1304**, 018 (2013) [arXiv:1211.2239 [hep-th]].

[65] C. Ahn and H. Kim, "Spin-5 Casimir operator its three-point functions with two scalars," JHEP **1401**, 012 (2014) [arXiv:1308.1726 [hep-th]].

[66] S. F. Prokushkin and M. A. Vasiliev, "Higher spin gauge interactions for massive matter fields in 3-D AdS space-time," Nucl. Phys. B **545**, 385 (1999) [hep-th/9806236].

[67] S. Prokushkin and M. A. Vasiliev, "3-d higher spin gauge theories with matter," hep-th/9812242.

[68] H. Georgi, "Lie Algebras In Particle Physics. From Isospin To Unified Theories,", 2nd Edition(1999), Front. Phys. **54**, 1 (1982).

[69] Philippe Di Francesco, Pierre Mathieu, and David Senrchal "Conformal Field Theory," Springer-Verlag (1997).

[70] Robert N. Cahn "Semi-Simple Lie Algebras and Their Representations,"THE BENJAMIN/CUMMINGS PUBLISHING COMPANY (1984).

$U(1)$ charge	Composite fields of spin-$\frac{7}{2}$
$\frac{(4+3k)}{(6+k)}$	$\hat{B}_-T_+^{(\frac{5}{2})}$, $\hat{B}_-\hat{B}_-\hat{G}_{22}$, $\hat{B}_-\hat{G}_{11}\hat{A}_-$, $\hat{B}_-\hat{G}_{21}\hat{A}_3$, $\hat{B}_-\hat{G}_{21}\hat{B}_3$, $\hat{B}_-\partial\hat{G}_{21}$, $\hat{G}_{21}\partial\hat{B}_-$
$\frac{(-4+3k)}{(6+k)}$	$\hat{B}_-U^{(\frac{5}{2})}$, $\hat{B}_-\hat{B}_-\hat{G}_{12}$, $\hat{B}_-\hat{G}_{11}\hat{A}_3$, $\hat{B}_-\hat{G}_{11}\hat{B}_3$, $\hat{B}_-\hat{G}_{21}\hat{A}_+$, $\hat{B}_-\partial\hat{G}_{11}$, $\hat{G}_{11}\partial\hat{B}_-$
$\frac{(12+k)}{(6+k)}$	$\hat{A}_-\partial\hat{G}_{21}$, $\hat{B}_-\hat{A}_-\hat{G}_{22}$, $\hat{G}_{11}\hat{A}_-\hat{A}_-$, $\hat{G}_{21}\hat{A}_-\hat{A}_3$, $\hat{G}_{21}\hat{A}_-\hat{B}_3$, $\hat{G}_{21}\partial\hat{A}_-$, $T_+^{(\frac{5}{2})}\hat{A}_-$
$\frac{(-12+k)}{(6+k)}$	$\hat{A}_+\partial\hat{G}_{11}$, $\hat{B}_-\hat{A}_+\hat{G}_{12}$, $\hat{G}_{11}\hat{A}_3\hat{A}_+$, $\hat{G}_{11}\hat{B}_3\hat{A}_+$, $\hat{G}_{11}\partial\hat{A}_+$, $\hat{G}_{21}\hat{A}_+\hat{A}_+$, $U^{(\frac{5}{2})}\hat{A}_+$
$\frac{(4+k)}{(6+k)}$	$W_+^{(\frac{7}{2})}$, $\hat{A}_3\partial\hat{G}_{21}$, $\hat{A}_-\partial\hat{G}_{11}$, $\hat{B}_3\partial\hat{G}_{21}$, $\hat{B}_-V^{(\frac{5}{2})}$, $\hat{B}_-\hat{A}_3\hat{G}_{22}$ $\hat{B}_-\hat{A}_-\hat{G}_{12}$, $\hat{B}_-\hat{B}_3\hat{G}_{22}$, $\hat{B}_-\hat{G}_{21}\hat{B}_+$, $\hat{B}_-\partial\hat{G}_{22}$, $\hat{G}_{11}\hat{A}_-\hat{A}_3$, $\hat{G}_{11}\hat{A}_-\hat{B}_3$, $\hat{G}_{11}\partial\hat{A}_-$, $\hat{G}_{21}T^{(2)}$, $\hat{G}_{21}\hat{T}$, $\hat{G}_{21}\hat{A}_3\hat{A}_3$, $\hat{G}_{21}\hat{A}_3\hat{B}_3$, $\hat{G}_{21}\hat{A}_-\hat{A}_+$, $\hat{G}_{21}\hat{B}_3\hat{B}_3$, $\hat{G}_{21}\partial\hat{A}_3$, $\hat{G}_{21}\partial\hat{B}_3$, $\hat{G}_{22}\partial\hat{B}_-$, $T_+^{(\frac{5}{2})}\hat{A}_3$, $T_+^{(\frac{5}{2})}\hat{B}_3$, $U^{(\frac{5}{2})}\hat{A}_-$, $\partial T_+^{(\frac{5}{2})}$, $\partial^2\hat{G}_{21}$
$\frac{(-4+k)}{(6+k)}$	$U^{(\frac{7}{2})}$, $\hat{A}_3\partial\hat{G}_{11}$, $\hat{A}_+\partial\hat{G}_{21}$, $\hat{B}_3\partial\hat{G}_{11}$, $\hat{B}_-\partial T_-^{(\frac{5}{2})}$, $\hat{B}_-\hat{A}_3\hat{G}_{12}$, $\hat{B}_-\hat{A}_+\hat{G}_{22}$, $\hat{B}_-\hat{B}_3\hat{G}_{12}$, $\hat{B}_-\hat{G}_{11}\hat{B}_+$, $\hat{B}_-\partial\hat{G}_{12}$, $\hat{G}_{11}T^{(2)}$, $\hat{G}_{11}\hat{T}$, $\hat{G}_{11}\hat{A}_3\hat{A}_3$, $\hat{G}_{11}\hat{A}_3\hat{B}_3$, $\hat{G}_{11}\hat{A}_-\hat{A}_+$, $\hat{G}_{11}\hat{B}_3\hat{B}_3$, $\hat{G}_{11}\partial\hat{A}_3$, $\hat{G}_{11}\partial\hat{B}_3$, $\hat{G}_{12}\partial\hat{B}_-$, $\hat{G}_{21}\hat{A}_3\hat{A}_+$, $\hat{G}_{21}\hat{B}_3\hat{A}_+$, $\hat{G}_{21}\partial\hat{A}_+$, $T_+^{(\frac{5}{2})}\hat{A}_+$, $U^{(\frac{5}{2})}\hat{A}_3$, $U^{(\frac{5}{2})}\hat{B}_3$, $\partial U^{(\frac{5}{2})}$, $\partial^2\hat{G}_{11}$
$-\frac{(-4+k)}{(6+k)}$	$V^{(\frac{7}{2})}$, $\hat{A}_3V^{(\frac{5}{2})}$, $\hat{A}_3\hat{A}_3\hat{G}_{22}$, $\hat{A}_3\hat{B}_3\hat{G}_{22}$, $\hat{A}_3\partial\hat{G}_{22}$, $\hat{A}_3T_-^{(\frac{5}{2})}$, $\hat{A}_-\hat{A}_3\hat{G}_{12}$, $\hat{A}_-\hat{A}_+\hat{G}_{22}$, $\hat{A}_-\hat{B}_3\hat{G}_{12}$, $\hat{A}_-\partial\hat{G}_{12}$, $\hat{B}_3V^{(\frac{5}{2})}$, $\hat{B}_3\hat{B}_3\hat{G}_{22}$, $\hat{B}_3\partial\hat{G}_{22}$, $\hat{B}_-\hat{G}_{22}\hat{B}_+$, $\hat{B}_+\partial\hat{G}_{21}$, $\hat{G}_{11}\hat{A}_-\hat{B}_+$, $\hat{G}_{12}\partial\hat{A}_-$, $\hat{G}_{21}\hat{A}_3\hat{B}_+$, $\hat{G}_{21}\hat{B}_3\hat{B}_+$, $\hat{G}_{21}\partial\hat{B}_+$, $\hat{G}_{22}\partial\hat{A}_3$, $\hat{G}_{22}\partial\hat{B}_3$, $T^{(2)}\hat{G}_{22}$, $T_+^{(\frac{5}{2})}\hat{B}_+$, $\hat{T}\hat{G}_{22}$, $\partial V^{(\frac{5}{2})}$, $\partial^2\hat{G}_{22}$
$-\frac{(4+k)}{(6+k)}$	$W_-^{(\frac{7}{2})}$, $\hat{A}_3T_-^{(\frac{5}{2})}$, $\hat{A}_3\hat{A}_3\hat{G}_{12}$, $\hat{A}_3\hat{A}_+\hat{G}_{22}$, $\hat{A}_3\hat{B}_3\hat{G}_{12}$, $\hat{A}_3\partial\hat{G}_{12}$, $\hat{A}_-\hat{A}_+\hat{G}_{12}$, $\hat{A}_+V^{(\frac{5}{2})}$, $\hat{A}_+\partial\hat{G}_{22}$, $\hat{B}_3T_-^{(\frac{5}{2})}$, $\hat{B}_3\hat{A}_+\hat{G}_{22}$, $\hat{B}_3\hat{B}_3\hat{G}_{12}$, $\hat{B}_3\partial\hat{G}_{12}$, $\hat{B}_-\hat{G}_{12}\hat{B}_+$, $\hat{B}_+\partial\hat{G}_{11}$, $\hat{G}_{11}\hat{A}_3\hat{B}_+$, $\hat{G}_{11}\hat{B}_3\hat{B}_+$, $\hat{G}_{11}\partial\hat{B}_+$, $\hat{G}_{12}\partial\hat{A}_3$, $\hat{G}_{12}\partial\hat{B}_3$, $\hat{G}_{21}\hat{A}_+\hat{B}_+$, $\hat{G}_{22}\partial\hat{A}_+$, $T^{(2)}\hat{G}_{12}$, $\hat{T}\hat{G}_{12}$, $U^{(\frac{5}{2})}\partial\hat{B}_+$, $\partial T_-^{(\frac{5}{2})}$, $\partial^2\hat{G}_{12}$
$-\frac{(-12+k)}{(6+k)}$	$\hat{A}_-V^{(\frac{5}{2})}$, $\hat{A}_-\hat{A}_3\hat{G}_{22}$, $\hat{A}_-\hat{A}_-\hat{G}_{12}$, $\hat{A}_-\hat{B}_3\hat{G}_{22}$, $\hat{A}_-\partial\hat{G}_{22}$, $\hat{G}_{21}\hat{A}_-\hat{B}_+$, $\hat{G}_{22}\partial\hat{A}_-$
$-\frac{(12+k)}{(6+k)}$	$\hat{A}_3\hat{A}_+\hat{G}_{12}$, $\hat{A}_+T_-^{(\frac{5}{2})}$, $\hat{A}_+\hat{A}_+\hat{G}_{22}$, $\hat{A}_+\partial\hat{G}_{12}$, $\hat{B}_3\hat{A}_+\hat{G}_{12}$, $\hat{G}_{11}\hat{A}_+\hat{B}_+$, $\hat{G}_{12}\partial\hat{A}_+$
$-\frac{(-4+3k)}{(6+k)}$	$\hat{A}_3\hat{G}_{22}\hat{B}_+$, $\hat{A}_-\hat{G}_{12}\hat{B}_+$, $\hat{B}_3\hat{G}_{22}\hat{B}_+$, $\hat{B}_+\partial\hat{G}_{22}$, $\hat{G}_{21}\hat{B}_+\hat{B}_+$, $\hat{G}_{22}\partial\hat{B}_+$, $V^{(\frac{5}{2})}\hat{B}_+$
$-\frac{(4+3k)}{(6+k)}$	$\hat{A}_3\hat{G}_{12}\hat{B}_+$, $\hat{A}_+\hat{G}_{22}\hat{B}_+$, $\hat{B}_3\hat{G}_{12}\hat{B}_+$, $\hat{B}_+\partial\hat{G}_{12}$, $\hat{G}_{11}\hat{B}_+\hat{B}_+$, $\hat{G}_{12}\partial\hat{B}_+$, $T_-^{(\frac{5}{2})}\hat{B}_+$

Table 6: The $U(1)$ charges for the spin-$\frac{7}{2}$ composite fields.

$U(1)$ charge	Composite fields of spin-4
$\frac{2k}{(6+k)}$	$\hat{A}_3\hat{A}_3\partial\hat{B}_-, \hat{A}_3\hat{B}_3\partial\hat{B}_-, \hat{A}_3\partial^2\hat{B}_-, \hat{A}_-\hat{A}_+\partial\hat{B}_-, \hat{B}_3\hat{B}_3\partial\hat{B}_-, \hat{B}_3\partial^2\hat{B}_-, \hat{B}_-T^{(3)}, \hat{B}_-W^{(3)}, \hat{B}_-\hat{A}_3T^{(2)}, \hat{B}_-\hat{A}_3\hat{T}, \hat{B}_-\hat{A}_3\hat{A}_3\hat{A}_3, \hat{B}_-\hat{A}_3\hat{A}_3\hat{B}_3, \hat{B}_-\hat{A}_3\hat{B}_3\hat{B}_3, \hat{B}_-\hat{A}_3\partial\hat{A}_3, \hat{B}_-\hat{A}_3\partial\hat{B}_3, \hat{B}_-\hat{A}_-\hat{A}_3\hat{A}_+, \hat{B}_-\hat{A}_-\hat{B}_3\hat{A}_+, \hat{B}_-\hat{A}_-\partial\hat{A}_+, \hat{B}_-\hat{A}_+\partial\hat{A}_-, \hat{B}_-\hat{B}_3T^{(2)}, \hat{B}_-\hat{B}_3\hat{T}, \hat{B}_-\hat{B}_3\hat{B}_3\hat{B}_3, \hat{B}_-\hat{B}_3\partial\hat{A}_3, \hat{B}_-\hat{B}_3\partial\hat{B}_3, \hat{B}_-\hat{B}_-\hat{A}_3\hat{B}_+, \hat{B}_-\hat{B}_-\hat{B}_3\hat{B}_+, \hat{B}_-\hat{B}_-\partial\hat{B}_+, \hat{B}_-\hat{B}_+\partial\hat{B}_-, \hat{B}_-\hat{G}_{11}\hat{G}_{22}, \hat{B}_-\hat{G}_{21}\hat{G}_{12}, \hat{B}_-\partial T^{(2)}, \hat{B}_-\partial\hat{T}, \hat{B}_-\partial^2\hat{A}_3, \hat{B}_-\partial^2\hat{B}_3, \hat{G}_{11}\hat{G}_{11}\hat{A}_-, \hat{G}_{11}\partial\hat{G}_{21}, \hat{G}_{21}U^{(\frac{5}{2})}, \hat{G}_{21}\hat{G}_{11}\hat{A}_3, \hat{G}_{21}\hat{G}_{11}\hat{B}_3, \hat{G}_{21}\hat{G}_{21}\hat{A}_+, \hat{G}_{21}\partial\hat{G}_{11}, T^{(2)}\partial\hat{B}_-, T_+^{(\frac{5}{2})}\hat{G}_{11}, \hat{T}\partial\hat{B}_-, U_+^{(3)}\hat{A}_3, U_+^{(3)}\hat{B}_3, \partial\hat{B}_-\partial\hat{A}_3, \partial\hat{B}_-\partial\hat{B}_3, \partial U_+^{(3)}, \partial^3\hat{B}_-$
$\frac{2(4+k)}{(6+k)}$	$\hat{A}_-\hat{A}_3\partial\hat{B}_-, \hat{A}_-\hat{B}_3\partial\hat{B}_-, \hat{A}_-\partial^2\hat{B}_-, \hat{B}_-V_+^{(3)}, \hat{B}_-\hat{A}_3\partial\hat{A}_-, \hat{B}_-\hat{A}_-T^{(2)}, \hat{B}_-\hat{A}_-\hat{T}, \hat{B}_-\hat{A}_-\hat{A}_3\hat{A}_3, \hat{B}_-\hat{A}_-\hat{A}_3\hat{B}_3, \hat{B}_-\hat{A}_-\hat{A}_-\hat{A}_+, \hat{B}_-\hat{A}_-\hat{B}_3\hat{B}_3, \hat{B}_-\hat{A}_-\partial\hat{A}_3, \hat{B}_-\hat{A}_-\partial\hat{B}_3, \hat{B}_-\hat{B}_3\partial\hat{A}_-, \hat{B}_-\hat{B}_-\hat{A}_-\hat{B}_+, \hat{B}_-\hat{G}_{21}\hat{G}_{22}, \hat{B}_-\partial^2\hat{A}_-, \hat{G}_{21}T_+^{(\frac{5}{2})}, \hat{G}_{21}\hat{G}_{11}\hat{A}_-, \hat{G}_{21}\hat{G}_{21}\hat{A}_3, \hat{G}_{21}\hat{G}_{21}\hat{B}_3, \hat{G}_{21}\partial\hat{G}_{21}, U_+^{(3)}\hat{A}_-, \partial\hat{B}_-\partial\hat{A}_-,$
$\frac{2(-4+k)}{(6+k)}$	$\hat{A}_3\hat{A}_+\partial\hat{B}_-, \hat{A}_+\partial^2\hat{B}_-, \hat{B}_3\hat{A}_+\partial\hat{B}_-, \hat{B}_-U_-^{(3)}, \hat{B}_-\hat{A}_3\hat{A}_3\hat{A}_+, \hat{B}_-\hat{A}_3\hat{B}_3\hat{A}_+, \hat{B}_-\hat{A}_3\partial\hat{A}_+, \hat{B}_-\hat{A}_-\hat{A}_+\hat{A}_+, \hat{B}_-\hat{A}_+\partial\hat{A}_3, \hat{B}_-\hat{A}_+\partial\hat{B}_3, \hat{B}_-\hat{B}_3\hat{B}_3\hat{A}_+, \hat{B}_-\hat{B}_3\partial\hat{A}_+, \hat{B}_-\hat{B}_-\hat{A}_+\hat{B}_+, \hat{B}_-\hat{G}_{11}\hat{G}_{12}, \hat{B}_-T^{(2)}\hat{A}_+, \hat{B}_-\hat{T}\hat{A}_+, \hat{B}_-\partial^2\hat{A}_+, \hat{G}_{11}U^{(\frac{5}{2})}, \hat{G}_{11}\hat{G}_{11}\hat{A}_3, \hat{G}_{11}\hat{G}_{11}\hat{B}_3, \hat{G}_{11}\partial\hat{G}_{11}, \hat{G}_{21}\hat{G}_{11}\hat{A}_+, U_+^{(3)}\hat{A}_+, \partial\hat{B}_-\partial\hat{A}_+,$
$\frac{8}{(6+k)}$	$\hat{A}_3\hat{A}_3\partial\hat{A}_-, \hat{A}_3\hat{B}_3\partial\hat{A}_-, \hat{A}_3\partial^2\hat{A}_-, \hat{A}_-T^{(3)}, \hat{A}_-W^{(3)}, \hat{A}_-\hat{A}_3T^{(2)}, \hat{A}_-\hat{A}_3\hat{T}, \hat{A}_-\hat{A}_3\hat{A}_3\hat{A}_3, \hat{A}_-\hat{A}_3\hat{A}_3\hat{B}_3, \hat{A}_-\hat{A}_3\hat{B}_3\hat{B}_3, \hat{A}_-\hat{A}_3\partial\hat{A}_3, \hat{A}_-\hat{A}_3\partial\hat{B}_3, \hat{A}_-\hat{A}_-\hat{A}_3\hat{A}_+, \hat{A}_-\hat{A}_-\hat{B}_3\hat{A}_+, \hat{A}_-\hat{A}_-\partial\hat{A}_+, \hat{A}_-\hat{A}_+\partial\hat{A}_-, \hat{A}_-\hat{B}_3T^{(2)}, \hat{A}_-\hat{B}_3\hat{T}, \hat{A}_-\hat{B}_3\hat{B}_3\hat{B}_3, \hat{A}_-\hat{B}_3\partial\hat{A}_3, \hat{A}_-\hat{B}_3\partial\hat{B}_3, \hat{A}_-\hat{B}_+\partial\hat{B}_-, \hat{A}_-\partial T^{(2)}, \hat{A}_-\partial\hat{T}, \hat{A}_-\partial^2\hat{A}_3, \hat{A}_-\partial^2\hat{B}_3, \hat{B}_3\hat{B}_3\partial\hat{A}_-, \hat{B}_3\partial^2\hat{A}_-, \hat{B}_-\hat{A}_-\hat{A}_3\hat{B}_+, \hat{B}_-\hat{A}_-\hat{B}_3\hat{B}_+, \hat{B}_-\hat{A}_-\partial\hat{B}_+, \hat{B}_-\hat{B}_+\partial\hat{A}_-, \hat{B}_-\hat{G}_{22}\hat{G}_{22}, \hat{G}_{11}\hat{A}_-\hat{G}_{22}, \hat{G}_{21}V^{(\frac{5}{2})}, \hat{G}_{21}\hat{A}_3\hat{G}_{22}, \hat{G}_{21}\hat{A}_-\hat{G}_{12}, \hat{G}_{21}\hat{B}_3\hat{G}_{22}, \hat{G}_{21}\hat{G}_{21}\hat{B}_+, \hat{G}_{21}\partial\hat{G}_{22}, \hat{G}_{22}\partial\hat{G}_{21}, T^{(2)}\partial\hat{A}_-, T_+^{(\frac{5}{2})}\hat{G}_{22}, \hat{T}\partial\hat{A}_-, V_+^{(3)}\hat{A}_3, V_+^{(3)}\hat{B}_3, \partial\hat{A}_-\partial\hat{A}_3, \partial\hat{A}_-\partial\hat{B}_3, \partial V_+^{(3)}, \partial^3\hat{A}_-$
0	$W^{(4)}, \hat{A}_3T^{(3)}, \hat{A}_3W^{(3)}, \hat{A}_3\hat{A}_3T^{(2)}, \hat{A}_3\hat{A}_3\hat{T}, \hat{A}_3\hat{A}_3\hat{A}_3\hat{A}_3, \hat{A}_3\hat{A}_3\hat{A}_3\hat{B}_3, \hat{A}_3\hat{A}_3\hat{B}_3\hat{B}_3, \hat{A}_3\hat{A}_3\partial\hat{A}_3, \hat{A}_3\hat{A}_3\partial\hat{B}_3, \hat{A}_3\hat{A}_+\partial\hat{A}_-, \hat{A}_3\hat{B}_3T^{(2)}, \hat{A}_3\hat{B}_3\hat{T}, \hat{A}_3\hat{B}_3\hat{B}_3\hat{B}_3, \hat{A}_3\hat{B}_3\partial\hat{A}_3, \hat{A}_3\hat{B}_3\partial\hat{B}_3, \hat{A}_3\hat{B}_+\partial\hat{B}_-, \hat{A}_3\partial T^{(2)}, \hat{A}_3\partial\hat{T}, \hat{A}_3\partial^2\hat{A}_3, \hat{A}_3\partial^2\hat{B}_3, \hat{A}_-U_-^{(3)}, \hat{A}_-\hat{A}_3\hat{A}_3\hat{A}_+, \hat{A}_-\hat{A}_3\hat{B}_3\hat{A}_+, \hat{A}_-\hat{A}_3\partial\hat{A}_+, \hat{A}_-\hat{A}_-\hat{A}_+\hat{A}_+, \hat{A}_-\hat{A}_+\partial\hat{A}_3, \hat{A}_-\hat{A}_+\partial\hat{B}_3, \hat{A}_-\hat{B}_3\hat{B}_3\hat{A}_+, \hat{A}_-\hat{B}_3\partial\hat{A}_+, \hat{A}_-T^{(2)}\hat{A}_+, \hat{A}_-\hat{T}\hat{A}_+, \hat{A}_-\partial^2\hat{A}_+, \hat{A}_+\partial^2\hat{A}_-, \hat{B}_3T^{(3)}, \hat{B}_3W^{(3)}, \hat{B}_3\hat{A}_+\partial\hat{A}_-, \hat{B}_3\hat{B}_3T^{(2)}, \hat{B}_3\hat{B}_3\hat{T}, \hat{B}_3\hat{B}_3\hat{B}_3\hat{B}_3, \hat{B}_3\hat{B}_3\partial\hat{A}_3, \hat{B}_3\hat{B}_3\partial\hat{B}_3, \hat{B}_3\hat{B}_+\partial\hat{B}_-, \hat{B}_3\partial T^{(2)}, \hat{B}_3\partial\hat{T}, \hat{B}_3\partial^2\hat{A}_3, \hat{B}_3\partial^2\hat{B}_3, \hat{B}_-V_-^{(3)}, \hat{B}_-\hat{A}_3\hat{A}_3\hat{B}_+, \hat{B}_-\hat{A}_3\hat{B}_3\hat{B}_+, \hat{B}_-\hat{A}_3\partial\hat{B}_+, \hat{B}_-\hat{A}_-\hat{A}_+\hat{B}_+, \hat{B}_-\hat{B}_3\hat{B}_3\hat{B}_+, \hat{B}_-\hat{B}_3\partial\hat{B}_+, \hat{B}_-\hat{B}_-\hat{B}_+\hat{B}_+, \hat{B}_-\hat{B}_+\partial\hat{A}_3, \hat{B}_-\hat{B}_+\partial\hat{B}_3, \hat{B}_-\hat{G}_{22}\hat{G}_{12}, \hat{B}_-T^{(2)}\hat{B}_+, \hat{B}_-\hat{T}\hat{B}_+, \hat{B}_-\partial^2\hat{B}_+, \hat{B}_+\partial^2\hat{B}_-, \hat{G}_{11}V^{(\frac{5}{2})}, \hat{G}_{11}\hat{A}_3\hat{G}_{22}, \hat{G}_{11}\hat{A}_-\hat{G}_{12}, \hat{G}_{11}\hat{B}_3\hat{G}_{22}, \hat{G}_{11}\partial\hat{G}_{22}, \hat{G}_{12}\partial\hat{G}_{21}, \hat{G}_{21}T_-^{(\frac{5}{2})}, \hat{G}_{21}\hat{A}_3\hat{G}_{12}, \hat{G}_{21}\hat{A}_+\hat{G}_{22}, \hat{G}_{21}\hat{B}_3\hat{G}_{12}, \hat{G}_{21}\hat{G}_{11}\hat{B}_+, \hat{G}_{21}\partial\hat{G}_{12}, \hat{G}_{22}\partial\hat{G}_{11}, T^{(2)}T^{(2)}, T^{(2)}\partial\hat{A}_3, T^{(2)}\partial\hat{B}_3, T_+^{(\frac{5}{2})}\hat{G}_{12}, \hat{T}T^{(2)}, \hat{T}\hat{T}, \hat{T}\partial\hat{A}_3, \hat{T}\partial\hat{B}_3, U_+^{(3)}\hat{B}_+, U^{(\frac{5}{2})}\hat{G}_{22}, V_+^{(3)}\hat{A}_+, \partial\hat{A}_3\partial\hat{A}_3, \partial\hat{A}_3\partial\hat{B}_3, \partial\hat{A}_-\partial\hat{A}_+, \partial\hat{B}_3\partial\hat{B}_3, \partial\hat{B}_-\partial\hat{B}_+, \partial T^{(3)}, \partial W^{(3)}, \partial^2T^{(2)}, \partial^2\hat{T}, \partial^3\hat{A}_3, \partial^3\hat{B}_3$

Table 7: The $U(1)$ charges for the spin-4 composite fields.

$U(1)$ charge	Composite fields of spin-4
$-\frac{8}{(8+k)}$	$\hat{A}_3U_-^{(3)}, \hat{A}_3\hat{A}_3\hat{A}_3\hat{A}_+, \hat{A}_3\hat{A}_3\hat{B}_3\hat{A}_+, \hat{A}_3\hat{A}_3\partial\hat{A}_+, \hat{A}_3\hat{A}_+\partial\hat{A}_3, \hat{A}_3\hat{A}_+\partial\hat{B}_3,$ $\hat{A}_3\hat{B}_3\hat{B}_3\hat{A}_+, \hat{A}_3\hat{B}_3\partial\hat{A}_+, \hat{A}_3T^{(2)}\hat{A}_+, \hat{A}_3\hat{T}\hat{A}_+, \hat{A}_3\partial^2\hat{A}_+, \hat{A}_-\hat{A}_3\hat{A}_+\hat{A}_+,$ $\hat{A}_-\hat{A}_+\partial\hat{A}_+, \hat{A}_-\hat{B}_3\hat{A}_+\hat{A}_+, \hat{A}_+\hat{A}_+\partial\hat{A}_-, \hat{A}_+\hat{B}_+\partial\hat{B}_-, \hat{A}_+\partial T^{(2)}, \hat{A}_+\partial\hat{T},$ $\hat{A}_+\partial^2\hat{A}_3, \hat{A}_+\partial^2\hat{B}_3, \hat{B}_3U_-^{(3)}, \hat{B}_3\hat{A}_+\partial\hat{A}_3, \hat{B}_3\hat{A}_+\partial\hat{B}_3, \hat{B}_3\hat{B}_3\hat{B}_3\hat{A}_+, \hat{B}_3\hat{B}_3\partial\hat{A}_+,$ $\hat{B}_3T^{(2)}\hat{A}_+, \hat{B}_3\hat{T}\hat{A}_+, \hat{B}_3\partial^2\hat{A}_+, \hat{B}_-\hat{A}_3\hat{A}_+\hat{B}_+, \hat{B}_-\hat{A}_+\partial\hat{B}_+, \hat{B}_-\hat{B}_3\hat{A}_+\hat{B}_+,$ $\hat{B}_-\hat{B}_+\partial\hat{A}_+, \hat{B}_-\hat{G}_{12}\hat{G}_{12}, \hat{G}_{11}T_-^{(\frac{5}{2})}, \hat{G}_{11}\hat{A}_3\hat{G}_{12}, \hat{G}_{11}\hat{A}_+\hat{G}_{22}, \hat{G}_{11}\hat{B}_3\hat{G}_{12},$ $\hat{G}_{11}\hat{G}_{11}\hat{B}_+, \hat{G}_{11}\partial\hat{G}_{12}, \hat{G}_{12}\partial\hat{G}_{11}, \hat{G}_{21}\hat{A}_+\hat{G}_{12}, T^{(2)}\partial\hat{A}_+, T^{(3)}\hat{A}_+, \hat{T}\partial\hat{A}_+,$ $U^{(\frac{5}{2})}\hat{G}_{12}, W^{(3)}\partial\hat{A}_+, \partial\hat{A}_3\partial\hat{A}_+, \partial\hat{B}_3\partial\hat{A}_+, \partial U_-^{(3)}, \partial^3\hat{A}_+$
$-\frac{2(-4+k)}{(6+k)}$	$\hat{A}_3\hat{B}_+\partial\hat{A}_-, \hat{A}_3\hat{G}_{22}\hat{G}_{22}, \hat{A}_-V_-^{(3)}, \hat{A}_-\hat{A}_3\hat{A}_3\hat{B}_+, \hat{A}_-\hat{A}_3\hat{B}_3\hat{B}_+, \hat{A}_-\hat{A}_3\partial\hat{B}_+,$ $\hat{A}_-\hat{A}_-\hat{A}_+\hat{B}_+, \hat{A}_-\hat{B}_3\hat{B}_3\hat{B}_+, \hat{A}_-\hat{B}_3\partial\hat{B}_+, \hat{A}_-\hat{B}_+\partial\hat{A}_3, \hat{A}_-\hat{B}_+\partial\hat{B}_3, \hat{A}_-\hat{G}_{22}\partial\hat{G}_{12},$ $\hat{A}_-T^{(2)}\hat{B}_+, \hat{A}_-\hat{T}\hat{B}_+, \hat{A}_-\partial^2\hat{B}_+, \hat{B}_3\hat{B}_+\partial\hat{A}_-, \hat{B}_3\hat{G}_{22}\hat{G}_{22}, \hat{B}_-\hat{A}_-\hat{B}_+\hat{B}_+,$ $\hat{B}_+\partial^2\hat{A}_-, \hat{G}_{21}\hat{G}_{22}\hat{B}_+, \hat{G}_{22}V^{(\frac{5}{2})}, \hat{G}_{22}\partial\hat{G}_{22}, V_+^{(3)}\hat{B}_+, \partial\hat{A}_-\partial\hat{B}_+$
$-\frac{2(4+k)}{(6+k)}$	$\hat{A}_3\hat{A}_3\hat{A}_+\hat{B}_+, \hat{A}_3\hat{A}_+\partial\hat{B}_+, \hat{A}_3\hat{B}_3\hat{A}_+\hat{B}_+, \hat{A}_3\hat{B}_+\partial\hat{A}_+, \hat{A}_3\hat{G}_{12}\hat{G}_{12}, \hat{A}_-\hat{A}_+\hat{A}_+\hat{B}_+,$ $\hat{A}_+V_-^{(3)}, \hat{A}_+\hat{B}_+\partial\hat{A}_3, \hat{A}_+\hat{B}_+\partial\hat{B}_3, \hat{A}_+\hat{G}_{22}\hat{G}_{12}, \hat{A}_+\partial^2\hat{B}_+, \hat{B}_3\hat{A}_+\partial\hat{B}_+,$ $\hat{B}_3\hat{B}_3\hat{A}_+\hat{B}_+, \hat{B}_3\hat{B}_+\partial\hat{A}_+, \hat{B}_3\hat{G}_{12}\hat{G}_{12}, \hat{B}_-\hat{A}_+\hat{B}_+\hat{B}_+, \hat{B}_+\partial^2\hat{A}_+, \hat{G}_{11}\hat{G}_{12}\hat{B}_+,$ $\hat{G}_{12}T_-^{(\frac{5}{2})}, \hat{G}_{12}\partial\hat{G}_{12}, T^{(2)}\hat{A}_+\hat{B}_+, \hat{T}\hat{A}_+\hat{B}_+, U_-^{(3)}\hat{B}_+, \partial\hat{A}_+\partial\hat{B}_+,$
$-\frac{2k}{(6+k)}$	$\hat{A}_3V_-^{(3)}, \hat{A}_3\hat{A}_3\hat{A}_3\hat{B}_+, \hat{A}_3\hat{A}_3\hat{B}_3\hat{B}_+, \hat{A}_3\hat{A}_3\partial\hat{B}_+, \hat{A}_3\hat{B}_3\hat{B}_3\hat{B}_+, \hat{A}_3\hat{B}_3\partial\hat{B}_+,$ $\hat{A}_3\hat{B}_+\partial\hat{A}_3, \hat{A}_3\hat{B}_+\partial\hat{B}_3, \hat{A}_3\hat{G}_{22}\hat{G}_{12}, \hat{A}_3T^{(2)}\hat{B}_+, \hat{A}_3\hat{T}\hat{B}_+, \hat{A}_3\partial^2\hat{B}_+,$ $\hat{A}_-\hat{A}_3\hat{A}_+\hat{B}_+, \hat{A}_-\hat{A}_+\partial\hat{B}_+, \hat{A}_-\hat{B}_3\hat{A}_+\hat{B}_+, \hat{A}_-\hat{B}_+\partial\hat{A}_+, \hat{A}_-\hat{G}_{12}\hat{G}_{12}, \hat{A}_+\hat{B}_+\partial\hat{A}_-,$ $\hat{A}_+\hat{G}_{22}\hat{G}_{22}, \hat{B}_3V_-^{(3)}, \hat{B}_3\hat{B}_3\hat{B}_3\hat{B}_+, \hat{B}_3\hat{B}_3\partial\hat{B}_+, \hat{B}_3\hat{B}_+\partial\hat{A}_3, \hat{B}_3\hat{B}_+\partial\hat{B}_3,$ $\hat{B}_3\hat{G}_{22}\hat{G}_{12}, \hat{B}_3T^{(2)}\hat{B}_+, \hat{B}_3\hat{T}\hat{B}_+, \hat{B}_3\partial^2\hat{B}_+, \hat{B}_-\hat{A}_3\hat{B}_+\hat{B}_+, \hat{B}_-\hat{B}_3\hat{B}_+\hat{B}_+,$ $\hat{B}_-\hat{B}_+\partial\hat{B}_+, \hat{B}_+\hat{B}_+\partial\hat{B}_-, \hat{B}_+\partial T^{(2)}, \hat{B}_+\partial\hat{T}, \hat{B}_+\partial^2\hat{A}_3, \hat{B}_+\partial^2\hat{B}_3,$ $\hat{G}_{11}\hat{G}_{22}\hat{B}_+, \hat{G}_{12}\partial\hat{G}_{22}, \hat{G}_{21}\hat{G}_{12}\hat{B}_+, \hat{G}_{22}T_-^{(\frac{5}{2})}, \hat{G}_{22}\partial\hat{G}_{12}, T^{(2)}\partial\hat{B}_+,$ $T^{(3)}\hat{B}_+, \hat{T}\partial\hat{B}_+, V^{(\frac{5}{2})}\hat{G}_{12}, W^{(3)}\hat{B}_+, \partial\hat{A}_3\partial\hat{B}_+, \partial\hat{B}_3\partial\hat{B}_+, \partial V_-^{(3)}, \partial^3\hat{B}_+$

Table 8: The $U(1)$ charges for the spin-4 composite fields.

$U(1)$ charge	Composite fields of spin-$\frac{9}{2}$
$\frac{(4+k)}{(6+k)}$	$\hat{A}_3\hat{A}_3\partial\hat{G}_{21}, \hat{A}_3\hat{B}_3\partial\hat{G}_{21}, \hat{A}_3\hat{G}_{22}\partial\hat{B}_-, \hat{A}_3\partial T_+^{(\frac{5}{2})}, \hat{A}_3\partial^2\hat{G}_{21}, \hat{A}_-\hat{A}_3\partial\hat{G}_{11},$ $\hat{A}_-\hat{A}_+\partial\hat{G}_{21}, \hat{A}_-\hat{B}_3\partial\hat{G}_{11}, \hat{A}_-\hat{G}_{12}\partial\hat{B}_-, \hat{A}_-\partial U^{(\frac{5}{2})}, \hat{A}_-\partial^2\hat{G}_{11}, \hat{B}_3\hat{B}_3\partial\hat{G}_{21},$ $\hat{B}_3\hat{G}_{22}\partial\hat{B}_-, \hat{B}_3\partial T_+^{(\frac{5}{2})}, \hat{B}_3\partial^2\hat{G}_{21}, \hat{B}_-V^{(\frac{7}{2})}, \hat{B}_-\hat{A}_3V^{(\frac{5}{2})}, \hat{B}_-\hat{A}_3\hat{A}_3\hat{G}_{22}, \hat{B}_-\hat{A}_3\hat{B}_3\hat{G}_{22},$ $\hat{B}_-\hat{A}_3\partial\hat{G}_{22}, \hat{B}_-\hat{A}_-T_-^{(\frac{5}{2})}, \hat{B}_-\hat{A}_-\hat{A}_3\hat{G}_{12}, \hat{B}_-\hat{A}_-\hat{A}_+\hat{G}_{22}, \hat{B}_-\hat{A}_-\hat{B}_3\hat{G}_{12},$ $\hat{B}_-\hat{A}_+\partial\hat{G}_{12}, \hat{B}_-\hat{B}_3V^{(\frac{5}{2})}, \hat{B}_-\hat{B}_3\hat{B}_3\hat{G}_{22}, \hat{B}_-\hat{B}_3\partial\hat{G}_{22}, \hat{B}_-\hat{B}_-\hat{G}_{22}\hat{B}_+, \hat{B}_-\hat{B}_+\partial\hat{G}_{21},$ $\hat{B}_-\hat{G}_{11}\hat{A}_-\hat{B}_+, \hat{B}_-\hat{G}_{12}\partial\hat{A}_-, \hat{B}_-\hat{G}_{21}\hat{A}_3\hat{B}_+, \hat{B}_-\hat{G}_{21}\hat{B}_3\hat{B}_+, \hat{B}_-\hat{G}_{21}\partial\hat{B}_+, \hat{B}_-\hat{G}_{22}\partial\hat{A}_3,$ $\hat{B}_-\hat{G}_{22}\partial\hat{B}_3, \hat{B}_-T^{(2)}\hat{G}_{22}, \hat{B}_-T_+^{(\frac{5}{2})}\hat{B}_+, \hat{B}_-\hat{T}\hat{G}_{22}, \hat{B}_-\partial V^{(\frac{5}{2})}, \hat{B}_-\partial^2\hat{G}_{22}, \hat{G}_{11}V_+^{(3)},$ $\hat{G}_{11}\hat{A}_3\partial\hat{A}_-, \hat{G}_{11}\hat{A}_-T^{(2)}, \hat{G}_{11}\hat{A}_-\hat{T}, \hat{G}_{11}\hat{A}_-\hat{A}_3\hat{A}_3, \hat{G}_{11}\hat{A}_-\hat{A}_3\hat{B}_3, \hat{G}_{11}\hat{A}_-\hat{A}_-\hat{A}_+,$ $\hat{G}_{11}\hat{A}_-\hat{B}_3\hat{B}_3, \hat{G}_{11}\hat{A}_-\partial\hat{A}_3, \hat{G}_{11}\hat{A}_-\partial\hat{B}_3, \hat{G}_{11}\hat{B}_3\partial\hat{A}_-, \hat{G}_{11}\partial^2\hat{A}_-, \hat{G}_{21}T^{(3)},$ $\hat{G}_{21}W^{(3)}, \hat{G}_{21}\hat{A}_3T^{(2)}, \hat{G}_{21}\hat{A}_3\hat{T}, \hat{G}_{21}\hat{A}_3\hat{A}_3\hat{A}_3, \hat{G}_{21}\hat{A}_3\hat{A}_3\hat{B}_3, \hat{G}_{21}\hat{A}_3\hat{B}_3\hat{B}_3,$ $\hat{G}_{21}\hat{A}_3\partial\hat{A}_3, \hat{G}_{21}\hat{A}_3\partial\hat{B}_3, \hat{G}_{21}\hat{A}_-\hat{A}_3\hat{A}_+, \hat{G}_{21}\hat{A}_-\hat{B}_3\hat{A}_+, \hat{G}_{21}\hat{A}_-\partial\hat{A}_+, \hat{G}_{21}\hat{A}_+\partial\hat{A}_-,$ $\hat{G}_{21}\hat{B}_3T^{(2)}, \hat{G}_{21}\hat{B}_3\hat{T}, \hat{G}_{21}\hat{B}_3\hat{B}_3\hat{B}_3, \hat{G}_{21}\hat{B}_3\partial\hat{A}_3, \hat{G}_{21}\hat{B}_3\partial\hat{B}_3, \hat{G}_{21}\hat{B}_+\partial\hat{B}_-,$ $\hat{G}_{21}\hat{G}_{11}\hat{G}_{22}, \hat{G}_{21}\hat{G}_{21}\hat{G}_{12}, \hat{G}_{21}\partial T^{(2)}, \hat{G}_{21}\partial\hat{T}, \hat{G}_{21}\partial^2\hat{A}_3, \hat{G}_{21}\partial^2\hat{B}_3, \hat{G}_{22}\partial^2\hat{B}_-,$ $T^{(2)}\partial\hat{G}_{21}, T_+^{(\frac{5}{2})}T^{(2)}, T_+^{(\frac{5}{2})}\hat{T}, T_+^{(\frac{5}{2})}\hat{A}_3\hat{A}_3, T_+^{(\frac{5}{2})}\hat{A}_3\hat{B}_3, T_+^{(\frac{5}{2})}\hat{A}_-\hat{A}_+, T_+^{(\frac{5}{2})}\hat{B}_3\hat{B}_3,$ $T_+^{(\frac{5}{2})}\partial\hat{A}_3, T_+^{(\frac{5}{2})}\partial\hat{B}_3, \hat{T}\partial\hat{G}_{21}, U_+^{(3)}\hat{G}_{22}, U^{(\frac{5}{2})}\hat{A}_-\hat{A}_3, U^{(\frac{5}{2})}\hat{A}_-\hat{B}_3, U^{(\frac{5}{2})}\partial\hat{A}_-,$ $U^{(\frac{7}{2})}\hat{A}_-, V^{(\frac{5}{2})}\partial\hat{B}_-, W_+^{(\frac{7}{2})}\hat{A}_3, W_+^{(\frac{7}{2})}\hat{B}_3, \partial\hat{B}_-\partial\hat{G}_{22}, \partial\hat{G}_{11}\partial\hat{A}_-, \partial\hat{G}_{21}\partial\hat{A}_3,$ $\partial\hat{G}_{21}\partial\hat{B}_3, \partial W_+^{(\frac{7}{2})}, \partial^2T_+^{(\frac{5}{2})}, \partial^3\hat{G}_{21}$
$\frac{(-4+k)}{(6+k)}$	$\hat{A}_3\hat{A}_3\partial\hat{G}_{11}, \hat{A}_3\hat{A}_+\partial\hat{G}_{21}, \hat{A}_3\hat{B}_3\partial\hat{G}_{11}, \hat{A}_3\hat{G}_{12}\partial\hat{B}_-, \hat{A}_3\partial U^{(\frac{5}{2})}, \hat{A}_3\partial^2\hat{G}_{11}, \hat{A}_-\hat{A}_+\partial\hat{G}_{11},$ $\hat{A}_+\hat{G}_{22}\partial\hat{B}_-, \hat{A}_+\partial T_+^{(\frac{5}{2})}, \hat{A}_+\partial^2\hat{G}_{21}, \hat{B}_3\hat{A}_+\partial\hat{G}_{21}, \hat{B}_3\hat{B}_3\partial\hat{G}_{11}, \hat{B}_3\hat{G}_{12}\partial\hat{B}_-,$ $\hat{B}_3\partial U^{(\frac{5}{2})}, \hat{B}_3\partial^2\hat{G}_{11}, \hat{B}_-W_-^{(\frac{7}{2})}, \hat{B}_-\hat{A}_3T_-^{(\frac{5}{2})}, \hat{B}_-\hat{A}_3\hat{A}_3\hat{G}_{12}, \hat{B}_-\hat{A}_3\hat{A}_+\hat{G}_{22},$ $\hat{B}_-\hat{A}_3\hat{B}_3\hat{G}_{12}, \hat{B}_-\hat{A}_3\partial\hat{G}_{12}, \hat{B}_-\hat{A}_-\hat{A}_+\hat{G}_{12}, \hat{B}_-\hat{A}_+V^{(\frac{5}{2})}, \hat{B}_-\hat{A}_+\partial\hat{G}_{22},$ $\hat{B}_-\hat{B}_3T_-^{(\frac{5}{2})}, \hat{B}_-\hat{B}_3\hat{A}_+\hat{G}_{22}, \hat{B}_-\hat{B}_3\hat{B}_3\hat{G}_{12}, \hat{B}_-\hat{B}_3\partial\hat{G}_{12}, \hat{B}_-\hat{B}_-\hat{G}_{12}\hat{B}_+, \hat{B}_-\hat{B}_+\partial\hat{G}_{11},$ $\hat{B}_-\hat{G}_{11}\hat{A}_3\hat{B}_+, \hat{B}_-\hat{G}_{11}\hat{B}_3\hat{B}_+, \hat{B}_-\hat{G}_{11}\partial\hat{B}_+, \hat{B}_-\hat{G}_{12}\partial\hat{A}_3, \hat{B}_-\hat{G}_{12}\partial\hat{B}_3, \hat{B}_-\hat{G}_{21}\hat{A}_+\hat{B}_+,$ $\hat{B}_-\hat{G}_{22}\partial\hat{A}_+, \hat{B}_-T^{(2)}\hat{G}_{12}, \hat{B}_-\hat{T}\hat{G}_{12}, \hat{B}_-U^{(\frac{5}{2})}\hat{B}_+, \hat{B}_-\partial T_-^{(\frac{5}{2})}, \hat{B}_-\partial^2\hat{G}_{12}, \hat{G}_{11}T^{(3)},$ $\hat{G}_{11}W^{(3)}, \hat{G}_{11}\hat{A}_3T^{(2)}, \hat{G}_{11}\hat{A}_3\hat{T}, \hat{G}_{11}\hat{A}_3\hat{A}_3\hat{A}_3, \hat{G}_{11}\hat{A}_3\hat{A}_3\hat{B}_3, \hat{G}_{11}\hat{A}_3\hat{B}_3\hat{B}_3,$ $\hat{G}_{11}\hat{A}_3\partial\hat{A}_3, \hat{G}_{11}\hat{A}_3\partial\hat{B}_3, \hat{G}_{11}\hat{A}_-\hat{A}_3\hat{A}_+, \hat{G}_{11}\hat{A}_-\hat{B}_3\hat{A}_+, \hat{G}_{11}\hat{A}_-\partial\hat{A}_+, \hat{G}_{11}\hat{A}_+\partial\hat{A}_-,$ $\hat{G}_{11}\hat{B}_3T^{(2)}, \hat{G}_{11}\hat{B}_3\hat{T}, \hat{G}_{11}\hat{B}_3\hat{B}_3\hat{B}_3, \hat{G}_{11}\hat{B}_3\partial\hat{A}_3, \hat{G}_{11}\hat{B}_3\partial\hat{B}_3, \hat{G}_{11}\hat{B}_+\partial\hat{B}_-,$ $\hat{G}_{11}\hat{G}_{11}\hat{G}_{22}, \hat{G}_{11}\partial T^{(2)}, \hat{G}_{11}\partial\hat{T}, \hat{G}_{11}\partial^2\hat{A}_3, \hat{G}_{11}\partial^2\hat{B}_3, \hat{G}_{12}\partial^2\hat{B}_-,$ $\hat{G}_{21}U_-^{(3)}, \hat{G}_{21}\hat{A}_3\hat{A}_3\hat{A}_+, \hat{G}_{21}\hat{A}_3\hat{B}_3\hat{A}_+, \hat{G}_{21}\hat{A}_3\partial\hat{A}_+, \hat{G}_{21}\hat{A}_-\hat{A}_+\hat{A}_+, \hat{G}_{21}\hat{A}_+\partial\hat{A}_3,$ $\hat{G}_{21}\hat{A}_+\partial\hat{B}_3, \hat{G}_{21}\hat{B}_3\hat{B}_3\hat{A}_+, \hat{G}_{21}\hat{B}_3\partial\hat{A}_+, \hat{G}_{21}\hat{G}_{11}\hat{G}_{12}, \hat{G}_{21}T^{(2)}\hat{A}_+, \hat{G}_{21}\hat{T}\hat{A}_+, \hat{G}_{21}\partial^2\hat{A}_+,$ $T^{(2)}\partial\hat{G}_{11}, T_-^{(\frac{5}{2})}\partial\hat{B}_-, T_+^{(\frac{5}{2})}\hat{A}_3\hat{A}_+, T_+^{(\frac{5}{2})}\hat{B}_3\hat{A}_+, T_+^{(\frac{5}{2})}\partial\hat{A}_+, \hat{T}\partial\hat{G}_{11}, U_+^{(3)}\hat{G}_{12},$ $U^{(\frac{5}{2})}T^{(2)}, U^{(\frac{5}{2})}\hat{T}, U^{(\frac{5}{2})}\hat{A}_3\hat{A}_3, U^{(\frac{5}{2})}\hat{A}_3\hat{B}_3, U^{(\frac{5}{2})}\hat{A}_-\hat{A}_+, U^{(\frac{5}{2})}\hat{B}_3\hat{B}_3, U^{(\frac{5}{2})}\partial\hat{A}_3,$ $U^{(\frac{5}{2})}\partial\hat{B}_3, U^{(\frac{7}{2})}\hat{A}_3, U^{(\frac{7}{2})}\hat{B}_3, W_+^{(\frac{7}{2})}\hat{A}_+, \partial\hat{B}_-\partial\hat{G}_{12}, \partial\hat{G}_{11}\partial\hat{A}_3,$ $\partial\hat{G}_{11}\partial\hat{B}_3, \partial\hat{G}_{21}\partial\hat{A}_+, \partial U^{(\frac{7}{2})}, \partial^2U^{(\frac{5}{2})}, \partial^3\hat{G}_{11}$

Table 9: The $U(1)$ charges for the spin-$\frac{9}{2}$ composite fields.

$U(1)$ charge	Composite fields of spin-$\frac{9}{2}$
$-\frac{(-4+k)}{(6+k)}$	$\hat{A}_3V^{(\frac{7}{2})}, \hat{A}_3\hat{A}_3V^{(\frac{5}{2})}, \hat{A}_3\hat{A}_3\hat{A}_3\hat{G}_{22}, \hat{A}_3\hat{A}_3\hat{B}_3\hat{G}_{22}, \hat{A}_3\hat{A}_3\partial\hat{G}_{22}, \hat{A}_3\hat{B}_3V^{(\frac{5}{2})}, \hat{A}_3\hat{B}_3\hat{B}_3\hat{G}_{22},$ $\hat{A}_3\hat{B}_3\partial\hat{G}_{22}, \hat{A}_3\hat{B}_+\partial\hat{G}_{21}, \hat{A}_3\hat{G}_{12}\partial\hat{A}_-, \hat{A}_3\hat{G}_{22}\partial\hat{A}_3, \hat{A}_3\hat{G}_{22}\partial\hat{B}_3, \hat{A}_3T^{(2)}\hat{G}_{22},$ $\hat{A}_3\hat{T}\hat{G}_{22}, \hat{A}_3\partial V^{(\frac{5}{2})}, \hat{A}_3\partial^2\hat{G}_{22}, \hat{A}_-W_-^{(\frac{7}{2})}, \hat{A}_-\hat{A}_3T_-^{(\frac{5}{2})}, \hat{A}_-\hat{A}_3\hat{A}_3\hat{G}_{12}, \hat{A}_-\hat{A}_3\hat{A}_+\hat{G}_{22},$ $\hat{A}_-\hat{A}_3\hat{B}_3\hat{G}_{12}, \hat{A}_-\hat{A}_3\partial\hat{G}_{12}, \hat{A}_-\hat{A}_-\hat{A}_+\hat{G}_{12}, \hat{A}_-\hat{A}_+V^{(\frac{5}{2})}, \hat{A}_-\hat{A}_+\partial\hat{G}_{22},$ $\hat{A}_-\hat{B}_3T_-^{(\frac{5}{2})}, \hat{A}_-\hat{B}_3\hat{A}_+\hat{G}_{22}, \hat{A}_-\hat{B}_3\hat{B}_3\hat{G}_{12}, \hat{A}_-\hat{B}_3\partial\hat{G}_{12}, \hat{A}_-\hat{B}_+\partial\hat{G}_{11}, \hat{A}_-\hat{G}_{12}\partial\hat{A}_3,$ $\hat{A}_-\hat{G}_{12}\partial\hat{B}_3, \hat{A}_-\hat{G}_{22}\partial\hat{A}_+, \hat{A}_-T^{(2)}\hat{G}_{12}, \hat{A}_-\hat{T}\hat{G}_{12}, \hat{A}_-\partial T_-^{(\frac{5}{2})}, \hat{A}_-\partial^2\hat{G}_{12},$ $\hat{A}_+\hat{G}_{22}\partial\hat{A}_-, \hat{B}_3V^{(\frac{7}{2})}, \hat{B}_3\hat{B}_3V^{(\frac{5}{2})}, \hat{B}_3\hat{B}_3\hat{B}_3\hat{G}_{22}, \hat{B}_3\hat{B}_3\partial\hat{G}_{22}, \hat{B}_3\hat{B}_+\partial\hat{G}_{21}, \hat{B}_3\hat{G}_{12}\partial\hat{A}_-,$ $\hat{B}_3\hat{G}_{22}\partial\hat{A}_3, \hat{B}_3\hat{G}_{22}\partial\hat{B}_3, \hat{B}_3T^{(2)}\hat{G}_{22}, \hat{B}_3\hat{T}\hat{G}_{22}, \hat{B}_3\partial V^{(\frac{5}{2})}, \hat{B}_3\partial^2\hat{G}_{22},$ $\hat{B}_-\hat{A}_3\hat{G}_{22}\hat{B}_+, \hat{B}_-\hat{A}_-\hat{G}_{12}\hat{B}_+, \hat{B}_-\hat{B}_3\hat{G}_{22}\hat{B}_+, \hat{B}_-\hat{B}_+\partial\hat{G}_{22}, \hat{B}_-\hat{G}_{21}\hat{B}_+\hat{B}_+, \hat{B}_-\hat{G}_{22}\partial\hat{B}_+,$ $\hat{B}_-V^{(\frac{5}{2})}\hat{B}_+, \hat{B}_+\partial T_+^{(\frac{5}{2})}, \hat{B}_+\partial^2\hat{G}_{21}, \hat{G}_{11}\hat{A}_-\hat{A}_3\hat{B}_+, \hat{G}_{11}\hat{A}_-\hat{B}_3\hat{B}_+, \hat{G}_{11}\hat{A}_-\partial\hat{B}_+,$ $\hat{G}_{11}\hat{B}_+\partial\hat{A}_-, \hat{G}_{11}\hat{G}_{22}\hat{G}_{22}, \hat{G}_{11}\partial^2\hat{A}_-, \hat{G}_{21}V_-^{(3)}, \hat{G}_{21}\hat{A}_3\hat{A}_3\hat{B}_+, \hat{G}_{21}\hat{A}_3\hat{B}_3\hat{B}_+,$ $\hat{G}_{21}\hat{A}_3\partial\hat{B}_+, \hat{G}_{21}\hat{A}_-\hat{A}_+\hat{B}_+, \hat{G}_{21}\hat{B}_3\hat{B}_3\hat{B}_+, \hat{G}_{21}\hat{B}_3\partial\hat{B}_+, \hat{G}_{21}\hat{B}_+\partial\hat{A}_3, \hat{G}_{21}\hat{B}_+\partial\hat{B}_3,$ $\hat{G}_{21}\hat{G}_{22}\hat{G}_{12}, \hat{G}_{21}T^{(2)}\hat{B}_+, \hat{G}_{21}\hat{T}\hat{B}_+, \hat{G}_{21}\partial^2\hat{B}_+, \hat{G}_{22}\hat{B}_+\partial\hat{B}_-, \hat{G}_{22}\partial T^{(2)}, \hat{G}_{22}\partial\hat{T}, \hat{G}_{22}\partial^2\hat{A}_3,$ $\hat{G}_{22}\partial^2\hat{B}_3, T^{(2)}V^{(\frac{5}{2})}, T^{(2)}\partial\hat{G}_{22}, T^{(3)}\hat{G}_{22}, T_-^{(\frac{5}{2})}\partial\hat{A}_-, T_+^{(\frac{5}{2})}\hat{A}_3\hat{B}_+, T_+^{(\frac{5}{2})}\hat{B}_3\hat{B}_+,$ $T_+^{(\frac{5}{2})}\partial\hat{B}_+, \hat{T}V^{(\frac{5}{2})}, \hat{T}\partial\hat{G}_{22}, U^{(\frac{5}{2})}\hat{A}_-\hat{B}_+, V_+^{(3)}\hat{G}_{12}, V^{(\frac{5}{2})}\partial\hat{A}_3, V^{(\frac{5}{2})}\partial\hat{B}_3,$ $W^{(3)}\hat{G}_{22}, W_+^{(\frac{7}{2})}\hat{B}_+, \partial\hat{A}_3\partial\hat{G}_{22}, \partial\hat{A}_-\partial\hat{G}_{12}, \partial\hat{B}_3\partial\hat{G}_{22}, \partial\hat{G}_{21}\partial\hat{B}_+, \partial V^{(\frac{7}{2})},$ $\partial^2V^{(\frac{5}{2})}, \partial^3\hat{G}_{22}$
$-\frac{(4+k)}{(6+k)}$	$\hat{A}_3W_-^{(\frac{7}{2})}, \hat{A}_3\hat{A}_3T_-^{(\frac{5}{2})}, \hat{A}_3\hat{A}_3\hat{A}_3\hat{G}_{12}, \hat{A}_3\hat{A}_3\hat{A}_+\hat{G}_{22}, \hat{A}_3\hat{A}_3\hat{B}_3\hat{G}_{12}, \hat{A}_3\hat{A}_3\partial\hat{G}_{12}, \hat{A}_3\hat{A}_+V^{(\frac{5}{2})},$ $\hat{A}_3\hat{A}_+\partial\hat{G}_{22}, \hat{A}_3\hat{B}_3T_-^{(\frac{5}{2})}, \hat{A}_3\hat{B}_3\hat{A}_+\hat{G}_{22}, \hat{A}_3\hat{B}_3\hat{B}_3\hat{G}_{12}, \hat{A}_3\hat{B}_3\partial\hat{G}_{12}, \hat{A}_3\hat{B}_+\partial\hat{G}_{11},$ $\hat{A}_3\hat{G}_{12}\partial\hat{A}_3, \hat{A}_3\hat{G}_{12}\partial\hat{B}_3, \hat{A}_3\hat{G}_{22}\partial\hat{A}_+, \hat{A}_3T^{(2)}\hat{G}_{12}, \hat{A}_3\hat{T}\hat{G}_{12}, \hat{A}_3\partial T_-^{(\frac{5}{2})},$ $\hat{A}_3\partial^2\hat{G}_{12}, \hat{A}_-\hat{A}_3\hat{A}_+\hat{G}_{12}, \hat{A}_-\hat{A}_+T_-^{(\frac{5}{2})}, \hat{A}_-\hat{A}_+\hat{A}_+\hat{G}_{22}, \hat{A}_-\hat{A}_+\partial\hat{G}_{12},$ $\hat{A}_-\hat{B}_3\hat{A}_+\hat{G}_{12}, \hat{A}_-\hat{G}_{12}\partial\hat{A}_+, \hat{A}_+V^{(\frac{7}{2})}, \hat{A}_+\hat{B}_+\partial\hat{G}_{21}, \hat{A}_+\hat{G}_{12}\partial\hat{A}_-, \hat{A}_+\hat{G}_{22}\partial\hat{A}_3,$ $\hat{A}_+\hat{G}_{22}\partial\hat{B}_3, \hat{A}_+\partial V^{(\frac{5}{2})}, \hat{A}_+\partial^2\hat{G}_{22}, \hat{B}_3W_-^{(\frac{7}{2})}, \hat{B}_3\hat{A}_+V^{(\frac{5}{2})}, \hat{B}_3\hat{A}_+\partial\hat{G}_{22}, \hat{B}_3\hat{B}_3T_-^{(\frac{5}{2})},$ $\hat{B}_3\hat{B}_3\hat{A}_+\hat{G}_{22}, \hat{B}_3\hat{B}_3\hat{B}_3\hat{G}_{12}, \hat{B}_3\hat{B}_3\partial\hat{G}_{12}, \hat{B}_3\hat{B}_+\partial\hat{G}_{11}, \hat{B}_3\hat{G}_{12}\partial\hat{A}_3, \hat{B}_3\hat{G}_{12}\partial\hat{B}_3,$ $\hat{B}_3\hat{G}_{22}\partial\hat{A}_+, \hat{B}_3T^{(2)}\hat{G}_{12}, \hat{B}_3\hat{T}\hat{G}_{12}, \hat{B}_3\partial T_-^{(\frac{5}{2})}, \hat{B}_3\partial^2\hat{G}_{12}, \hat{B}_-\hat{A}_3\hat{G}_{12}\hat{B}_+,$ $\hat{B}_-\hat{A}_+\hat{G}_{22}\hat{B}_+, \hat{B}_-\hat{B}_3\hat{G}_{12}\hat{B}_+, \hat{B}_-\hat{B}_+\partial\hat{G}_{12}, \hat{B}_-\hat{G}_{11}\hat{B}_+\hat{B}_+, \hat{B}_-\hat{G}_{12}\partial\hat{B}_+, \hat{B}_-T_-^{(\frac{5}{2})}\hat{B}_+,$ $\hat{B}_+\partial U^{(\frac{5}{2})}, \hat{B}_+\partial^2\hat{G}_{11}, \hat{G}_{11}V_-^{(3)}, \hat{G}_{11}\hat{A}_3\hat{A}_3\hat{B}_+, \hat{G}_{11}\hat{A}_3\hat{B}_3\hat{B}_+, \hat{G}_{11}\hat{A}_3\partial\hat{B}_+,$ $\hat{G}_{11}\hat{A}_-\hat{A}_+\hat{B}_+, \hat{G}_{11}\hat{B}_3\hat{B}_3\hat{B}_+, \hat{G}_{11}\hat{B}_3\partial\hat{B}_+, \hat{G}_{11}\hat{B}_+\partial\hat{A}_3, \hat{G}_{11}\hat{B}_+\partial\hat{B}_3, \hat{G}_{11}\hat{G}_{22}\hat{G}_{12},$ $\hat{G}_{11}T^{(2)}\hat{B}_+, \hat{G}_{11}\hat{T}\hat{B}_+, \hat{G}_{11}\partial^2\hat{B}_+, \hat{G}_{12}\hat{B}_+\partial\hat{B}_-, \hat{G}_{12}\partial T^{(2)}, \hat{G}_{12}\partial\hat{T}, \hat{G}_{12}\partial^2\hat{A}_3,$ $\hat{G}_{12}\partial^2\hat{B}_3, \hat{G}_{21}\hat{A}_3\hat{A}_+\hat{B}_+, \hat{G}_{21}\hat{A}_+\partial\hat{B}_+, \hat{G}_{21}\hat{B}_3\hat{A}_+\hat{B}_+, \hat{G}_{21}\hat{B}_+\partial\hat{A}_+, \hat{G}_{21}\hat{G}_{12}\hat{G}_{12},$ $\hat{G}_{22}\partial^2\hat{A}_+, T^{(2)}T_-^{(\frac{5}{2})}, T^2\hat{A}_+\hat{G}_{22}, T^{(2)}\partial\hat{G}_{12}, T^{(3)}\hat{G}_{12}, T_-^{(\frac{5}{2})}\partial\hat{A}_3, T_-^{(\frac{5}{2})}\partial\hat{B}_3,$ $T_+^{(\frac{5}{2})}\hat{A}_+\hat{B}_+, \hat{T}T_-^{(\frac{5}{2})}, \hat{T}\hat{A}_+\hat{G}_{22}, \hat{T}\partial\hat{G}_{12}, U_-^{(3)}\hat{G}_{22}, U^{(\frac{5}{2})}\hat{A}_3\hat{B}_+, U^{(\frac{5}{2})}\hat{B}_3\hat{B}_+,$ $U^{(\frac{5}{2})}\partial\hat{B}_+, U^{(\frac{7}{2})}\hat{B}_+, V^{(\frac{5}{2})}\partial\hat{A}_+, W^{(3)}\hat{G}_{12}, \partial\hat{A}_3\partial\hat{G}_{12}, \partial\hat{A}_+\partial\hat{G}_{22}, \partial\hat{B}_3\partial\hat{G}_{12},$ $\partial\hat{G}_{11}\partial\hat{B}_+, \partial W_-^{(\frac{7}{2})}, \partial^2T_-^{(\frac{5}{2})}, \partial^3\hat{G}_{12}$

Table 10: The $U(1)$ charges for the spin-$\frac{9}{2}$ composite fields.

values	(A,B,C) indices of $f_{AB}{}^{C}$
1	$(1,8,\bar{11})$, $(1,14,9)$, $(2,7,\bar{11})$, $(2,13,9)$, $(4,7,13)$, $(4,8,14)$ $(4,\bar{11},9)$ $(4,\bar{12},\bar{10})$, $(7,14,6)$, $(8,13,6)$, $(\bar{10},1,13)$ $(\bar{10},2,14)$, $(\bar{11},\bar{10},6)$, $(\bar{12},1,7)$, $(\bar{12},2,8)$ $(\bar{12},9,6)$, $(6,\bar{9},\bar{12})$, $(6,\bar{10},\bar{11})$, $(6,\bar{13},18)$, $(6,\bar{14},7)$, $(7,\bar{1},\bar{12})$ $(8,\bar{2},\bar{12})$ $(9,\bar{11},4)$, $(9,\bar{13},2)$, $(9,\bar{14},1)$, $(\bar{10},\bar{12},4)$ $(\bar{11},\bar{7},2)$, $(\bar{11},\bar{8},1)$, $(\bar{13},\bar{1},\bar{10})$, $(\bar{13},\bar{7},4)$ $(\bar{14},\bar{2},\bar{10})$ $(\bar{14},\bar{8},4)$, $(1,\bar{7},\bar{12})$, $(1,\bar{3},\bar{10})$, $(2,\bar{8},\bar{12})$, $(2,\bar{14},\bar{10})$, $(7,\bar{11},\bar{2})$, $(7,\bar{13},\bar{4})$, $(8,\bar{11},\bar{1})$ $(8,\bar{14},\bar{4})$, $(9,\bar{6},\bar{12})$, $(\bar{10},\bar{6},\bar{11})$, $(\bar{11},\bar{9},\bar{4})$, $(\bar{12},\bar{10},\bar{4})$ $(\bar{13},\bar{6},\bar{8})$, $(\bar{13},\bar{9},\bar{2})$, $(\bar{14},\bar{6},\bar{7})$, $(\bar{14},\bar{9},\bar{1})$, $(\bar{1},\bar{10},\bar{13})$, $(\bar{1},\bar{12},\bar{7})$, $(\bar{2},\bar{10},\bar{14})$, $(\bar{12},\bar{12},\bar{8})$, $(\bar{7},\bar{2},\bar{11})$, $(\bar{7},\bar{4},\bar{13})$ $(\bar{8},\bar{1},\bar{11})$ $(\bar{8},\bar{4},\bar{14})$, $(\bar{9},\bar{12},\bar{6})$, $(\bar{10},\bar{11},\bar{6})$ $(\bar{11},\bar{4},\bar{9})$, $(\bar{12},\bar{4},\bar{10})$, $(\bar{13},\bar{2},\bar{9})$, $(\bar{13},\bar{8},\bar{6})$ $(\bar{14},\bar{1},\bar{9})$ $(\bar{14},\bar{7},\bar{6})$, $(\bar{6},7,\bar{14})$, $(\bar{6},8,\bar{13})$, $(\bar{6},11,\bar{10})$, $(\bar{6},12,\bar{9})$, $(\bar{7},12,\bar{1})$, $(\bar{8},12,\bar{2})$ $(\bar{9},1,\bar{14})$ $(\bar{9},2,\bar{13})$, $(\bar{9},4,\bar{11})$, $(\bar{10},4,\bar{12})$ $(\bar{11},1,\bar{8})$, $(\bar{11},2,\bar{7})$, $(\bar{13},4,\bar{7})$, $(\bar{13},10,\bar{1})$ $(\bar{14},4,\bar{8})$ $(\bar{14},10,\bar{2})$, $(\bar{1},9,14)$, $(\bar{1},11,8)$, $(\bar{2},9,13)$, $(\bar{2},11,7)$, $(\bar{4},9,11)$, $(\bar{4},10,12)$ $(\bar{4},13,7)$ $(\bar{4},14,8)$, $(\bar{7},6,14)$, $(\bar{8},6,13)$ $(\bar{10},13,1)$, $(\bar{10},14,2)$, $(\bar{11},6,10)$, $(\bar{12},6,9)$ $(\bar{12},7,1)$ $(\bar{12},8,2)$
$\frac{1}{\sqrt{2}}$	$(3,7,7)$, $(3,13,13)$ $(5,7,7)$ $(5,8,8)$ $(5,11,11)$, $(5,12,12)$, $(8,3,8)$, $(14,3,14)$, $(7,\bar{7},3)$, $(7,\bar{7},5)$ $(8,\bar{3},8)$ $(8,\bar{8},5)$ $(11,\bar{11},5)$, $(12,\bar{12},5)$, $(13,\bar{13},3)$, $(14,\bar{3},14)$, $(3,\bar{8},\bar{8})$, $(3,\bar{14},\bar{14})$, $(7,\bar{7},\bar{3})$ $(7,\bar{7},\bar{5})$, $(8,\bar{8},\bar{5})$, $(11,\bar{11},\bar{5})$, $(12,\bar{12},\bar{5})$ $(13,\bar{13},\bar{3})$, $(\bar{3},\bar{8},\bar{8})$, $(\bar{3},\bar{14},\bar{14})$ $(\bar{7},\bar{3},\bar{7})$, $(\bar{7},\bar{5},\bar{7})$, $(\bar{8},\bar{5},\bar{8})$, $(\bar{11},\bar{5},\bar{11})$ $(\bar{12},\bar{5},\bar{12})$ $(\bar{13},\bar{3},\bar{13})$, $(\bar{7},3,\bar{7})$, $(\bar{7},5,\bar{7})$ $(\bar{8},5,\bar{8})$, $(\bar{8},8,\bar{3})$, $(\bar{11},5,\bar{11})$, $(\bar{12},5,\bar{12})$ $(\bar{13},3,\bar{13})$ $(\bar{14},14,\bar{3})$, $(\bar{3},7,7)$, $(\bar{3},13,13)$ $(\bar{5},7,7)$, $(\bar{5},8,8)$, $(\bar{5},11,11)$, $(\bar{5},12,12)$ $(\bar{8},8,3)$ $(\bar{14},14,3)$
$\frac{i}{\sqrt{2}}$	$(3,9,9)$, $(3,11,11)$ $(5,9,9)$ $(5,10,10)$ $(5,13,13)$, $(5,14,14)$, $(10,3,10)$, $(12,3,12)$, $(9,\bar{3},9)$, $(9,\bar{5},9)$ $(10,\bar{5},10)$ $(10,\bar{10},3)$ $(11,\bar{3},11)$, $(12,\bar{12},3)$, $(13,\bar{5},13)$, $(14,\bar{5},14)$, $(3,\bar{10},\bar{10})$, $(3,\bar{12},\bar{12})$ $(9,\bar{9},\bar{3})$ $(9,\bar{9},\bar{5})$ $(10,\bar{10},\bar{5})$, $(11,\bar{11},\bar{3})$, $(13,\bar{13},\bar{5})$, $(14,\bar{14},\bar{5})$, $(\bar{3},\bar{9},\bar{9})$, $(\bar{3},\bar{11},\bar{11})$ $(\bar{5},\bar{9},\bar{9})$, $(\bar{5},\bar{10},\bar{10})$, $(\bar{5},\bar{13},\bar{13})$, $(\bar{5},\bar{14},\bar{14})$ $(\bar{10},\bar{3},\bar{10})$ $(\bar{12},\bar{3},\bar{12})$, $(\bar{9},3,\bar{9})$, $(\bar{9},5,\bar{9})$ $(\bar{10},5,\bar{10})$, $(\bar{10},10,\bar{3})$, $(\bar{11},3,\bar{11})$, $(\bar{12},12,\bar{3})$ $(\bar{13},5,\bar{13})$ $(\bar{14},5,\bar{14})$, $(\bar{3},10,10)$, $(\bar{3},12,12)$ $(\bar{9},9,3)$, $(\bar{9},9,5)$, $(\bar{10},10,5)$, $(\bar{11},11,3)$ $(\bar{13},13,5)$ $(\bar{14},14,5)$
$\frac{(1+i)}{\sqrt{2}}$	$(3,1,1)$, $(5,6,6)$, $(2,\bar{3},2)$, $(4,\bar{5},4)$, $(1,\bar{1},\bar{3})$, $(6,\bar{6},\bar{5})$, $(\bar{3},\bar{2},\bar{2})$, $(\bar{5},\bar{4},\bar{4})$, $(\bar{1},3,\bar{1})$, $(\bar{6},5,\bar{6})$, $(\bar{2},2,3)$, $(\bar{4},4,5)$
$\frac{(1-i)}{\sqrt{2}}$	$(2,3,2)$, $(4,5,4)$, $(1,\bar{1},3)$, $(6,\bar{6},5)$, $(3,\bar{2},\bar{2})$, $(5,\bar{4},\bar{4})$, $(\bar{1},\bar{3},\bar{1})$, $(\bar{6},\bar{5},\bar{6})$, $(\bar{2},2,\bar{3})$, $(\bar{4},4,\bar{5})$, $(\bar{3},1,1)$, $(\bar{5},6,6)$

Table 11: The nonzero structure constants of $SO(8)$.